



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 09:38 pm BST

PDB ID : 4ITR  
Title : Crystal Structure of IbpAFic2-H3717A in complex with adenylylated Cdc42  
Authors : Xiao, J.; Dixon, J.E.  
Deposited on : 2013-01-18  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div>2%</div> <div>79% 14% • 5%</div> </div>
1	B	316	<div> <div>81% 11% • 6%</div> </div>
2	C	191	<div> <div>82% 10% 8%</div> </div>
2	D	191	<div> <div>2%</div> <div>75% 17% • 7%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8384 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosine monophosphate-protein transferase and cysteine protease IbpA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	0	0
			2355	1483	396	473	3			
1	B	296	Total	C	N	O	S	0	0	0
			2329	1468	392	466	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3717	ALA	HIS	ENGINEERED MUTATION	UNP Q06277
B	3717	ALA	HIS	ENGINEERED MUTATION	UNP Q06277

- Molecule 2 is a protein called Cell division control protein 42 homolog.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	176	Total	C	N	O	S	0	0	0
			1370	883	218	263	6			
2	D	177	Total	C	N	O	S	0	0	0
			1377	888	219	264	6			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula:  $C_{10}H_{14}N_5O_7P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
4	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula:  $C_{10}H_{15}N_5O_{11}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
5	D	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	269	Total	O	0	0
			269	269		

Continued on next page...


*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	280	Total 280	O 280	0	0
7	C	163	Total 163	O 163	0	0
7	D	117	Total 117	O 117	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Adenosine monophosphate-protein transferase and cysteine protease IbpA

Chain A: 





- Molecule 1: Adenosine monophosphate-protein transferase and cysteine protease IbpA

Chain B: 




- Molecule 2: Cell division control protein 42 homolog


Chain C: 



- Molecule 2: Cell division control protein 42 homolog

Chain D: 







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.49Å 91.00Å 190.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.95 – 2.30 45.72 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (44.95-2.30) 99.6 (45.72-2.30)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.66 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, $R_{free}$	0.178 , 0.226 0.180 , 0.227	Depositor DCC
$R_{free}$ test set	2525 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 42.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8384	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, AMP, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/2387	0.42	0/3219
1	B	0.24	0/2361	0.41	0/3185
2	C	0.26	0/1400	0.47	0/1906
2	D	0.25	0/1408	0.47	0/1918
All	All	0.25	0/7556	0.44	0/10228

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2355	0	2390	32	0
1	B	2329	0	2367	25	0
2	C	1370	0	1383	9	0
2	D	1377	0	1390	19	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	1	0
4	C	23	0	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	23	0	12	0	0
5	C	28	0	12	0	0
5	D	28	0	12	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	269	0	0	10	0
7	B	280	0	0	7	0
7	C	163	0	0	2	0
7	D	117	0	0	5	0
All	All	8384	0	7578	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3534:SER:OG	7:B:3974:HOH:O	1.84	0.90
1:A:3655:ASP:OD1	7:A:4162:HOH:O	1.96	0.83
1:A:3586:THR:O	7:A:4016:HOH:O	2.00	0.80
1:A:3622:GLU:OE2	7:A:4139:HOH:O	2.00	0.79
1:A:3591:LYS:HE3	1:A:3591:LYS:H	1.48	0.78
1:A:3773:GLU:OE2	7:A:4026:HOH:O	2.03	0.76
1:B:3489:ASN:HD21	1:B:3491:GLN:HB2	1.52	0.74
1:B:3617:THR:OG1	7:B:4103:HOH:O	2.06	0.72
2:D:31:GLU:OE1	7:D:376:HOH:O	2.08	0.71
2:D:98:VAL:HG11	2:D:149:LEU:HD13	1.74	0.69
1:B:3656:GLU:HG2	1:B:3660:ARG:HG3	1.74	0.68
2:D:95:GLU:OE1	7:D:380:HOH:O	2.13	0.67
2:C:98:VAL:HG11	2:C:149:LEU:HD13	1.77	0.66
1:B:3488:VAL:N	7:B:3932:HOH:O	2.28	0.65
1:B:3584:VAL:HA	1:B:3587:LEU:HD22	1.78	0.65
2:C:96:LYS:NZ	7:C:343:HOH:O	2.26	0.62
1:A:3537:GLU:CD	1:A:3537:GLU:H	2.03	0.60
1:A:3508:ASP:OD2	7:A:4141:HOH:O	2.16	0.60
1:A:3584:VAL:HA	1:A:3587:LEU:HD22	1.87	0.57
2:C:98:VAL:HG13	2:C:99:PRO:HD3	1.87	0.55
1:A:3694:ARG:NH1	7:A:3928:HOH:O	2.42	0.53
2:D:120:ARG:NH2	2:D:156:GLU:OE1	2.42	0.52
1:B:3489:ASN:ND2	1:B:3491:GLN:HB2	2.23	0.52
2:C:93:VAL:O	2:C:98:VAL:HG12	2.10	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3507:ASP:OD1	1:B:3571:ARG:NH1	2.43	0.52
1:B:3592:VAL:HG12	7:B:3905:HOH:O	2.09	0.51
1:A:3593:ILE:H	1:A:3593:ILE:HD13	1.76	0.51
1:A:3757:GLN:HG3	1:A:3762:THR:HG21	1.93	0.51
1:B:3754:ILE:HG13	1:B:3755:ILE:HG23	1.91	0.51
2:C:131:LYS:NZ	7:C:438:HOH:O	2.45	0.50
1:B:3599:LYS:NZ	7:B:4046:HOH:O	2.44	0.49
1:B:3655:ASP:OD2	7:B:4176:HOH:O	2.20	0.49
1:B:3537:GLU:O	1:B:3541:LYS:HG2	2.14	0.48
2:D:147:ARG:NH2	7:D:325:HOH:O	2.43	0.47
2:D:66:ARG:HD2	7:D:401:HOH:O	2.13	0.47
2:D:93:VAL:O	2:D:98:VAL:HG12	2.14	0.47
2:C:120:ARG:NH2	2:C:156:GLU:OE1	2.47	0.47
1:A:3696:LYS:HA	7:A:4046:HOH:O	2.13	0.46
1:A:3781:LYS:HD3	1:A:3781:LYS:HA	1.67	0.46
1:B:3670:LYS:HD2	2:C:32:TYR:CE1	2.50	0.46
1:A:3588:PRO:O	1:A:3741:PRO:HD3	2.16	0.46
1:B:3732:ASN:O	1:B:3736:LEU:HG	2.14	0.46
1:A:3534:SER:HB3	2:D:71:SER:OG	2.16	0.45
2:D:98:VAL:HG13	2:D:99:PRO:HD3	1.98	0.45
1:A:3721:GLU:OE2	7:A:4127:HOH:O	2.21	0.45
1:A:3728:ARG:NH2	1:A:3745:LYS:HG2	2.32	0.44
1:B:3497:LEU:HB2	1:B:3499:LEU:HD13	1.99	0.44
1:A:3488:VAL:N	7:A:4066:HOH:O	2.50	0.44
2:D:67:LEU:HD23	2:D:70:LEU:HD12	1.99	0.44
1:A:3541:LYS:HD3	1:A:3541:LYS:HA	1.76	0.44
1:A:3506:GLU:HG3	1:A:3570:PHE:HB3	1.98	0.44
2:D:37:PHE:HB3	2:D:57:ASP:HB2	2.00	0.44
1:B:3607:LEU:HA	1:B:3607:LEU:HD12	1.84	0.44
3:D:204:SO4:O2	7:D:381:HOH:O	2.21	0.44
2:C:164:GLY:O	2:C:168:VAL:HG23	2.18	0.43
1:A:3695:ALA:HA	1:A:3700:ILE:HD12	1.99	0.43
2:D:83:SER:HB3	2:D:86:SER:HB3	2.01	0.43
1:A:3529:GLU:O	1:A:3533:LYS:HG3	2.19	0.42
1:A:3728:ARG:HH21	1:A:3745:LYS:HG2	1.84	0.42
1:A:3786:ASP:OD2	7:A:4158:HOH:O	2.22	0.42
2:D:46:ILE:HB	2:D:51:TYR:CE2	2.54	0.42
1:B:3695:ALA:HA	1:B:3700:ILE:HD12	2.02	0.42
2:D:61:GLN:O	2:D:68:ARG:NH2	2.53	0.42
1:B:3728:ARG:HH21	1:B:3745:LYS:HG2	1.85	0.42
1:A:3507:ASP:OD1	1:A:3571:ARG:NH1	2.54	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3701:GLU:HA	1:A:3702:PRO:HD3	1.88	0.41
1:A:3710:TYR:HB2	1:A:3731:VAL:HG21	2.02	0.41
1:A:3663:TYR:HA	1:A:3719:PHE:HA	2.02	0.41
1:B:3583:LEU:O	1:B:3587:LEU:HD13	2.20	0.41
1:B:3744:THR:OG1	1:B:3745:LYS:N	2.54	0.41
2:C:49:GLU:HA	2:C:50:PRO:HD3	1.79	0.41
1:A:3563:ARG:NH2	2:D:63:ASP:OD2	2.54	0.41
1:A:3750:PHE:CD1	1:A:3773:GLU:HG2	2.56	0.41
1:B:3586:THR:O	7:B:4128:HOH:O	2.22	0.41
1:B:3670:LYS:HG3	1:B:3671:GLU:HG3	2.03	0.41
1:A:3662:LEU:CD2	1:B:3521:GLU:HG2	2.52	0.40
1:A:3732:ASN:O	1:A:3736:LEU:HG	2.21	0.40
2:D:105:CYS:HB3	2:D:108:THR:OG1	2.21	0.40
2:D:164:GLY:O	2:D:168:VAL:HG23	2.21	0.40
2:D:49:GLU:HA	2:D:50:PRO:HD3	1.88	0.40
1:B:3710:TYR:HB2	1:B:3731:VAL:HG21	2.04	0.40
2:D:98:VAL:CG1	2:D:99:PRO:HD3	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/316 (94%)	294 (99%)	3 (1%)	0	100	100
1	B	294/316 (93%)	290 (99%)	4 (1%)	0	100	100
2	C	174/191 (91%)	170 (98%)	4 (2%)	0	100	100
2	D	175/191 (92%)	170 (97%)	5 (3%)	0	100	100
All	All	940/1014 (93%)	924 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/279 (94%)	251 (95%)	12 (5%)	27	38
1	B	260/279 (93%)	252 (97%)	8 (3%)	40	55
2	C	156/171 (91%)	149 (96%)	7 (4%)	27	39
2	D	157/171 (92%)	150 (96%)	7 (4%)	27	39
All	All	836/900 (93%)	802 (96%)	34 (4%)	30	43

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3488	VAL
1	A	3534	SER
1	A	3541	LYS
1	A	3542	GLN
1	A	3548	GLU
1	A	3553	LEU
1	A	3587	LEU
1	A	3591	LYS
1	A	3592	VAL
1	A	3593	ILE
1	A	3640	LEU
1	A	3643	GLN
1	B	3491	GLN
1	B	3505	LEU
1	B	3553	LEU
1	B	3587	LEU
1	B	3607	LEU
1	B	3640	LEU
1	B	3643	GLN
1	B	3671	GLU
2	C	33	VAL
2	C	67	LEU
2	C	74	GLN
2	C	111	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	C	132	ASN
2	C	141	THR
2	C	145	LEU
2	D	33	VAL
2	D	67	LEU
2	D	79	LEU
2	D	111	LEU
2	D	135	LYS
2	D	145	LEU
2	D	162	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	3489	ASN
2	D	162	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	B	3801	-	4,4,4	0.16	0	6,6,6	0.04	0
3	SO4	D	204	-	4,4,4	0.15	0	6,6,6	0.05	0
5	GDP	C	202	6	24,30,30	1.20	2 (8%)	31,47,47	1.93	8 (25%)
5	GDP	D	202	6	24,30,30	1.19	2 (8%)	31,47,47	1.86	7 (22%)
4	AMP	C	201	2	22,25,25	0.89	1 (4%)	25,38,38	1.21	2 (8%)
3	SO4	A	3801	-	4,4,4	0.16	0	6,6,6	0.06	0
3	SO4	C	204	-	4,4,4	0.16	0	6,6,6	0.06	0
4	AMP	D	201	2	22,25,25	0.91	1 (4%)	25,38,38	1.22	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GDP	C	202	6	-	0/12/32/32	0/3/3/3
5	GDP	D	202	6	-	0/12/32/32	0/3/3/3
4	AMP	C	201	2	-	3/6/26/26	0/3/3/3
4	AMP	D	201	2	-	2/6/26/26	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	202	GDP	C6-C5	4.26	1.48	1.41
5	D	202	GDP	C6-C5	4.21	1.48	1.41
4	D	201	AMP	C5-C4	2.56	1.47	1.40
4	C	201	AMP	C5-C4	2.53	1.47	1.40
5	C	202	GDP	C5-C4	2.40	1.47	1.40
5	D	202	GDP	C5-C4	2.35	1.47	1.40

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	202	GDP	C2-N3-C4	4.89	120.95	115.36
5	D	202	GDP	C2-N3-C4	4.70	120.72	115.36
5	D	202	GDP	C5-C6-N1	-3.99	117.97	123.43
5	C	202	GDP	C5-C6-N1	-3.89	118.11	123.43
5	C	202	GDP	C6-C5-C4	-3.89	117.09	120.80
5	D	202	GDP	C6-N1-C2	3.85	122.05	115.93
5	C	202	GDP	C6-N1-C2	3.81	121.99	115.93
5	D	202	GDP	C6-C5-C4	-3.60	117.36	120.80

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	201	AMP	C4-C5-N7	-3.12	106.15	109.40
5	C	202	GDP	PA-O3A-PB	-3.11	122.15	132.83
4	D	201	AMP	C4-C5-N7	-3.07	106.20	109.40
5	C	202	GDP	N3-C2-N1	-3.05	123.15	127.22
5	D	202	GDP	PA-O3A-PB	-3.00	122.53	132.83
4	D	201	AMP	N3-C2-N1	-2.99	124.00	128.68
5	D	202	GDP	N3-C2-N1	-2.96	123.27	127.22
5	D	202	GDP	C4-C5-N7	-2.94	106.33	109.40
5	C	202	GDP	C4-C5-N7	-2.93	106.35	109.40
4	C	201	AMP	N3-C2-N1	-2.91	124.13	128.68
5	C	202	GDP	C3'-C2'-C1'	2.33	104.48	100.98

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	C	201	AMP	C5'-O5'-P-O1P
4	C	201	AMP	C5'-O5'-P-O2P
4	D	201	AMP	C5'-O5'-P-O1P
4	D	201	AMP	C5'-O5'-P-O2P
4	C	201	AMP	C5'-O5'-P-O3P


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	204	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	299/316 (94%)	-0.31	3 (1%) 82 86	18, 28, 49, 82	0
1	B	296/316 (93%)	-0.29	1 (0%) 94 96	16, 28, 44, 66	0
2	C	176/191 (92%)	-0.43	0 100 100	17, 27, 43, 54	0
2	D	177/191 (92%)	-0.28	4 (2%) 60 67	20, 30, 57, 77	0
All	All	948/1014 (93%)	-0.32	8 (0%) 86 89	16, 28, 48, 82	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3655	ASP	2.9
2	D	49	GLU	2.8
2	D	48	GLY	2.7
1	A	3785	GLU	2.7
1	B	3655	ASP	2.4
2	D	47	GLY	2.3
2	D	45	MET	2.1
1	A	3636	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	MG	C	203	1/1	0.81	0.10	26,26,26,26	0
6	MG	D	203	1/1	0.90	0.12	28,28,28,28	0
4	AMP	D	201	23/23	0.97	0.12	27,31,37,39	0
5	GDP	D	202	28/28	0.98	0.10	15,24,29,31	0
4	AMP	C	201	23/23	0.98	0.10	24,31,39,41	0
3	SO4	A	3801	5/5	0.98	0.09	31,44,51,56	0
3	SO4	D	204	5/5	0.98	0.12	41,54,65,66	0
5	GDP	C	202	28/28	0.98	0.10	16,25,31,34	0
3	SO4	B	3801	5/5	0.99	0.08	29,38,46,46	0
3	SO4	C	204	5/5	0.99	0.13	29,33,40,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around GDP D 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around AMP C 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.