



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 06:26 pm BST

PDB ID : 4IX0
Title : Computational Design of an Unnatural Amino Acid Metalloprotein with Atomic Level Accuracy
Authors : Mills, J.; Bolduc, J.; Khare, S.; Stoddard, B.; Baker, D.
Deposited on : 2013-01-24
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unnatural Amino Acid Mediated Metalloprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	0	2	0
			1999	1269	342	382	6			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	GLU	LYS	ENGINEERED MUTATION	UNP Q06121
A	22	VAL	PHE	ENGINEERED MUTATION	UNP Q06121
A	70	ALA	SER	ENGINEERED MUTATION	UNP Q06121
A	110	MET	LYS	ENGINEERED MUTATION	UNP Q06121
A	133	BP5	ILE	ENGINEERED MUTATION	UNP Q06121
A	161	THR	ASN	ENGINEERED MUTATION	UNP Q06121
A	180	SER	ASN	ENGINEERED MUTATION	UNP Q06121
A	182	GLN	ARG	ENGINEERED MUTATION	UNP Q06121
A	184	ASP	LEU	ENGINEERED MUTATION	UNP Q06121
A	210	ASP	GLU	ENGINEERED MUTATION	UNP Q06121
A	246	LEU	PHE	ENGINEERED MUTATION	UNP Q06121
A	249	GLU	-	EXPRESSION TAG	UNP Q06121
A	250	GLY	-	EXPRESSION TAG	UNP Q06121
A	251	SER	-	EXPRESSION TAG	UNP Q06121
A	252	LEU	-	EXPRESSION TAG	UNP Q06121
A	253	GLU	-	EXPRESSION TAG	UNP Q06121
A	254	HIS	-	EXPRESSION TAG	UNP Q06121
A	255	HIS	-	EXPRESSION TAG	UNP Q06121
A	256	HIS	-	EXPRESSION TAG	UNP Q06121
A	257	HIS	-	EXPRESSION TAG	UNP Q06121
A	258	HIS	-	EXPRESSION TAG	UNP Q06121
A	259	HIS	-	EXPRESSION TAG	UNP Q06121

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ni 1 1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	39	Total O 39 39	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Unnatural Amino Acid Mediated Metalloprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.57Å 62.79Å 92.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.99 – 2.50 35.05 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.9 (51.99-2.50) 96.9 (35.05-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.200 , 0.257 0.196 , 0.250	Depositor DCC
R_{free} test set	526 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	37.9	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2049	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NI, BP5, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.13	4/2012 (0.2%)	1.07	9/2708 (0.3%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	39	GLU	CB-CG	6.33	1.64	1.52
1	A	68	GLU	CD-OE1	5.67	1.31	1.25
1	A	39	GLU	CG-CD	5.25	1.59	1.51
1	A	86	GLU	CB-CG	-5.22	1.42	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	248	LEU	CA-CB-CG	7.13	131.69	115.30
1	A	97	ARG	NE-CZ-NH1	7.07	123.83	120.30
1	A	64	ARG	NE-CZ-NH2	-6.74	116.93	120.30
1	A	19	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	A	171	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	A	97	ARG	NE-CZ-NH2	-5.99	117.31	120.30
1	A	64	ARG	NE-CZ-NH1	5.81	123.20	120.30
1	A	238	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	142	LEU	CA-CB-CG	5.20	127.26	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1999	0	2055	41	0
2	A	1	0	0	0	0
3	A	10	0	0	0	0
4	A	39	0	0	3	0
All	All	2049	0	2055	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:GLU:OE2	1:A:171:ARG:HD2	1.56	1.05
1:A:97:ARG:HG2	1:A:97:ARG:HH11	1.37	0.87
1:A:67:ILE:HD11	1:A:98:LYS:HB3	1.57	0.85
1:A:142:LEU:HD13	1:A:172:ILE:HG12	1.64	0.79
1:A:52:TYR:OH	1:A:54:ARG:NH1	2.20	0.74
1:A:34:ASN:HD22	1:A:34:ASN:H	1.39	0.70
1:A:18:ARG:NH1	4:A:410:HOH:O	2.26	0.68
1:A:182:GLN:HB2	1:A:189:ILE:HD13	1.81	0.63
1:A:67:ILE:HD11	1:A:98:LYS:CB	2.29	0.61
1:A:186:THR:OG1	1:A:188:GLU:HB2	2.04	0.58
1:A:55:LYS:O	1:A:89:PHE:HA	2.04	0.57
1:A:205:VAL:O	1:A:207:LYS:NZ	2.39	0.56
1:A:207:LYS:HE3	4:A:438:HOH:O	2.06	0.56
1:A:55:LYS:HE2	1:A:90:ASN:HD22	1.70	0.55
1:A:22:VAL:HG11	1:A:125:LEU:HD13	1.91	0.53
1:A:160:ILE:HD12	1:A:197:LEU:HD21	1.89	0.52
1:A:25:SER:OG	1:A:27:GLN:NE2	2.43	0.52
1:A:34:ASN:H	1:A:34:ASN:ND2	2.06	0.51
1:A:167[A]:ASP:OD2	1:A:171:ARG:NH2	2.44	0.51
1:A:182:GLN:HB2	1:A:189:ILE:CD1	2.41	0.50
1:A:18:ARG:HH11	1:A:18:ARG:HG3	1.78	0.49
1:A:211:SER:HB2	1:A:213:ILE:HG13	1.95	0.49
1:A:145:LEU:HD13	4:A:413:HOH:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:ARG:CG	1:A:97:ARG:HH11	2.19	0.47
1:A:186:THR:O	1:A:187:LEU:HB2	2.14	0.47
1:A:36:ARG:NH1	1:A:78:VAL:O	2.47	0.46
1:A:24:ALA:HB3	1:A:124:ASN:HB3	1.98	0.46
1:A:100:ALA:HA	1:A:107:ILE:CD1	2.45	0.46
1:A:162:ASP:C	1:A:162:ASP:OD1	2.55	0.45
1:A:146:LEU:HD23	1:A:146:LEU:C	2.38	0.44
1:A:179:ILE:O	1:A:194:GLN:NE2	2.51	0.43
1:A:97:ARG:NH1	1:A:97:ARG:HG2	2.15	0.43
1:A:57:PRO:HD3	1:A:89:PHE:CZ	2.54	0.42
1:A:55:LYS:HE2	1:A:90:ASN:ND2	2.35	0.42
1:A:26:ARG:CZ	1:A:30:ILE:HD13	2.50	0.41
1:A:121:ASP:O	1:A:125:LEU:HD22	2.19	0.41
1:A:74:GLU:HA	1:A:105:ILE:HD13	2.03	0.41
1:A:97:ARG:NH1	1:A:97:ARG:CG	2.83	0.41
1:A:97:ARG:HD3	1:A:125:LEU:HD12	2.03	0.40
1:A:15:LEU:HD11	1:A:87:LYS:HB2	2.04	0.40
1:A:202:PRO:HB2	1:A:204:ASN:ND2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	245/258 (95%)	233 (95%)	12 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	222/232 (96%)	193 (87%)	29 (13%)	4 7

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	GLU
1	A	12	VAL
1	A	25	SER
1	A	31	ILE
1	A	32	SER
1	A	34	ASN
1	A	38	LEU
1	A	44	ASN
1	A	58	SER
1	A	75	ARG
1	A	90	ASN
1	A	96	LEU
1	A	97	ARG
1	A	125	LEU
1	A	142	LEU
1	A	144	SER
1	A	145	LEU
1	A	170	LEU
1	A	182	GLN
1	A	187	LEU
1	A	197	LEU
1	A	204	ASN
1	A	210	ASP
1	A	222	LEU
1	A	225	LEU
1	A	231	LEU
1	A	241	GLU
1	A	242	LYS
1	A	248	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	34	ASN
1	A	41	ASN
1	A	90	ASN
1	A	164	ASN
1	A	182	GLN
1	A	204	ASN
1	A	217	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	BP5	A	133	1,2	17,18,19	1.26	2 (11%)	20,23,25	1.94	6 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	BP5	A	133	1,2	-	3/9/10/12	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	BP5	C4-N1	2.23	1.39	1.34
1	A	133	BP5	C3-N1	2.07	1.39	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	133	BP5	C12-C13-C14	-4.43	103.17	111.47
1	A	133	BP5	C4-N1-C3	3.55	122.21	117.23
1	A	133	BP5	C11-N2-C6	2.86	121.67	117.90
1	A	133	BP5	C7-C6-N2	-2.48	118.56	122.26
1	A	133	BP5	C3-C6-N2	2.45	121.11	116.77
1	A	133	BP5	C5-C4-N1	-2.12	119.97	123.43

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	133	BP5	C2-C3-C6-N2
1	A	133	BP5	N1-C3-C6-N2
1	A	133	BP5	C2-C3-C6-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	302	-	4,4,4	0.21	0	6,6,6	0.65	0
3	SO4	A	303	-	4,4,4	0.29	0	6,6,6	0.51	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	246/258 (95%)	0.04	12 (4%)	29 31	16, 30, 57, 68	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	GLU	3.5
1	A	192[A]	GLU	2.9
1	A	27	GLN	2.6
1	A	157	LEU	2.5
1	A	208	VAL	2.4
1	A	216	ARG	2.2
1	A	248	LEU	2.2
1	A	231	LEU	2.2
1	A	186	THR	2.1
1	A	131	LEU	2.1
1	A	199	SER	2.1
1	A	195	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	BP5	A	133	17/18	0.98	0.14	16,20,26,26	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	302	5/5	0.91	0.18	59,60,63,63	0
3	SO4	A	303	5/5	0.99	0.10	39,39,41,43	0
2	NI	A	301	1/1	0.99	0.04	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.