



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 07:57 PM BST

PDB ID : 6J1A
Title : Photoswitchable fluorescent protein Gamillus, off-state
Authors : Nakashima, R.; Sakurai, K.; shinoda, H.; Matsuda, T.; Nagai, T.
Deposited on : 2018-12-28
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div></div> <div>72%</div> <div>12%</div> <div>15%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2003 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	1	0
			1822	1154	311	346	11			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Cl	0	0
			3	3		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	161	Total	O	0	0
			161	161		

- Molecule 1: Green fluorescent protein

Response	Percentage
Yes	72%
No	12%
Don't know	15%



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	161.54Å 161.54Å 161.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	114.23 – 1.96 80.77 – 1.96	Depositor EDS
% Data completeness (in resolution range)	99.8 (114.23-1.96) 99.8 (80.77-1.96)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.00 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.152 , 0.180 0.163 , 0.187	Depositor DCC
R_{free} test set	2492 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2003	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, QYG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.67	16/1843 (0.9%)	1.43	17/2495 (0.7%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	38	GLU	CD-OE2	8.33	1.34	1.25
1	A	23	GLU	CD-OE2	8.28	1.34	1.25
1	A	238	TYR	CG-CD1	-8.19	1.28	1.39
1	A	104[A]	GLU	CD-OE2	-7.76	1.17	1.25
1	A	104[B]	GLU	CD-OE2	-7.76	1.17	1.25
1	A	23	GLU	CG-CD	6.40	1.61	1.51
1	A	166	SER	CA-CB	6.32	1.62	1.52
1	A	116	SER	CB-OG	-6.12	1.34	1.42
1	A	67	SER	CA-CB	5.69	1.61	1.52
1	A	79	TYR	CE2-CZ	-5.44	1.31	1.38
1	A	238	TYR	CE2-CZ	-5.42	1.31	1.38
1	A	223	GLU	CG-CD	5.41	1.60	1.51
1	A	106	ASP	CG-OD2	-5.31	1.13	1.25
1	A	43	SER	CA-CB	5.23	1.60	1.52
1	A	27	GLU	CG-CD	5.23	1.59	1.51
1	A	166	SER	CB-OG	5.00	1.48	1.42

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	140	MET	CG-SD-CE	-13.50	78.60	100.20
1	A	46	ARG	NE-CZ-NH1	-10.72	114.94	120.30
1	A	46	ARG	NE-CZ-NH2	10.38	125.49	120.30
1	A	134	ASP	CB-CG-OD2	-8.46	110.68	118.30
1	A	98	ASP	CB-CG-OD2	-7.88	111.20	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	98	ASP	CB-CG-OD1	-6.40	112.54	118.30
1	A	99	ARG	NE-CZ-NH2	6.37	123.48	120.30
1	A	164	LEU	CB-CG-CD1	6.36	121.80	111.00
1	A	214	ASP	CB-CG-OD1	5.75	123.47	118.30
1	A	119	ASP	CB-CG-OD1	-5.62	113.25	118.30
1	A	82	ASP	CB-CG-OD1	5.51	123.26	118.30
1	A	98	ASP	OD1-CG-OD2	5.45	133.65	123.30
1	A	183	ASP	CB-CG-OD2	5.37	123.13	118.30
1	A	34	LYS	CD-CE-NZ	-5.34	99.42	111.70
1	A	46	ARG	CD-NE-CZ	-5.22	116.30	123.60
1	A	99	ARG	NE-CZ-NH1	-5.18	117.71	120.30
1	A	27	GLU	OE1-CD-OE2	-5.05	117.25	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1822	0	1751	3	0
2	A	3	0	0	0	0
3	A	12	0	16	0	0
4	A	5	0	0	0	0
5	A	161	0	0	1	0
All	All	2003	0	1767	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:ILE:HG12	5:A:541:HOH:O	1.83	0.77
1:A:164:LEU:HB2	1:A:184:CYS:HB2	1.92	0.52
1:A:211:ASP:OD1	1:A:212:PRO:HD2	2.17	0.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/271 (84%)	221 (97%)	6 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/231 (86%)	191 (96%)	7 (4%)	36	24

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LYS
1	A	47	PHE
1	A	75	MET
1	A	150	ASN
1	A	164	LEU
1	A	193	LYS
1	A	198	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	205	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	QYG	A	70	1	25,25,26	2.61	6 (24%)	31,34,36	4.06	14 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	QYG	A	70	1	-	0/13/32/33	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	70	QYG	CB2-CA2	8.23	1.42	1.35
1	A	70	QYG	C1-N2	5.49	1.40	1.32
1	A	70	QYG	C1-N3	4.55	1.44	1.37
1	A	70	QYG	CA3-N3	-3.62	1.40	1.47
1	A	70	QYG	CG2-CB2	-2.88	1.41	1.46
1	A	70	QYG	CD2-CG2	-2.07	1.35	1.39

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	70	QYG	CA2-C2-N3	12.35	109.21	103.37
1	A	70	QYG	O2-C2-CA2	-9.77	125.47	130.96
1	A	70	QYG	CA1-C1-N3	-8.93	113.21	124.85
1	A	70	QYG	C2-N3-C1	-8.35	103.74	107.97
1	A	70	QYG	CA3-N3-C1	6.10	134.49	127.16
1	A	70	QYG	CA1-C1-N2	3.14	129.88	123.56
1	A	70	QYG	CE1-CD1-CG2	-3.09	117.22	121.25
1	A	70	QYG	CB2-CA2-N2	2.95	132.92	128.83
1	A	70	QYG	N3-C1-N2	2.89	113.45	111.45
1	A	70	QYG	CB2-CA2-C2	-2.85	118.87	122.28
1	A	70	QYG	CB1-CG1-CD3	-2.66	103.57	112.59
1	A	70	QYG	CD1-CG2-CB2	-2.12	114.00	121.22
1	A	70	QYG	CD1-CG2-CD2	2.07	120.71	117.64
1	A	70	QYG	CD1-CE1-CZ	2.05	122.13	119.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	305	-	5,5,5	0.88	0	5,5,5	0.93	0
3	GOL	A	304	-	5,5,5	0.73	0	5,5,5	1.37	1 (20%)
4	PO4	A	306	-	4,4,4	0.66	0	6,6,6	2.39	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	305	-	-	2/4/4/4	-
3	GOL	A	304	-	-	0/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	306	PO4	O3-P-O1	-4.12	95.83	110.89
4	A	306	PO4	O2-P-O1	2.92	121.59	110.89
3	A	304	GOL	C3-C2-C1	-2.88	100.50	111.70

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	305	GOL	C1-C2-C3-O3
3	A	305	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/271 (84%)	-0.53	1 (0%) 92 95	19, 26, 47, 63	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	191	GLY	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	QYG	A	70	24/25	0.98	0.09	18,23,40,47	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	305	6/6	0.87	0.13	56,68,74,77	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	304	6/6	0.94	0.19	36,50,51,55	0
2	CL	A	302	1/1	0.98	0.04	31,31,31,31	0
2	CL	A	303	1/1	0.99	0.06	30,30,30,30	0
2	CL	A	301	1/1	0.99	0.09	41,41,41,41	0
4	PO4	A	306	5/5	1.00	0.08	25,34,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.