



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 02:09 pm BST

PDB ID : 6J1H
Title : Crystal structure of HypX from Aquifex aeolicus, Q15A-R131A-S194A-Q195
A-N306A-R542A variant
Authors : Muraki, N.; Aono, S.
Deposited on : 2018-12-28
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	582	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 90% 7% </div> </div>
1	B	582	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 15%, orange 1%, yellow 1%, green 82%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 15% 82% 14% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9240 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrogenase regulation HoxX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	564	Total	C	N	O	S	0	2	0
			4584	2950	785	831	18			
1	B	562	Total	C	N	O	S	0	19	0
			4568	2937	784	830	17			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP O67224
A	-18	GLY	-	expression tag	UNP O67224
A	-17	SER	-	expression tag	UNP O67224
A	-16	SER	-	expression tag	UNP O67224
A	-15	HIS	-	expression tag	UNP O67224
A	-14	HIS	-	expression tag	UNP O67224
A	-13	HIS	-	expression tag	UNP O67224
A	-12	HIS	-	expression tag	UNP O67224
A	-11	HIS	-	expression tag	UNP O67224
A	-10	HIS	-	expression tag	UNP O67224
A	-9	SER	-	expression tag	UNP O67224
A	-8	SER	-	expression tag	UNP O67224
A	-7	GLY	-	expression tag	UNP O67224
A	-6	LEU	-	expression tag	UNP O67224
A	-5	VAL	-	expression tag	UNP O67224
A	-4	PRO	-	expression tag	UNP O67224
A	-3	ARG	-	expression tag	UNP O67224
A	-2	GLY	-	expression tag	UNP O67224
A	-1	SER	-	expression tag	UNP O67224
A	0	HIS	-	expression tag	UNP O67224
A	15	ALA	GLN	engineered mutation	UNP O67224
A	131	ALA	ARG	engineered mutation	UNP O67224
A	194	ALA	SER	engineered mutation	UNP O67224
A	195	ALA	GLN	engineered mutation	UNP O67224
A	306	ALA	ASN	engineered mutation	UNP O67224

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	542	ALA	ARG	engineered mutation	UNP O67224
B	-19	MET	-	initiating methionine	UNP O67224
B	-18	GLY	-	expression tag	UNP O67224
B	-17	SER	-	expression tag	UNP O67224
B	-16	SER	-	expression tag	UNP O67224
B	-15	HIS	-	expression tag	UNP O67224
B	-14	HIS	-	expression tag	UNP O67224
B	-13	HIS	-	expression tag	UNP O67224
B	-12	HIS	-	expression tag	UNP O67224
B	-11	HIS	-	expression tag	UNP O67224
B	-10	HIS	-	expression tag	UNP O67224
B	-9	SER	-	expression tag	UNP O67224
B	-8	SER	-	expression tag	UNP O67224
B	-7	GLY	-	expression tag	UNP O67224
B	-6	LEU	-	expression tag	UNP O67224
B	-5	VAL	-	expression tag	UNP O67224
B	-4	PRO	-	expression tag	UNP O67224
B	-3	ARG	-	expression tag	UNP O67224
B	-2	GLY	-	expression tag	UNP O67224
B	-1	SER	-	expression tag	UNP O67224
B	0	HIS	-	expression tag	UNP O67224
B	15	ALA	GLN	engineered mutation	UNP O67224
B	131	ALA	ARG	engineered mutation	UNP O67224
B	194	ALA	SER	engineered mutation	UNP O67224
B	195	ALA	GLN	engineered mutation	UNP O67224
B	306	ALA	ASN	engineered mutation	UNP O67224
B	542	ALA	ARG	engineered mutation	UNP O67224

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	49	Total	O	0	0
			49	49		
3	B	27	Total	O	0	0
			27	27		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Hydrogenase regulation HoxX



• Molecule 1: Hydrogenase regulation HoxX



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	79.92Å 124.41Å 290.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.25 – 2.10 48.43 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.25-2.10) 94.0 (48.43-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 2.10Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.200 , 0.235 0.201 , 0.237	Depositor DCC
R_{free} test set	1998 reflections (2.37%)	wwPDB-VP
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.346	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9240	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/4707	0.55	0/6360
1	B	14.86	4/4701 (0.1%)	1.98	8/6355 (0.1%)
All	All	10.51	4/9408 (0.0%)	1.45	8/12715 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	546[A]	PHE	CB-CG	542.07	10.72	1.51
1	B	546[B]	PHE	CB-CG	542.07	10.72	1.51
1	B	547[A]	ARG	CB-CG	474.23	14.32	1.52
1	B	547[B]	ARG	CB-CG	474.23	14.32	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	546[A]	PHE	CB-CG-CD1	-102.69	48.92	120.80
1	B	546[B]	PHE	CB-CG-CD1	-102.69	48.92	120.80
1	B	547[A]	ARG	CA-CB-CG	-26.67	54.72	113.40
1	B	547[B]	ARG	CA-CB-CG	-26.67	54.72	113.40
1	B	546[A]	PHE	CB-CG-CD2	-17.49	108.55	120.80
1	B	546[B]	PHE	CB-CG-CD2	-17.49	108.55	120.80
1	B	546[A]	PHE	CA-CB-CG	-7.71	95.41	113.90
1	B	546[B]	PHE	CA-CB-CG	-7.71	95.41	113.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4584	0	4526	20	0
1	B	4568	0	4465	50	0
2	A	6	0	8	0	0
2	B	6	0	8	1	0
3	A	49	0	0	0	0
3	B	27	0	0	0	0
All	All	9240	0	9007	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:5:PHE:HB2	1:B:30:VAL:HG22	1.65	0.77
1:B:543[A]:LYS:HD3	1:B:544[A]:PRO:HD2	1.68	0.74
1:B:253:GLU:O	1:B:541:ARG:NH2	2.22	0.72
1:B:448:ARG:NH1	1:B:546[B]:PHE:O	2.25	0.63
1:B:330:LYS:HD2	1:B:485:ILE:HD12	1.81	0.63
1:A:54:PRO:HA	1:A:73:ILE:HD13	1.85	0.59
1:B:308:ALA:HB1	1:B:346:ASN:HA	1.84	0.58
1:A:36:PRO:HB3	1:A:61:PRO:HG3	1.84	0.58
1:B:113:VAL:HG22	1:B:157:THR:O	2.04	0.57
1:B:54:PRO:HA	1:B:73:ILE:HD13	1.88	0.56
1:B:3:ILE:HD12	1:B:21:LEU:HD13	1.89	0.55
1:A:8:TYR:CE1	1:A:57:LYS:HD2	2.42	0.55
1:B:9:ARG:HB3	1:B:547[B]:ARG:NH2	2.20	0.55
1:B:351:GLU:HG2	1:B:540:VAL:HG21	1.90	0.54
1:B:53:ALA:HB1	1:B:56:LEU:HG	1.90	0.54
1:B:103:GLU:HG3	1:B:114:TRP:HE1	1.74	0.53
1:B:39:THR:HG22	1:B:64:VAL:HG21	1.91	0.52
1:B:69:LYS:HG2	1:B:149:PHE:HE1	1.74	0.52
1:A:497:LYS:HG2	1:A:501:ARG:HG3	1.93	0.51
1:A:253:GLU:O	1:A:541:ARG:NH1	2.43	0.50
1:B:3:ILE:O	1:B:28:VAL:HA	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:334:LEU:HD11	1:B:374:ILE:HD11	1.94	0.49
1:A:182:THR:HG23	1:A:242:VAL:HG22	1.93	0.49
1:B:18:TYR:CE1	1:B:22:THR:HG21	2.47	0.49
1:B:91:MET:O	1:B:187:ARG:HD2	2.12	0.49
1:A:18:TYR:CZ	1:A:22:THR:HG21	2.47	0.49
1:B:448:ARG:HD2	1:B:551[A]:TYR:CE2	2.48	0.48
1:B:8:TYR:CD2	1:B:9:ARG:HG2	2.48	0.48
1:B:276:LYS:HB2	1:B:279:GLU:HG3	1.96	0.48
1:B:322:LYS:NZ	1:B:373:GLU:OE1	2.47	0.47
1:B:44:GLU:O	1:B:47:LYS:HG3	2.14	0.47
1:B:6:LEU:HD23	1:B:6:LEU:HA	1.74	0.47
1:B:20:GLU:HG2	1:B:139:VAL:HG13	1.95	0.47
1:B:31:GLU:OE1	1:B:555[A]:HIS:NE2	2.44	0.47
1:B:9:ARG:HB3	1:B:547[B]:ARG:HH22	1.79	0.47
1:B:130:TYR:HA	1:B:134:VAL:HB	1.97	0.47
1:B:10:PHE:O	1:B:547[B]:ARG:NH1	2.28	0.46
1:B:56:LEU:HA	1:B:56:LEU:HD23	1.79	0.46
1:A:53:ALA:HB1	1:A:56:LEU:HD13	1.96	0.46
1:B:8:TYR:HA	1:B:33:ASP:HB3	1.98	0.46
1:A:308:ALA:HB1	1:A:346:ASN:HA	1.97	0.45
1:A:267:LEU:O	1:A:270:VAL:HG12	2.16	0.45
1:B:216:GLU:HG3	1:B:284:LYS:HB2	1.97	0.45
1:A:395:VAL:HG11	1:A:414:PRO:HB3	1.99	0.44
1:B:314:CYS:SG	1:B:367:ILE:HG12	2.57	0.44
1:B:395:VAL:HG11	1:B:414:PRO:HB3	2.00	0.44
1:A:501:ARG:HD3	1:A:506:TRP:CZ2	2.52	0.44
1:A:334:LEU:HD11	1:A:374:ILE:HD11	1.99	0.44
1:B:491:TYR:O	1:B:495:GLU:HG2	2.17	0.44
1:B:555[A]:HIS:HA	1:B:560[A]:PHE:CD1	2.53	0.44
1:A:193:ASP:OD1	1:A:194:ALA:HA	2.18	0.44
1:B:538:TYR:HD2	1:B:546[B]:PHE:CE1	2.36	0.43
1:B:299:PHE:CG	2:B:701:GOL:H12	2.53	0.43
1:A:103:GLU:O	1:A:111:GLY:HA3	2.19	0.43
1:B:103:GLU:HG3	1:B:114:TRP:NE1	2.34	0.43
1:A:221:LYS:HE3	1:A:221:LYS:HB2	1.74	0.42
1:B:52:ILE:O	1:B:54:PRO:HD3	2.20	0.42
1:A:212:ALA:HA	1:A:243:TRP:O	2.20	0.42
1:B:4:LEU:HD12	1:B:29:SER:O	2.20	0.42
1:B:371:CYS:HB3	1:B:400:THR:HG21	2.02	0.41
1:B:506:TRP:CE3	1:B:507:LEU:HD13	2.54	0.41
1:B:201:LYS:HB2	1:B:201:LYS:HE3	1.83	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:212:ALA:HA	1:B:243:TRP:O	2.19	0.41
1:B:516:HIS:CE1	1:B:520:LYS:HE2	2.55	0.41
1:B:56:LEU:HD11	1:B:60:ILE:HD11	2.01	0.41
1:B:309:MET:HE2	1:B:309:MET:HB3	1.47	0.41
1:A:130:TYR:HA	1:A:134:VAL:HB	2.01	0.41
1:A:314:CYS:HB2	1:A:366:ALA:HB1	2.04	0.40
1:B:38:LEU:HA	1:B:38:LEU:HD12	1.85	0.40
1:A:112:ASP:HB3	1:A:156:PRO:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	564/582 (97%)	553 (98%)	11 (2%)	0	100	100
1	B	565/582 (97%)	535 (95%)	27 (5%)	3 (0%)	29	26
All	All	1129/1164 (97%)	1088 (96%)	38 (3%)	3 (0%)	47	41

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	268	LYS
1	B	545[A]	HIS
1	B	545[B]	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	480/505 (95%)	475 (99%)	5 (1%)	76	82
1	B	474/505 (94%)	464 (98%)	10 (2%)	53	59
All	All	954/1010 (94%)	939 (98%)	15 (2%)	67	69

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	LEU
1	A	45	LEU
1	A	108	TYR
1	A	363	ASN
1	A	558	LEU
1	B	63	GLU
1	B	150	GLU
1	B	363	ASN
1	B	419	ILE
1	B	448	ARG
1	B	507	LEU
1	B	546[A]	PHE
1	B	546[B]	PHE
1	B	547[A]	ARG
1	B	547[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	701	-	5,5,5	1.10	0	5,5,5	0.88	0
2	GOL	A	701	-	5,5,5	1.07	0	5,5,5	1.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	701	-	-	0/4/4/4	-
2	GOL	A	701	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	GOL	O1-C1-C2-O2
2	A	701	GOL	O1-C1-C2-C3
2	A	701	GOL	O2-C2-C3-O3
2	A	701	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	701	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	564/582 (96%)	-0.16	5 (0%) 84 86	35, 47, 71, 95	0
1	B	562/582 (96%)	0.76	87 (15%) 2 2	37, 62, 127, 157	14 (2%)
All	All	1126/1164 (96%)	0.30	92 (8%) 11 15	35, 52, 117, 157	14 (1%)

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	38	LEU	11.1
1	B	154	PHE	10.1
1	B	149	PHE	9.0
1	B	146	LEU	8.8
1	B	46	TYR	7.8
1	B	51	ILE	7.5
1	B	48	PRO	7.5
1	B	32	LEU	7.3
1	B	557[A]	ARG	6.9
1	B	163	TRP	6.8
1	B	546[A]	PHE	6.7
1	B	45	LEU	6.6
1	B	561[A]	SER	6.5
1	B	64	VAL	6.5
1	B	156	PRO	6.2
1	B	-1	SER	5.9
1	B	70	THR	5.9
1	B	547[A]	ARG	5.7
1	B	58	ARG	5.3
1	B	6	LEU	5.3
1	B	115	ALA	5.2
1	B	2	ARG	5.1
1	B	528	PHE	5.1
1	B	71	LEU	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	548[A]	THR	4.9
1	B	107	GLU	4.7
1	B	50	LEU	4.6
1	B	35	HIS	4.5
1	B	72	ILE	4.3
1	B	114	TRP	4.3
1	B	544[A]	PRO	3.9
1	B	63	GLU	3.8
1	B	105	SER	3.8
1	B	560[A]	PHE	3.8
1	B	68	TYR	3.7
1	B	558[A]	LEU	3.7
1	A	558	LEU	3.7
1	B	102	LEU	3.7
1	B	37	ASP	3.5
1	B	4	LEU	3.5
1	B	33	ASP	3.5
1	B	159	GLN	3.5
1	B	555[A]	HIS	3.5
1	B	150	GLU	3.4
1	B	106	GLU	3.3
1	B	552[A]	LEU	3.3
1	B	164	TRP	3.3
1	B	110	ALA	3.3
1	B	36	PRO	3.2
1	B	549[A]	PRO	3.2
1	B	59	LYS	3.2
1	B	551[A]	TYR	3.1
1	B	34	VAL	3.1
1	A	538	TYR	3.1
1	B	169	GLU	3.0
1	B	103	GLU	3.0
1	B	61	PRO	3.0
1	B	109	ASP	2.9
1	B	62	GLN	2.9
1	B	166	PRO	2.9
1	B	554[A]	ILE	2.8
1	A	108	TYR	2.8
1	B	42	ALA	2.8
1	B	116	TYR	2.8
1	B	172	LEU	2.8
1	B	39	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	56	LEU	2.7
1	B	157	THR	2.7
1	B	30	VAL	2.7
1	B	3	ILE	2.6
1	B	60	ILE	2.6
1	B	171	GLU	2.6
1	B	253	GLU	2.5
1	B	160	LYS	2.5
1	B	0	HIS	2.5
1	B	545[A]	HIS	2.4
1	B	162	HIS	2.3
1	A	153	ASP	2.3
1	B	104	ALA	2.3
1	B	170	GLN	2.3
1	B	152	GLY	2.3
1	B	1	MET	2.3
1	B	108	TYR	2.3
1	B	151	ARG	2.3
1	B	556[A]	ARG	2.3
1	B	26	HIS	2.2
1	A	149	PHE	2.2
1	B	344	GLY	2.2
1	B	306	ALA	2.1
1	B	543[A]	LYS	2.1
1	B	271	LYS	2.1
1	B	27	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	701	6/6	0.82	0.17	61,64,65,66	0
2	GOL	B	701	6/6	0.94	0.12	56,66,72,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.