



## Full wwPDB EM Validation Report ⓘ

Nov 14, 2022 – 03:27 PM JST

PDB ID : 6J3Q  
EMDB ID : EMD-9774  
Title : Capsid structure of a freshwater cyanophage Siphoviridae Mic1  
Authors : Jin, H.; Jiang, Y.L.; Yang, F.; Zhang, J.T.; Li, W.F.; Zhou, K.; Ju, J.; Chen, Y.; Zhou, C.Z.  
Deposited on : 2019-01-05  
Resolution : 3.53 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	<div> <div>12%</div> <div>96%</div> </div>
1	B	348	<div> <div>14%</div> <div>97%</div> </div>
1	C	348	<div> <div>9%</div> <div>98%</div> </div>
1	D	348	<div> <div>10%</div> <div>98%</div> </div>
1	E	348	<div> <div>16%</div> <div>98%</div> </div>
1	F	348	<div> <div>14%</div> <div>97%</div> </div>
1	U	348	<div> <div>11%</div> <div>97%</div> </div>
1	V	348	<div> <div>10%</div> <div>97%</div> </div>
1	W	348	<div> <div>11%</div> <div>97%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	X	348	
1	Y	348	
1	Z	348	
1	a	348	
2	0	99	
2	1	99	
2	2	99	
2	3	99	
2	4	99	
2	5	99	
2	6	99	
2	7	99	
2	8	99	
2	9	99	
2	b	99	
2	c	99	
2	d	99	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 43481 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called major capsid protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	B	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	C	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	D	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	E	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	F	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	U	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	V	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	W	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	X	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	Y	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	Z	346	Total	C	N	O	S	0	0
			2647	1674	448	515	10		
1	a	335	Total	C	N	O	S	0	0
			2565	1623	433	499	10		

- Molecule 2 is a protein called cement protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	0	97	Total	C	N	O	S	0	0
			704	447	116	140	1		
2	3	97	Total	C	N	O	S	0	0
			704	447	116	140	1		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
2	1	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	4	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	2	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	6	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	5	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	7	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	b	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	8	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	c	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	9	97	Total 704	C 447	N 116	O 140	S 1	0	0
2	d	97	Total 704	C 447	N 116	O 140	S 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: major capsid protein



- Molecule 1: major capsid protein




- Molecule 1: major capsid protein




- Molecule 1: major capsid protein






- Molecule 1: major capsid protein

Chain E: 16% 98%




- Molecule 1: major capsid protein

Chain F: 14% 97%




- Molecule 1: major capsid protein


Chain U: 11% 97%





- Molecule 1: major capsid protein

Chain V: 10% 97%









- Molecule 1: major capsid protein





- Molecule 1: major capsid protein



- Molecule 1: major capsid protein




- Molecule 1: major capsid protein




- Molecule 1: major capsid protein







• Molecule 2: cement protein




• Molecule 2: cement protein




• Molecule 2: cement protein




• Molecule 2: cement protein




• Molecule 2: cement protein




## • Molecule 2: cement protein




## • Molecule 2: cement protein



## • Molecule 2: cement protein




## • Molecule 2: cement protein



## • Molecule 2: cement protein




## • Molecule 2: cement protein




## • Molecule 2: cement protein





• Molecule 2: cement protein



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	9702	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE; CTFFIND4	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	25	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	59000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.140	Depositor
Minimum map value	-0.066	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.04	Depositor
Map size ( $\text{\AA}$ )	1088.0, 1088.0, 1088.0	wwPDB
Map dimensions	800, 800, 800	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.36, 1.36, 1.36	Depositor

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.37	0/2688	0.71	3/3647 (0.1%)
1	B	0.37	0/2688	0.69	2/3647 (0.1%)
1	C	0.37	0/2688	0.70	1/3647 (0.0%)
1	D	0.37	0/2688	0.71	1/3647 (0.0%)
1	E	0.37	1/2688 (0.0%)	0.68	0/3647
1	F	0.37	0/2688	0.70	2/3647 (0.1%)
1	U	0.38	0/2688	0.73	3/3647 (0.1%)
1	V	0.37	0/2688	0.70	2/3647 (0.1%)
1	W	0.37	0/2688	0.72	1/3647 (0.0%)
1	X	0.37	0/2688	0.69	1/3647 (0.0%)
1	Y	0.36	0/2688	0.69	1/3647 (0.0%)
1	Z	0.36	0/2688	0.69	1/3647 (0.0%)
1	a	0.34	0/2604	0.69	4/3531 (0.1%)
2	0	0.37	0/716	0.64	0/982
2	1	0.33	0/716	0.59	0/982
2	2	0.37	0/716	0.66	0/982
2	3	0.38	0/716	0.61	0/982
2	4	0.35	0/716	0.59	0/982
2	5	0.35	0/716	0.63	0/982
2	6	0.36	0/716	0.67	0/982
2	7	0.37	0/716	0.64	0/982
2	8	0.37	0/716	0.67	0/982
2	9	0.38	0/716	0.66	0/982
2	b	0.37	0/716	0.60	0/982
2	c	0.38	0/716	0.63	0/982
2	d	0.38	0/716	0.65	0/982
All	All	0.37	1/44168 (0.0%)	0.69	22/60061 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	D	0	2
1	F	0	3
1	U	0	1
1	V	0	2
1	W	0	2
1	Y	0	2
1	Z	0	4
1	a	0	1
2	2	0	2
2	5	0	3
2	6	0	2
2	7	0	2
2	8	0	4
2	9	0	2
2	b	0	3
2	c	0	3
2	d	0	2
All	All	0	45

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	336	ARG	CA-C	6.04	1.68	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	V	30	LEU	CA-CB-CG	7.55	132.68	115.30
1	F	30	LEU	CA-CB-CG	6.55	130.37	115.30
1	Z	6	LEU	CA-CB-CG	6.47	130.17	115.30
1	V	76	ILE	CG1-CB-CG2	-6.26	97.62	111.40
1	C	318	LEU	CA-CB-CG	6.19	129.53	115.30
1	A	248	MET	CA-CB-CG	6.17	123.79	113.30
1	W	136	ASP	CB-CG-OD1	6.09	123.78	118.30
1	U	309	LEU	CA-CB-CG	6.08	129.28	115.30
1	F	136	ASP	CB-CG-OD1	6.00	123.70	118.30
1	a	205	ILE	CG1-CB-CG2	-5.80	98.65	111.40
1	U	136	ASP	CB-CG-OD1	5.77	123.49	118.30
1	B	331	LEU	CA-CB-CG	5.43	127.79	115.30
1	A	161	LEU	CA-CB-CG	5.34	127.58	115.30
1	U	205	ILE	CG1-CB-CG2	-5.32	99.69	111.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	30	LEU	CA-CB-CG	5.27	127.42	115.30
1	D	237	LEU	CA-CB-CG	5.26	127.39	115.30
1	Y	309	LEU	CA-CB-CG	5.18	127.20	115.30
1	A	331	LEU	CA-CB-CG	5.17	127.19	115.30
1	X	136	ASP	CB-CG-OD1	5.17	122.95	118.30
1	a	331	LEU	CA-CB-CG	5.16	127.16	115.30
1	a	161	LEU	CA-CB-CG	5.07	126.95	115.30
1	a	228	MET	CA-CB-CG	5.06	121.90	113.30

There are no chirality outliers.

All (45) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	2	26	TYR	Peptide
2	2	30	GLY	Peptide
2	5	26	TYR	Peptide
2	5	30	GLY	Peptide
2	5	86	PRO	Peptide
2	6	26	TYR	Peptide
2	6	30	GLY	Peptide
2	7	26	TYR	Peptide
2	7	30	GLY	Peptide
2	8	26	TYR	Peptide
2	8	30	GLY	Peptide
2	8	85	THR	Peptide
2	8	86	PRO	Peptide
2	9	26	TYR	Peptide
2	9	30	GLY	Peptide
1	A	132	GLY	Peptide
1	A	178	ASP	Peptide
1	A	198	ARG	Peptide
1	B	237	LEU	Peptide
1	B	278	ASP	Peptide
1	D	198	ARG	Peptide
1	D	240	LEU	Peptide
1	F	181	GLN	Peptide
1	F	206	LEU	Peptide
1	F	73	GLN	Peptide
1	U	278	ASP	Peptide
1	V	278	ASP	Peptide
1	V	334	THR	Peptide
1	W	223	ARG	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	W	278	ASP	Peptide
1	Y	181	GLN	Peptide
1	Y	76	ILE	Peptide
1	Z	180	ARG	Peptide
1	Z	278	ASP	Peptide
1	Z	297	VAL	Peptide
1	Z	76	ILE	Peptide
1	a	198	ARG	Peptide
2	b	26	TYR	Peptide
2	b	30	GLY	Peptide
2	b	85	THR	Peptide
2	c	26	TYR	Peptide
2	c	30	GLY	Peptide
2	c	86	PRO	Peptide
2	d	26	TYR	Peptide
2	d	30	GLY	Peptide

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/348 (99%)	272 (79%)	70 (20%)	2 (1%)	25	65
1	B	344/348 (99%)	272 (79%)	72 (21%)	0	100	100
1	C	344/348 (99%)	274 (80%)	69 (20%)	1 (0%)	41	75
1	D	344/348 (99%)	251 (73%)	93 (27%)	0	100	100
1	E	344/348 (99%)	279 (81%)	65 (19%)	0	100	100
1	F	344/348 (99%)	282 (82%)	62 (18%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	U	344/348 (99%)	267 (78%)	77 (22%)	0	100	100
1	V	344/348 (99%)	273 (79%)	70 (20%)	1 (0%)	41	75
1	W	344/348 (99%)	266 (77%)	77 (22%)	1 (0%)	41	75
1	X	344/348 (99%)	266 (77%)	78 (23%)	0	100	100
1	Y	344/348 (99%)	275 (80%)	69 (20%)	0	100	100
1	Z	344/348 (99%)	273 (79%)	71 (21%)	0	100	100
1	a	331/348 (95%)	275 (83%)	56 (17%)	0	100	100
2	0	95/99 (96%)	84 (88%)	11 (12%)	0	100	100
2	1	95/99 (96%)	80 (84%)	15 (16%)	0	100	100
2	2	95/99 (96%)	80 (84%)	15 (16%)	0	100	100
2	3	95/99 (96%)	82 (86%)	13 (14%)	0	100	100
2	4	95/99 (96%)	81 (85%)	14 (15%)	0	100	100
2	5	95/99 (96%)	82 (86%)	13 (14%)	0	100	100
2	6	95/99 (96%)	81 (85%)	14 (15%)	0	100	100
2	7	95/99 (96%)	81 (85%)	14 (15%)	0	100	100
2	8	95/99 (96%)	79 (83%)	16 (17%)	0	100	100
2	9	95/99 (96%)	76 (80%)	18 (19%)	1 (1%)	14	54
2	b	95/99 (96%)	83 (87%)	12 (13%)	0	100	100
2	c	95/99 (96%)	81 (85%)	14 (15%)	0	100	100
2	d	95/99 (96%)	77 (81%)	18 (19%)	0	100	100
All	All	5694/5811 (98%)	4572 (80%)	1116 (20%)	6 (0%)	54	84

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	151	TYR
1	A	279	GLU
2	9	47	ASN
1	A	336	ARG
1	V	207	THR
1	W	233	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/285 (100%)	281 (99%)	3 (1%)	73	88
1	B	284/285 (100%)	280 (99%)	4 (1%)	67	85
1	C	284/285 (100%)	280 (99%)	4 (1%)	67	85
1	D	284/285 (100%)	282 (99%)	2 (1%)	84	93
1	E	284/285 (100%)	279 (98%)	5 (2%)	59	81
1	F	284/285 (100%)	279 (98%)	5 (2%)	59	81
1	U	284/285 (100%)	281 (99%)	3 (1%)	73	88
1	V	284/285 (100%)	280 (99%)	4 (1%)	67	85
1	W	284/285 (100%)	281 (99%)	3 (1%)	73	88
1	X	284/285 (100%)	278 (98%)	6 (2%)	53	79
1	Y	284/285 (100%)	278 (98%)	6 (2%)	53	79
1	Z	284/285 (100%)	281 (99%)	3 (1%)	73	88
1	a	275/285 (96%)	275 (100%)	0	100	100
2	0	73/74 (99%)	73 (100%)	0	100	100
2	1	73/74 (99%)	73 (100%)	0	100	100
2	2	73/74 (99%)	73 (100%)	0	100	100
2	3	73/74 (99%)	73 (100%)	0	100	100
2	4	73/74 (99%)	73 (100%)	0	100	100
2	5	73/74 (99%)	73 (100%)	0	100	100
2	6	73/74 (99%)	73 (100%)	0	100	100
2	7	73/74 (99%)	73 (100%)	0	100	100
2	8	73/74 (99%)	72 (99%)	1 (1%)	67	85
2	9	73/74 (99%)	73 (100%)	0	100	100
2	b	73/74 (99%)	73 (100%)	0	100	100
2	c	73/74 (99%)	73 (100%)	0	100	100
2	d	73/74 (99%)	73 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	4632/4667 (99%)	4583 (99%)	49 (1%)	74 88

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ARG
1	A	234	ARG
1	A	347	VAL
1	B	16	LEU
1	B	74	LEU
1	B	113	ARG
1	B	228	MET
1	C	223	ARG
1	C	228	MET
1	C	260	ARG
1	C	347	VAL
1	D	223	ARG
1	D	347	VAL
1	E	16	LEU
1	E	114	ASN
1	E	179	LYS
1	E	183	LEU
1	E	189	LEU
1	F	74	LEU
1	F	223	ARG
1	F	226	GLN
1	F	234	ARG
1	F	240	LEU
1	U	16	LEU
1	U	74	LEU
1	U	113	ARG
1	V	16	LEU
1	V	223	ARG
1	V	228	MET
1	V	347	VAL
1	W	16	LEU
1	W	74	LEU
1	W	228	MET
1	X	16	LEU
1	X	74	LEU
1	X	113	ARG
1	X	223	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	X	228	MET
1	X	347	VAL
1	Y	16	LEU
1	Y	74	LEU
1	Y	76	ILE
1	Y	113	ARG
1	Y	223	ARG
1	Y	347	VAL
1	Z	16	LEU
1	Z	74	LEU
1	Z	211	VAL
2	8	47	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (33) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	322	ASN
1	B	73	GLN
1	B	163	ASN
1	C	40	ASN
1	C	73	GLN
1	C	226	GLN
1	D	226	GLN
1	E	40	ASN
1	F	42	GLN
1	F	73	GLN
1	U	163	ASN
1	U	226	GLN
1	U	330	GLN
1	W	73	GLN
1	W	114	ASN
1	W	226	GLN
1	X	73	GLN
1	Y	40	ASN
1	Y	42	GLN
1	Y	73	GLN
1	Z	42	GLN
1	Z	73	GLN
1	Z	226	GLN
1	a	40	ASN
2	2	35	GLN
2	5	72	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	7	53	GLN
2	b	72	ASN
2	8	53	GLN
2	9	28	ASN
2	9	53	GLN
2	d	53	GLN
2	d	72	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-9774. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 400



Y Index: 400




Z Index: 400


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 684



Y Index: 684




Z Index: 116


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.04. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

## 6.5 Mask visualisation


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 10399 nm<sup>3</sup>; this corresponds to an approximate mass of 9393 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.283 Å<sup>-1</sup>

## 8 Fourier-Shell correlation ⓘ


This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-9774 and PDB model 6J3Q. Per-residue inclusion information can be found in section 3 on page 6.

### 9.1 Map-model overlays

#### 9.1.1 Map-model overlay [i](#)




#### 9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.04 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.04).























































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.04) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5928	 0.5020
0	 0.6182	 0.5030
1	 0.5159	 0.4960
2	 0.6081	 0.4990
3	 0.6210	 0.5050
4	 0.4870	 0.4990
5	 0.5821	 0.4980
6	 0.5749	 0.5000
7	 0.6153	 0.5000
8	 0.5994	 0.4950
9	 0.6095	 0.4970
A	 0.5995	 0.5020
B	 0.5933	 0.5070
C	 0.6091	 0.5040
D	 0.5922	 0.5010
E	 0.5756	 0.5020
F	 0.5825	 0.5040
U	 0.6072	 0.5050
V	 0.6064	 0.5010
W	 0.6022	 0.4990
X	 0.6022	 0.5050
Y	 0.6075	 0.5010
Z	 0.6149	 0.5050
a	 0.5208	 0.4990
b	 0.6081	 0.5070
c	 0.6052	 0.4970
d	 0.6297	 0.4990

