



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 11:51 am BST

PDB ID : 6J6X
Title : Crystal structure of apo GGTaseIII
Authors : Goto-Ito, S.; Yamagata, A.; Sato, Y.; Fukai, S.
Deposited on : 2019-01-16
Resolution : 2.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	<div> <div>2%</div> <div>67%</div> <div>9%</div> <div>24%</div> </div>
2	B	336	<div> <div>8%</div> <div>70%</div> <div>18%</div> <div>11%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4681 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein prenyltransferase alpha subunit repeat-containing protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	278	Total	As	C	N	O	S	0	0	0
			2318	2	1490	414	405	7			

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	328	GLY	-	insertion	UNP Q7Z6K3
A	329	ARG	-	insertion	UNP Q7Z6K3
A	330	PHE	-	insertion	UNP Q7Z6K3
A	331	PRO	-	insertion	UNP Q7Z6K3
A	332	HIS	-	insertion	UNP Q7Z6K3
A	333	SER	-	insertion	UNP Q7Z6K3
A	334	MET	-	insertion	UNP Q7Z6K3
A	335	THR	-	insertion	UNP Q7Z6K3
A	336	GLN	-	insertion	UNP Q7Z6K3
A	337	LEU	-	insertion	UNP Q7Z6K3
A	338	SER	-	insertion	UNP Q7Z6K3
A	339	PRO	-	insertion	UNP Q7Z6K3
A	340	ALA	-	insertion	UNP Q7Z6K3
A	341	ASP	-	insertion	UNP Q7Z6K3
A	342	SER	-	insertion	UNP Q7Z6K3
A	343	PRO	-	insertion	UNP Q7Z6K3
A	344	GLY	-	insertion	UNP Q7Z6K3
A	345	GLY	-	insertion	UNP Q7Z6K3
A	346	THR	-	insertion	UNP Q7Z6K3
A	347	LEU	-	insertion	UNP Q7Z6K3
A	348	SER	-	insertion	UNP Q7Z6K3
A	349	ASP	-	insertion	UNP Q7Z6K3
A	350	LEU	-	insertion	UNP Q7Z6K3
A	351	HIS	-	insertion	UNP Q7Z6K3
A	352	LEU	-	insertion	UNP Q7Z6K3
A	353	ILE	-	insertion	UNP Q7Z6K3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	354	PRO	-	insertion	UNP Q7Z6K3


- Molecule 2 is a protein called Geranylgeranyl transferase type-2 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	298	Total	C	N	O	S	0	0	0
			2346	1500	388	438	20			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP P53611
B	-3	PRO	-	expression tag	UNP P53611
B	-2	LEU	-	expression tag	UNP P53611
B	-1	GLY	-	expression tag	UNP P53611
B	0	SER	-	expression tag	UNP P53611

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 1: Protein prenyltransferase alpha subunit repeat-containing protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	88.54Å 88.54Å 647.58Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.48 – 2.96 49.47 – 2.96	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.48-2.96) 100.0 (49.47-2.96)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 2.96Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.266 , 0.297 0.271 , 0.302	Depositor DCC
R_{free} test set	1567 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	91.0	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 57.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4681	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, MG, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/2358	0.35	0/3201
2	B	0.21	0/2400	0.36	0/3248
All	All	0.20	0/4758	0.35	0/6449

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2318	0	2319	17	0
2	B	2346	0	2282	40	0
3	A	8	0	14	1	0
3	B	8	0	14	1	0
4	B	1	0	0	0	0
All	All	4681	0	4629	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:56:VAL:HG11	2:B:292:LEU:HA	1.77	0.66
2:B:278:PHE:HB2	2:B:291:THR:HA	1.79	0.64
1:A:112:ILE:HG23	1:A:119:PRO:HG3	1.82	0.62
2:B:168:ILE:HD13	2:B:206:GLN:HG3	1.83	0.60
2:B:106:THR:HG23	2:B:322:ARG:HH21	1.66	0.60
2:B:250:LYS:HG3	2:B:300:LEU:HD22	1.83	0.60
1:A:41:ARG:HB2	1:A:44:ARG:HB3	1.85	0.59
2:B:80:HIS:NE2	2:B:86:SER:OG	2.35	0.59
3:B:402:MPD:O2	3:B:402:MPD:O4	2.13	0.59
2:B:239:VAL:HG23	2:B:280:ASP:HA	1.84	0.59
2:B:243:TRP:HB2	2:B:293:PHE:HB3	1.87	0.57
2:B:264:ARG:NH2	2:B:303:GLU:OE1	2.34	0.56
2:B:250:LYS:HG2	2:B:255:LEU:HD22	1.88	0.56
1:A:29:ILE:HD13	1:A:54:LEU:HD23	1.89	0.55
1:A:84:ARG:HG3	1:A:116:THR:HG23	1.89	0.55
2:B:100:SER:HA	2:B:103:GLN:HE21	1.72	0.54
2:B:43:GLU:HA	2:B:46:ARG:HD3	1.88	0.54
1:A:69:HIS:NE2	2:B:43:GLU:OE1	2.40	0.53
1:A:123:LEU:O	1:A:143:ARG:NH2	2.42	0.53
2:B:142:ASP:OD2	2:B:144:ARG:NH1	2.42	0.53
2:B:239:VAL:HB	2:B:290:HIS:CD2	2.44	0.52
2:B:292:LEU:O	2:B:296:ALA:N	2.28	0.51
2:B:150:VAL:HG23	2:B:159:LEU:HD11	1.94	0.48
2:B:52:TRP:HA	2:B:292:LEU:HD11	1.94	0.48
2:B:57:MET:O	2:B:62:GLN:N	2.34	0.48
2:B:272:ASP:HB3	2:B:276:GLY:H	1.79	0.48
2:B:49:GLY:HA2	2:B:52:TRP:CE3	2.49	0.48
2:B:222:ARG:HB3	2:B:241:TYR:OH	2.13	0.48
1:A:57:GLU:OE2	1:A:59:TRP:NE1	2.40	0.48
2:B:142:ASP:OD1	2:B:143:THR:N	2.47	0.48
1:A:111:LEU:HD22	1:A:116:THR:HG21	1.95	0.47
1:A:75:TYR:HB2	1:A:82:LEU:HD13	1.98	0.46
2:B:239:VAL:HG12	2:B:293:PHE:HB2	1.98	0.46
2:B:237:PRO:HB2	2:B:240:CYS:SG	2.57	0.45
3:A:401:MPD:O4	3:A:401:MPD:O2	2.34	0.44
2:B:168:ILE:HA	2:B:200:PHE:HE1	1.83	0.44
1:A:74:LEU:HB3	1:A:80:GLN:HB3	1.99	0.44
2:B:293:PHE:HA	2:B:296:ALA:HB3	1.98	0.44
2:B:102:VAL:HG21	2:B:152:THR:HG23	1.99	0.44
1:A:50:VAL:HG22	1:A:53:LYS:O	2.18	0.44
1:A:119:PRO:HB2	1:A:150:LEU:HD21	2.00	0.43
1:A:118:ASN:HB3	1:A:121:LYS:HG3	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:280:ASP:HB3	2:B:290:HIS:CE1	2.53	0.43
2:B:54:LEU:HD11	2:B:71:ILE:HD13	2.00	0.43
1:A:289:PRO:O	1:A:293:GLU:HG2	2.19	0.42
2:B:15:PRO:HB2	2:B:16:ASP:H	1.68	0.42
2:B:211:ASN:HB3	2:B:214:LEU:HB3	2.01	0.42
2:B:316:PRO:HB2	2:B:319:VAL:HG23	2.00	0.42
1:A:12:VAL:HG21	1:A:67:TYR:CD2	2.55	0.42
2:B:215:LEU:HD23	2:B:252:ILE:HD11	2.02	0.42
2:B:53:GLY:O	2:B:57:MET:HG2	2.19	0.42
2:B:52:TRP:O	2:B:55:THR:HG22	2.20	0.41
2:B:246:LEU:HD11	2:B:263:LEU:HD21	2.03	0.41
2:B:239:VAL:HA	2:B:242:SER:HB2	2.03	0.41
1:A:145:TRP:O	1:A:149:GLN:HG2	2.20	0.41
1:A:37:CYS:HB3	1:A:57:GLU:HB3	2.02	0.40
2:B:249:LEU:HB3	2:B:255:LEU:HA	2.03	0.40
2:B:295:ILE:HG23	2:B:306:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/366 (74%)	265 (98%)	5 (2%)	0	100	100
2	B	294/336 (88%)	281 (96%)	13 (4%)	0	100	100
All	All	564/702 (80%)	546 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/332 (77%)	254 (99%)	2 (1%)	81	92
2	B	254/286 (89%)	253 (100%)	1 (0%)	91	96
All	All	510/618 (82%)	507 (99%)	3 (1%)	86	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	PHE
1	A	170	THR
2	B	18	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	315	HIS
2	B	23	HIS
2	B	103	GLN
2	B	271	GLN
2	B	290	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	92	1	5,8,9	1.35	0	1,9,11	0.87	0
1	CAS	A	60	1	5,8,9	1.32	0	1,9,11	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	92	1	-	0/0/7/9	-
1	CAS	A	60	1	-	0/0/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MPD	B	402	-	7,7,7	0.61	0	9,10,10	0.37	0
3	MPD	A	401	-	7,7,7	0.61	0	9,10,10	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	B	402	-	-	1/5/5/5	-
3	MPD	A	401	-	-	1/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	MPD	O2-C2-C3-C4
3	B	402	MPD	C2-C3-C4-O4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	MPD	1	0
3	A	401	MPD	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/366 (75%)	0.29	8 (2%) 51 35	60, 96, 132, 163	0
2	B	298/336 (88%)	0.50	27 (9%) 9 5	51, 81, 149, 158	0
All	All	574/702 (81%)	0.40	35 (6%) 21 12	51, 91, 144, 163	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	289	PHE	6.8
2	B	293	PHE	6.3
2	B	288	PRO	5.1
2	B	217	TRP	4.5
1	A	2	ALA	4.4
2	B	285	MET	4.3
1	A	327	ASN	4.1
1	A	3	GLU	4.0
2	B	266	PHE	4.0
2	B	237	PRO	3.9
2	B	323	VAL	3.6
2	B	241	TYR	3.5
2	B	292	LEU	3.4
2	B	34	LYS	3.4
2	B	322	ARG	3.2
2	B	275	THR	3.1
1	A	4	THR	2.9
1	A	252	VAL	2.9
2	B	278	PHE	2.5
2	B	35	ASP	2.4
2	B	251	ILE	2.4
2	B	261	GLU	2.4
2	B	284	ASP	2.4
1	A	208	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	238	ASP	2.3
2	B	46	ARG	2.3
2	B	86	SER	2.2
2	B	215	LEU	2.2
2	B	290	HIS	2.2
2	B	257	TRP	2.1
1	A	152	GLN	2.1
2	B	267	ILE	2.1
2	B	39	TYR	2.1
2	B	221	GLU	2.1
1	A	5	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	A	60	9/10	0.89	0.28	104,115,148,171	0
1	CAS	A	92	9/10	0.96	0.17	73,81,120,159	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	A	401	8/8	0.79	0.13	112,119,127,131	0
4	MG	B	401	1/1	0.88	0.33	67,67,67,67	0
3	MPD	B	402	8/8	0.90	0.22	68,95,101,106	0

6.5 Other polymers [i](#)

There are no such residues in this entry.