



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 09:05 am BST

PDB ID : 1JE8
Title : Two-Component response regulator NarL/DNA Complex: DNA Bending Found in a High Affinity Site
Authors : Maris, A.E.; Sawaya, M.R.; Kaczor-Grzeskowiak, M.; Jarvis, M.R.; Bearson, S.M.D.; Kopka, M.L.; Schroder, I.; Gunsalus, R.P.; Dickerson, R.E.
Deposited on : 2001-06-15
Resolution : 2.12 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	20	<div> <div>95%</div> <div>5%</div> </div>
1	D	20	<div> <div>70%</div> <div>25%</div> <div>5%</div> </div>
1	G	20	<div> <div>75%</div> <div>25%</div> </div>
1	H	20	<div> <div>85%</div> <div>15%</div> </div>
2	A	82	<div> <div>4%</div> <div>61%</div> <div>20%</div> <div>20%</div> </div>
2	B	82	<div> <div>2%</div> <div>55%</div> <div>23%</div> <div>20%</div> </div>
2	E	82	<div> <div>4%</div> <div>56%</div> <div>23%</div> <div>18%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	82	<p>2% 61% 18% 20%</p>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*CP*GP*TP*AP*CP*CP*CP*AP*TP*TP*AP*AP*TP*GP*GP*GP*TP*AP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	20	Total	C	N	O	P	0	0	0
			407	195	75	118	19			
1	D	20	Total	C	N	O	P	0	0	0
			407	195	75	118	19			
1	G	20	Total	C	N	O	P	0	0	0
			407	195	75	118	19			
1	H	20	Total	C	N	O	P	0	0	0
			407	195	75	118	19			

- Molecule 2 is a protein called Nitrate/Nitrite Response Regulator Protein NARL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	66	Total	C	N	O	Se	0	0	0
			548	347	107	91	3			
2	B	66	Total	C	N	O	Se	0	0	0
			548	347	107	91	3			
2	E	67	Total	C	N	O	Se	0	0	0
			557	352	108	94	3			
2	F	66	Total	C	N	O	Se	1	0	0
			548	347	107	91	3			

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MSE	-	EXPRESSION TAG	UNP P10957
A	-11	ARG	-	EXPRESSION TAG	UNP P10957
A	-10	GLY	-	EXPRESSION TAG	UNP P10957
A	-9	SER	-	EXPRESSION TAG	UNP P10957
A	-8	HIS	-	EXPRESSION TAG	UNP P10957
A	-7	HIS	-	EXPRESSION TAG	UNP P10957
A	-6	HIS	-	EXPRESSION TAG	UNP P10957

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP P10957
A	-4	HIS	-	EXPRESSION TAG	UNP P10957
A	-3	HIS	-	EXPRESSION TAG	UNP P10957
A	-2	GLY	-	EXPRESSION TAG	UNP P10957
A	-1	SER	-	EXPRESSION TAG	UNP P10957
A	175	MSE	MET	MODIFIED RESIDUE	UNP P10957
A	194	MSE	MET	MODIFIED RESIDUE	UNP P10957
A	198	MSE	MET	MODIFIED RESIDUE	UNP P10957
B	-12	MSE	-	EXPRESSION TAG	UNP P10957
B	-11	ARG	-	EXPRESSION TAG	UNP P10957
B	-10	GLY	-	EXPRESSION TAG	UNP P10957
B	-9	SER	-	EXPRESSION TAG	UNP P10957
B	-8	HIS	-	EXPRESSION TAG	UNP P10957
B	-7	HIS	-	EXPRESSION TAG	UNP P10957
B	-6	HIS	-	EXPRESSION TAG	UNP P10957
B	-5	HIS	-	EXPRESSION TAG	UNP P10957
B	-4	HIS	-	EXPRESSION TAG	UNP P10957
B	-3	HIS	-	EXPRESSION TAG	UNP P10957
B	-2	GLY	-	EXPRESSION TAG	UNP P10957
B	-1	SER	-	EXPRESSION TAG	UNP P10957
B	175	MSE	MET	MODIFIED RESIDUE	UNP P10957
B	194	MSE	MET	MODIFIED RESIDUE	UNP P10957
B	198	MSE	MET	MODIFIED RESIDUE	UNP P10957
E	-12	MSE	-	EXPRESSION TAG	UNP P10957
E	-11	ARG	-	EXPRESSION TAG	UNP P10957
E	-10	GLY	-	EXPRESSION TAG	UNP P10957
E	-9	SER	-	EXPRESSION TAG	UNP P10957
E	-8	HIS	-	EXPRESSION TAG	UNP P10957
E	-7	HIS	-	EXPRESSION TAG	UNP P10957
E	-6	HIS	-	EXPRESSION TAG	UNP P10957
E	-5	HIS	-	EXPRESSION TAG	UNP P10957
E	-4	HIS	-	EXPRESSION TAG	UNP P10957
E	-3	HIS	-	EXPRESSION TAG	UNP P10957
E	-2	GLY	-	EXPRESSION TAG	UNP P10957
E	-1	SER	-	EXPRESSION TAG	UNP P10957
E	175	MSE	MET	MODIFIED RESIDUE	UNP P10957
E	194	MSE	MET	MODIFIED RESIDUE	UNP P10957
E	198	MSE	MET	MODIFIED RESIDUE	UNP P10957
F	-12	MSE	-	EXPRESSION TAG	UNP P10957
F	-11	ARG	-	EXPRESSION TAG	UNP P10957
F	-10	GLY	-	EXPRESSION TAG	UNP P10957
F	-9	SER	-	EXPRESSION TAG	UNP P10957

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-8	HIS	-	EXPRESSION TAG	UNP P10957
F	-7	HIS	-	EXPRESSION TAG	UNP P10957
F	-6	HIS	-	EXPRESSION TAG	UNP P10957
F	-5	HIS	-	EXPRESSION TAG	UNP P10957
F	-4	HIS	-	EXPRESSION TAG	UNP P10957
F	-3	HIS	-	EXPRESSION TAG	UNP P10957
F	-2	GLY	-	EXPRESSION TAG	UNP P10957
F	-1	SER	-	EXPRESSION TAG	UNP P10957
F	175	MSE	MET	MODIFIED RESIDUE	UNP P10957
F	194	MSE	MET	MODIFIED RESIDUE	UNP P10957
F	198	MSE	MET	MODIFIED RESIDUE	UNP P10957

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	49	Total	O	0	0
			49	49		
4	D	47	Total	O	0	0
			47	47		
4	G	57	Total	O	0	0
			57	57		
4	H	53	Total	O	0	0
			53	53		
4	A	32	Total	O	0	0
			32	32		
4	B	30	Total	O	0	0
			30	30		
4	E	27	Total	O	0	0
			27	27		
4	F	35	Total	O	0	0
			35	35		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*CP*GP*TP*AP*CP*CP*CP*AP*TP*TP*AP*AP*TP*GP*GP*GP*TP*AP*CP*G)-3'

Chain C: 



- Molecule 1: 5'-D(*CP*GP*TP*AP*CP*CP*CP*AP*TP*TP*AP*AP*TP*GP*GP*GP*TP*AP*CP*G)-3'

Chain D: 




- Molecule 1: 5'-D(*CP*GP*TP*AP*CP*CP*CP*AP*TP*TP*AP*AP*TP*GP*GP*GP*TP*AP*CP*G)-3'

Chain G: 



- Molecule 1: 5'-D(*CP*GP*TP*AP*CP*CP*CP*AP*TP*TP*AP*AP*TP*GP*GP*GP*TP*AP*CP*G)-3'

Chain H: 




- Molecule 2: Nitrate/Nitrite Response Regulator Protein NARL

Chain A: 



- Molecule 2: Nitrate/Nitrite Response Regulator Protein NARL



- Molecule 2: Nitrate/Nitrite Response Regulator Protein NARL



- Molecule 2: Nitrate/Nitrite Response Regulator Protein NARL



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.26Å 52.74Å 83.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.97 – 2.12 56.42 – 2.12	Depositor EDS
% Data completeness (in resolution range)	96.2 (8.97-2.12) 96.1 (56.42-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.71 (at 2.12Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.228 , 0.273 0.231 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.801	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.479 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4199	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.63 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6581e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.54	0/456	0.85	0/702
1	D	0.49	0/456	0.83	2/702 (0.3%)
1	G	0.50	0/456	0.78	0/702
1	H	0.53	0/456	0.83	0/702
2	A	0.38	0/552	0.56	0/732
2	B	0.36	0/552	0.58	0/732
2	E	0.36	0/561	0.57	0/744
2	F	0.38	0/552	0.58	0/732
All	All	0.44	0/4041	0.71	2/5748 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	25	DC	O4'-C4'-C3'	-5.45	102.32	104.50
1	D	25	DC	C1'-O4'-C4'	-5.04	105.06	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	407	0	227	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	407	0	227	5	0
1	G	407	0	227	3	0
1	H	407	0	227	2	0
2	A	548	0	596	22	0
2	B	548	0	596	19	0
2	E	557	0	602	17	0
2	F	548	0	596	14	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
3	E	10	0	0	0	0
3	F	10	0	0	0	0
4	A	32	0	0	0	0
4	B	30	0	0	1	0
4	C	49	0	0	0	0
4	D	47	0	0	0	0
4	E	27	0	0	0	0
4	F	35	0	0	0	0
4	G	57	0	0	0	0
4	H	53	0	0	0	0
All	All	4199	0	3298	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:153:VAL:HA	2:A:198:MSE:HE1	1.26	1.10
2:F:210:VAL:HA	2:F:215:ILE:HD13	1.47	0.95
2:A:214:ARG:HG3	2:A:214:ARG:HH21	1.51	0.74
2:E:153:VAL:HG21	2:E:215:ILE:HG21	1.69	0.74
2:A:153:VAL:HA	2:A:198:MSE:CE	2.13	0.73

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	64/82 (78%)	64 (100%)	0	0	100	100
2	B	64/82 (78%)	64 (100%)	0	0	100	100
2	E	65/82 (79%)	62 (95%)	2 (3%)	1 (2%)	10	5
2	F	64/82 (78%)	63 (98%)	1 (2%)	0	100	100
All	All	257/328 (78%)	253 (98%)	3 (1%)	1 (0%)	34	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	152	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	61/70 (87%)	58 (95%)	3 (5%)	25	22
2	B	61/70 (87%)	59 (97%)	2 (3%)	38	39
2	E	62/70 (89%)	61 (98%)	1 (2%)	62	68
2	F	61/70 (87%)	58 (95%)	3 (5%)	25	22
All	All	245/280 (88%)	236 (96%)	9 (4%)	34	34

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	200	LEU
2	F	200	LEU
2	F	151	ARG
2	A	195	LEU
2	E	155	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	155	GLN
2	F	212	GLN
2	E	155	GLN
2	B	154	ASN
2	B	211	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	F	408	-	4,4,4	0.26	0	6,6,6	0.18	0
3	SO4	E	407	-	4,4,4	0.29	0	6,6,6	0.12	0
3	SO4	B	406	-	4,4,4	0.24	0	6,6,6	0.14	0
3	SO4	E	403	-	4,4,4	0.18	0	6,6,6	0.20	0
3	SO4	B	402	-	4,4,4	0.19	0	6,6,6	0.26	0
3	SO4	A	405	-	4,4,4	0.24	0	6,6,6	0.12	0
3	SO4	A	401	-	4,4,4	0.15	0	6,6,6	0.17	0
3	SO4	F	404	-	4,4,4	0.36	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	20/20 (100%)	-0.49	0 100 100	22, 33, 38, 39	0
1	D	20/20 (100%)	-0.46	0 100 100	21, 34, 39, 39	0
1	G	20/20 (100%)	-0.53	0 100 100	21, 34, 39, 39	0
1	H	20/20 (100%)	-0.49	0 100 100	21, 34, 38, 39	0
2	A	63/82 (76%)	0.12	3 (4%) 30 35	21, 32, 60, 70	0
2	B	63/82 (76%)	0.05	2 (3%) 47 54	21, 31, 54, 63	0
2	E	64/82 (78%)	0.05	3 (4%) 31 36	22, 31, 60, 71	0
2	F	63/82 (76%)	0.04	2 (3%) 47 54	21, 31, 61, 65	1 (1%)
All	All	333/408 (81%)	-0.07	10 (3%) 50 56	21, 32, 60, 71	1 (0%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	216	PHE	4.6
2	B	216	PHE	3.3
2	F	216	PHE	3.1
2	A	215	ILE	2.9
2	B	214	ARG	2.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	405	5/5	0.93	0.14	76,76,76,77	0
3	SO4	F	408	5/5	0.94	0.10	73,73,74,74	0
3	SO4	B	406	5/5	0.95	0.12	69,70,70,70	0
3	SO4	E	407	5/5	0.97	0.10	66,67,67,67	0
3	SO4	B	402	5/5	0.99	0.09	31,31,32,33	0
3	SO4	E	403	5/5	0.99	0.09	32,33,34,35	0
3	SO4	A	401	5/5	0.99	0.11	30,30,34,34	0
3	SO4	F	404	5/5	0.99	0.07	31,32,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.