



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2020 – 05:04 PM EST

PDB ID : 7JI2  
Title : Crystal Structure of H2-Kb in complex with a OVA mutant peptide  
Authors : Li, X.; Mallis, R.J.; Mizsei, R.; Tan, K.; Reinherz, E.L.; Wang, J.  
Deposited on : 2020-07-22  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.15.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.15.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	<div> <div>6%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>..</div> </div> </div>
1	D	281	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>10%</div> <div>..</div> </div> </div>
2	B	100	<div> <div></div> <div> <div></div> <div>88%</div> <div>10%</div> <div>.</div> </div> </div>
2	E	100	<div> <div></div> <div> <div></div> <div>77%</div> <div>20%</div> <div>..</div> </div> </div>
3	C	8	<div> <div></div> <div> <div></div> <div>100%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	8	 100%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6866 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	278	Total	C	N	O	S	0	2	0
			2241	1417	394	421	9			
1	D	278	Total	C	N	O	S	0	4	0
			2289	1443	405	432	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P01901
D	0	MET	-	initiating methionine	UNP P01901

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	98	Total	C	N	O	S	0	0	0
			809	515	136	151	7			
2	E	98	Total	C	N	O	S	0	2	0
			827	525	138	157	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P01887
B	85	ASP	ALA	conflict	UNP P01887
E	0	MET	-	initiating methionine	UNP P01887
E	85	ASP	ALA	conflict	UNP P01887

- Molecule 3 is a protein called OVA mutant peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	8	Total	C	N	O	0	0	0
			70	46	11	13			

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	8	Total	C	N	O	0	0	0
			70	46	11	13			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	166	Total 166	O 166	0	0
5	B	83	Total 83	O 83	0	0
5	D	176	Total 177	O 177	0	1
5	E	71	Total 71	O 71	0	0
5	C	9	Total 9	O 9	0	0
5	F	6	Total 6	O 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: OVA mutant peptide

Chain C:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: OVA mutant peptide

Chain F:  100%

There are no outlier residues recorded for this chain.



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.43Å 90.01Å 89.67Å 90.00° 111.70° 90.00°	Depositor
Resolution (Å)	45.01 – 1.95 45.01 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.0 (45.01-1.95) 98.0 (45.01-1.95)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	131.37 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.170 , 0.185 0.168 , 0.180	Depositor DCC
$R_{free}$ test set	3564 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtriage
Anisotropy	0.549	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 29.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	0.277 for h,-k,-h-l	Xtriage
Reported twinning fraction	0.330 for h,-k,-h-l	Depositor
Outliers	1 of 70388 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6866	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.35 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2173e-04.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	1/2305 (0.0%)	0.60	0/3139
1	D	0.41	0/2353	0.60	0/3200
2	B	0.32	0/835	0.56	0/1133
2	E	0.40	0/853	0.58	0/1157
3	C	0.72	0/71	0.75	0/93
3	F	0.70	0/71	0.72	0/93
All	All	0.41	1/6488 (0.0%)	0.59	0/8815

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	61	GLU	C-N	-5.74	1.20	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2241	0	2106	22	0
1	D	2289	0	2171	19	0
2	B	809	0	771	8	0
2	E	827	0	781	13	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	70	0	70	0	0
3	F	70	0	70	0	0
4	A	12	0	16	1	0
4	B	12	0	16	2	0
4	C	6	0	8	0	0
4	D	18	0	24	2	0
5	A	166	0	0	2	0
5	B	83	0	0	1	0
5	C	9	0	0	0	0
5	D	177	0	0	1	0
5	E	71	0	0	3	0
5	F	6	0	0	0	0
All	All	6866	0	6033	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ARG:HG2	1:A:194:ARG:HH21	1.24	0.99
1:A:111:ARG:HG2	1:A:111:ARG:HH21	1.41	0.83
1:A:194:ARG:NH2	1:A:194:ARG:HG2	1.97	0.72
1:D:219:LEU:O	1:D:222:GLU:HG2	1.91	0.70
2:E:17:ASN:ND2	5:E:101:HOH:O	2.26	0.68
1:A:111:ARG:NH2	1:A:111:ARG:HG2	2.13	0.64
1:A:224:LEU:O	1:A:228:MET:HE3	1.99	0.62
1:D:228:MET:HE2	1:D:245:ALA:HB1	1.82	0.61
2:B:23:LEU:HB2	2:B:70:PHE:CD1	2.37	0.59
1:D:220:ASN:N	4:D:303:GOL:O1	2.31	0.58
1:A:28:VAL:HG13	1:A:33:PHE:CD1	2.41	0.54
1:A:230:LEU:HD12	1:A:230:LEU:O	2.06	0.54
2:B:58:LYS:HD3	2:B:58:LYS:N	2.22	0.54
1:D:194[A]:ARG:HH21	1:D:194[A]:ARG:HG2	1.72	0.53
1:D:220:ASN:H	4:D:303:GOL:HO1	1.56	0.53
2:E:40:LEU:HD23	2:E:45:LYS:HA	1.93	0.51
1:D:214:THR:HB	1:D:262:TYR:HB2	1.91	0.51
2:E:50:GLU:OE1	2:E:67:HIS:NE2	2.44	0.51
1:D:194[A]:ARG:NH2	1:D:194[A]:ARG:HG2	2.26	0.50
2:B:23:LEU:HB2	2:B:70:PHE:CE1	2.47	0.50
1:A:45:TYR:HE1	1:A:67:ALA:HB2	1.76	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:176:ASN:O	1:D:180:LEU:HG	2.12	0.50
2:E:28:THR:HG22	2:E:63:TYR:HB2	1.94	0.50
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.48	0.49
2:E:38:GLN:HG3	2:E:81:ARG:HB3	1.95	0.49
1:A:224:LEU:O	1:A:228:MET:CE	2.61	0.48
1:D:209:TYR:CD1	1:D:210:PRO:HA	2.50	0.47
1:A:166:GLU:OE1	4:A:301:GOL:O1	2.24	0.47
1:D:261:VAL:HB	1:D:270:LEU:HB2	1.95	0.46
1:A:268:GLU:HB3	1:D:169:ARG:HD3	1.96	0.46
2:B:2:GLN:HE22	2:B:85:ASP:HB3	1.81	0.46
1:D:21:ARG:NH2	5:D:411:HOH:O	2.49	0.45
1:A:75:ARG:NH1	5:A:411:HOH:O	2.49	0.45
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.52	0.45
2:E:31:HIS:ND1	5:E:102:HOH:O	2.36	0.45
1:A:93:HIS:NE2	5:A:403:HOH:O	2.36	0.45
2:B:11:SER:O	4:B:102:GOL:O1	2.32	0.44
2:B:70:PHE:HE2	2:B:72:PRO:HB3	1.83	0.44
2:B:28:THR:HG22	2:B:63:TYR:HB2	2.00	0.44
1:A:261:VAL:HB	1:A:270:LEU:HB2	1.99	0.43
1:D:217:TRP:N	1:D:228:MET:HE3	2.34	0.43
1:A:54:GLN:H	1:A:54:GLN:HG2	1.51	0.43
1:A:16:GLY:O	1:A:17:LEU:HD23	2.19	0.43
1:D:48:ARG:NH2	2:E:53:ASP:OD2	2.52	0.42
2:E:23:LEU:HD23	2:E:39:MET:CE	2.49	0.42
1:A:230:LEU:C	1:A:230:LEU:HD12	2.40	0.42
1:D:22:TYR:CD2	1:D:71:GLU:HB2	2.54	0.42
4:B:102:GOL:H12	5:B:218:HOH:O	2.20	0.42
2:E:2:GLN:HG2	2:E:32:PRO:CD	2.49	0.42
1:A:159:TYR:CD1	1:A:163:THR:HB	2.55	0.41
1:D:41:GLU:HG2	1:D:41:GLU:H	1.58	0.41
2:E:38:GLN:HG2	5:E:139:HOH:O	2.18	0.41
2:E:57:SER:OG	2:E:59:ASP:OD1	2.39	0.41
1:D:201:LEU:O	1:D:246:SER:HA	2.21	0.41
1:A:14:ARG:NH2	1:A:39:ASP:OD1	2.50	0.41
1:A:151:GLY:O	1:A:154:GLU:HB2	2.22	0.40
1:D:127:ASN:OD1	1:D:134:THR:OG1	2.23	0.40
2:E:13:HIS:HB3	2:E:14:PRO:HD2	2.02	0.40
1:A:3:HIS:ND1	1:A:29:ASP:OD2	2.49	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/281 (99%)	269 (97%)	9 (3%)	0	100	100
1	D	280/281 (100%)	273 (98%)	7 (2%)	0	100	100
2	B	96/100 (96%)	93 (97%)	3 (3%)	0	100	100
2	E	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
3	C	6/8 (75%)	6 (100%)	0	0	100	100
3	F	6/8 (75%)	6 (100%)	0	0	100	100
All	All	764/778 (98%)	743 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/239 (96%)	224 (98%)	5 (2%)	52	44
1	D	238/239 (100%)	232 (98%)	6 (2%)	47	38
2	B	92/95 (97%)	92 (100%)	0	100	100
2	E	94/95 (99%)	92 (98%)	2 (2%)	53	46
3	C	8/8 (100%)	8 (100%)	0	100	100
3	F	8/8 (100%)	8 (100%)	0	100	100
All	All	669/684 (98%)	656 (98%)	13 (2%)	59	50

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ARG
1	A	45	TYR
1	A	54	GLN
1	A	194	ARG
1	A	220	ASN
1	D	30	ASP
1	D	41	GLU
1	D	45	TYR
1	D	149	GLN
1	D	194[A]	ARG
1	D	194[B]	ARG
2	E	38	GLN
2	E	70	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	2	GLN
2	E	38	GLN
3	C	4	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	301	-	5,5,5	0.91	0	5,5,5	0.99	0
4	GOL	C	101	-	5,5,5	0.12	0	5,5,5	0.35	0
4	GOL	B	101	-	5,5,5	0.15	0	5,5,5	0.28	0
4	GOL	B	102	-	5,5,5	0.88	0	5,5,5	1.05	0
4	GOL	D	301	-	5,5,5	0.91	0	5,5,5	0.98	0
4	GOL	D	302	-	5,5,5	0.89	0	5,5,5	1.01	0
4	GOL	D	303	-	5,5,5	0.88	0	5,5,5	1.04	0
4	GOL	A	302	-	5,5,5	0.92	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	301	-	-	0/4/4/4	-
4	GOL	C	101	-	-	2/4/4/4	-
4	GOL	B	101	-	-	1/4/4/4	-
4	GOL	B	102	-	-	0/4/4/4	-
4	GOL	D	301	-	-	1/4/4/4	-
4	GOL	D	302	-	-	2/4/4/4	-
4	GOL	D	303	-	-	3/4/4/4	-
4	GOL	A	302	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	303	GOL	O1-C1-C2-C3
4	A	302	GOL	C1-C2-C3-O3
4	D	303	GOL	O1-C1-C2-O2
4	A	302	GOL	O2-C2-C3-O3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	C	101	GOL	O1-C1-C2-C3
4	D	302	GOL	O1-C1-C2-C3
4	D	303	GOL	C1-C2-C3-O3
4	C	101	GOL	O1-C1-C2-O2
4	D	301	GOL	O2-C2-C3-O3
4	D	302	GOL	O1-C1-C2-O2
4	B	101	GOL	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	301	GOL	1	0
4	B	102	GOL	2	0
4	D	303	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	278/281 (98%)	0.25	17 (6%) 21 29	20, 32, 53, 65	0
1	D	278/281 (98%)	0.23	15 (5%) 25 34	19, 31, 49, 76	0
2	B	98/100 (98%)	-0.00	0 100 100	21, 31, 44, 49	0
2	E	98/100 (98%)	0.05	0 100 100	22, 32, 45, 53	0
3	C	8/8 (100%)	-0.03	0 100 100	24, 30, 34, 35	0
3	F	8/8 (100%)	-0.19	0 100 100	21, 25, 34, 37	0
All	All	768/778 (98%)	0.18	32 (4%) 36 45	19, 32, 49, 76	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	225	ILE	4.8
1	A	225	ILE	3.5
1	A	179	LEU	3.5
1	D	180	LEU	3.4
1	D	41	GLU	3.2
1	A	226	GLN	3.2
1	A	220	ASN	3.2
1	D	278	PRO	3.0
1	D	181	ARG	2.9
1	A	181	ARG	2.7
1	A	195	PRO	2.7
1	D	121	CYS	2.7
1	A	42	ASN	2.7
1	A	227	ASP	2.6
1	A	121	CYS	2.6
1	D	195	PRO	2.6
1	D	42	ASN	2.5
1	A	193	SER	2.4
1	A	173	LYS	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	1	GLY	2.3
1	D	176	ASN	2.3
1	D	226	GLN	2.3
1	D	227	ASP	2.3
1	D	193[A]	SER	2.2
1	A	278	PRO	2.1
1	A	40	ALA	2.1
1	D	145	HIS	2.1
1	A	145	HIS	2.1
1	A	180	LEU	2.1
1	A	54	GLN	2.0
1	D	16	GLY	2.0
1	D	138	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	D	301	6/6	0.55	0.26	32,37,39,40	0
4	GOL	A	302	6/6	0.73	0.29	39,45,50,50	0
4	GOL	D	302	6/6	0.83	0.26	41,42,45,46	0
4	GOL	B	102	6/6	0.85	0.24	28,32,33,37	0
4	GOL	C	101	6/6	0.86	0.21	34,45,50,59	0
4	GOL	D	303	6/6	0.87	0.39	35,39,42,46	0
4	GOL	A	301	6/6	0.88	0.26	33,35,37,42	0
4	GOL	B	101	6/6	0.89	0.18	35,36,40,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.