



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 09:08 am BST

PDB ID : 6JL8
Title : Crystal structure of GMP reductase C318A from Trypanosoma brucei
Authors : Mase, H.; Imamura, A.; Nishimura, S.; Inui, T.
Deposited on : 2019-03-04
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	504	
1	B	504	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6134 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GMP reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	453	Total	C	N	O	S	0	0	0
			3149	1949	581	595	24			
1	B	442	Total	C	N	O	S	0	0	0
			2966	1831	535	578	22			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	318	ALA	CYS	engineered mutation	UNP Q57ZS7
A	492	LYS	-	expression tag	UNP Q57ZS7
A	493	LEU	-	expression tag	UNP Q57ZS7
A	494	ALA	-	expression tag	UNP Q57ZS7
A	495	ALA	-	expression tag	UNP Q57ZS7
A	496	ALA	-	expression tag	UNP Q57ZS7
A	497	LEU	-	expression tag	UNP Q57ZS7
A	498	GLU	-	expression tag	UNP Q57ZS7
A	499	HIS	-	expression tag	UNP Q57ZS7
A	500	HIS	-	expression tag	UNP Q57ZS7
A	501	HIS	-	expression tag	UNP Q57ZS7
A	502	HIS	-	expression tag	UNP Q57ZS7
A	503	HIS	-	expression tag	UNP Q57ZS7
A	504	HIS	-	expression tag	UNP Q57ZS7
B	318	ALA	CYS	engineered mutation	UNP Q57ZS7
B	492	LYS	-	expression tag	UNP Q57ZS7
B	493	LEU	-	expression tag	UNP Q57ZS7
B	494	ALA	-	expression tag	UNP Q57ZS7
B	495	ALA	-	expression tag	UNP Q57ZS7
B	496	ALA	-	expression tag	UNP Q57ZS7
B	497	LEU	-	expression tag	UNP Q57ZS7
B	498	GLU	-	expression tag	UNP Q57ZS7
B	499	HIS	-	expression tag	UNP Q57ZS7
B	500	HIS	-	expression tag	UNP Q57ZS7
B	501	HIS	-	expression tag	UNP Q57ZS7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	502	HIS	-	expression tag	UNP Q57ZS7
B	503	HIS	-	expression tag	UNP Q57ZS7
B	504	HIS	-	expression tag	UNP Q57ZS7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	16	Total	O	0	0
			16	16		
2	B	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GMP reductase



• Molecule 1: GMP reductase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	124.56 Å 124.56 Å 540.17 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	21.81 – 2.80 49.51 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (21.81-2.80) 99.9 (49.51-2.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.81 Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.223 , 0.258 0.224 , 0.258	Depositor DCC
R_{free} test set	2642 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	62.1	Xtriage
Anisotropy	0.172	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 68.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6134	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/3177	0.68	0/4312
1	B	0.39	0/2991	0.56	0/4075
All	All	0.46	0/6168	0.63	0/8387

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3149	0	3090	49	0
1	B	2966	0	2786	39	0
2	A	16	0	0	0	0
2	B	3	0	0	0	0
All	All	6134	0	5876	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:PRO:HD2	1:B:139:ARG:HH22	1.46	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:84:GLN:HG2	1:A:248:LEU:HD21	1.63	0.80
1:B:95:GLN:HG3	1:B:229:LEU:HG	1.65	0.77
1:A:88:LEU:HD11	1:A:253:ALA:HB2	1.73	0.70
1:B:268:CYS:O	1:B:272:VAL:HG23	1.95	0.67
1:A:134:ASN:HB3	1:A:142:LEU:HD11	1.76	0.67
1:A:217:LYS:O	1:A:219:LYS:N	2.29	0.66
1:A:51:ALA:HB2	1:A:66:MET:HG3	1.78	0.66
1:A:301:ILE:HG12	1:A:346:VAL:HG21	1.78	0.65
1:B:80:SER:HG	1:B:82:GLU:HG2	1.62	0.65
1:B:80:SER:OG	1:B:82:GLU:HG2	1.96	0.65
1:B:53:ASN:HB3	1:B:74:ILE:HG22	1.80	0.63
1:A:171:VAL:HG12	1:A:172:SER:N	2.13	0.61
1:A:34:ASN:HD21	1:A:36:THR:HG23	1.65	0.61
1:A:383:ALA:O	1:A:396:LYS:NZ	2.35	0.59
1:B:167:ASP:OD1	1:B:168:LYS:N	2.36	0.59
1:B:95:GLN:CG	1:B:229:LEU:HG	2.32	0.59
1:B:171:VAL:HG12	1:B:172:SER:N	2.16	0.59
1:A:39:LEU:HD22	1:A:72:ILE:HG21	1.84	0.58
1:A:177:ILE:HG12	1:A:181:GLU:HB3	1.85	0.58
1:B:465:GLU:OE1	1:B:469:ARG:NH2	2.37	0.57
1:B:178:SER:OG	1:B:181:GLU:N	2.20	0.57
1:B:376:MET:HE1	1:B:447:LEU:HD21	1.86	0.57
1:A:131:LEU:HD11	1:A:206:LEU:HD13	1.86	0.56
1:A:260:ILE:HD11	1:A:268:CYS:HB2	1.88	0.56
1:B:269:ILE:HG23	1:B:303:ALA:HB2	1.88	0.56
1:B:80:SER:HG	1:B:83:GLU:H	1.53	0.55
1:A:92:LYS:NZ	1:A:254:ASP:OD2	2.34	0.55
1:A:464:GLU:OE2	1:A:468:ARG:NH2	2.40	0.55
1:B:351:ASP:HA	1:B:372:MET:HB3	1.90	0.54
1:B:217:LYS:O	1:B:219:LYS:N	2.41	0.54
1:B:84:GLN:HG2	1:B:248:LEU:HD21	1.90	0.54
1:B:127:VAL:O	1:B:147:ARG:NH2	2.42	0.53
1:B:178:SER:HG	1:B:181:GLU:H	1.54	0.53
1:A:257:VAL:HG22	1:A:287:ILE:HB	1.91	0.52
1:B:302:ASP:OD1	1:B:344:ARG:NH2	2.43	0.51
1:B:287:ILE:HD13	1:B:349:ILE:HD11	1.92	0.51
1:A:34:ASN:ND2	1:A:36:THR:HG23	2.25	0.50
1:A:119:ASN:ND2	1:A:147:ARG:HH22	2.10	0.49
1:A:171:VAL:CG1	1:A:172:SER:N	2.75	0.49
1:A:242:MET:CG	1:A:274:ARG:HD2	2.42	0.49
1:B:274:ARG:HB2	1:B:274:ARG:HH11	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:297:ALA:O	1:B:301:ILE:HG13	2.13	0.49
1:B:58:CYS:HA	1:B:62:MET:HB3	1.94	0.48
1:A:84:GLN:CG	1:A:248:LEU:HD21	2.41	0.48
1:B:171:VAL:CG1	1:B:172:SER:N	2.76	0.48
1:B:145:ILE:HD11	1:B:150:LEU:HD21	1.95	0.48
1:A:451:LEU:O	1:A:455:MET:HG3	2.14	0.47
1:A:40:SER:C	1:A:347:PRO:HG2	2.34	0.47
1:B:301:ILE:HG23	1:B:346:VAL:HG21	1.95	0.47
1:A:396:LYS:O	1:A:433:SER:HA	2.15	0.46
1:A:29:SER:O	1:A:32:GLU:HG2	2.15	0.46
1:A:390:LYS:HB3	1:A:390:LYS:HE3	1.88	0.46
1:A:291:ILE:HD11	1:A:308:LEU:HD13	1.98	0.46
1:A:99:ILE:HB	1:A:206:LEU:O	2.16	0.45
1:B:184:HIS:NE2	1:B:188:LYS:HD3	2.31	0.45
1:B:360:ILE:HD12	1:B:447:LEU:HD22	1.98	0.45
1:A:105:ILE:O	1:A:132:VAL:HA	2.16	0.45
1:A:360:ILE:HG12	1:A:371:VAL:HG21	1.98	0.45
1:A:60:GLN:O	1:A:64:VAL:HG23	2.17	0.45
1:A:55:ASP:HA	1:A:76:HIS:CD2	2.51	0.45
1:A:40:SER:CA	1:A:347:PRO:HG2	2.47	0.45
1:B:97:PHE:HD2	1:B:97:PHE:H	1.64	0.45
1:B:315:GLY:H	1:B:318:ALA:HB2	1.82	0.45
1:B:51:ALA:HB2	1:B:66:MET:SD	2.57	0.44
1:B:257:VAL:HG22	1:B:287:ILE:HB	1.98	0.44
1:B:77:ARG:HB3	1:B:234:ALA:O	2.17	0.44
1:A:330:GLN:OE1	1:A:330:GLN:HA	2.17	0.44
1:A:92:LYS:HE2	1:A:252:GLY:O	2.17	0.44
1:A:13:LEU:HD22	1:A:17:ASP:HB3	2.00	0.43
1:A:193:ASN:OD1	1:A:208:THR:HG23	2.18	0.43
1:A:179:LEU:O	1:A:183:THR:OG1	2.29	0.43
1:A:37:THR:HG21	1:A:48:PRO:HB3	2.00	0.43
1:A:242:MET:HG2	1:A:274:ARG:HD2	2.00	0.43
1:A:131:LEU:HD13	1:A:203:LEU:HD11	2.02	0.42
1:A:302:ASP:OD1	1:A:344:ARG:NH2	2.50	0.42
1:A:341:ALA:HB1	1:A:346:VAL:O	2.20	0.42
1:A:463:ILE:HD13	1:A:463:ILE:HA	1.96	0.42
1:B:258:VAL:HG22	1:B:288:ALA:HA	2.01	0.42
1:B:131:LEU:HD13	1:B:203:LEU:HD11	2.01	0.41
1:B:262:HIS:CE1	1:B:264:HIS:HB3	2.55	0.41
1:A:73:GLY:O	1:A:74:ILE:HD13	2.20	0.41
1:B:441:GLY:O	1:B:445:ARG:HG3	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:VAL:HG23	1:A:362:LYS:HD3	2.02	0.41
1:A:206:LEU:HG	1:A:207:VAL:N	2.34	0.41
1:B:269:ILE:HG12	1:B:300:LEU:HD23	2.03	0.41
1:A:51:ALA:HB3	1:A:73:GLY:HA2	2.03	0.41
1:A:330:GLN:O	1:A:334:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/504 (89%)	412 (92%)	33 (7%)	4 (1%)	17	46
1	B	436/504 (86%)	402 (92%)	31 (7%)	3 (1%)	22	53
All	All	885/1008 (88%)	814 (92%)	64 (7%)	7 (1%)	19	49

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	217	LYS
1	A	218	ASN
1	B	218	ASN
1	B	314	PRO
1	A	292	ALA
1	B	121	LYS
1	A	400	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/403 (75%)	302 (100%)	1 (0%)	92	98
1	B	272/403 (68%)	266 (98%)	6 (2%)	52	83
All	All	575/806 (71%)	568 (99%)	7 (1%)	71	92

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	B	46	SER
1	B	66	MET
1	B	135	CYS
1	B	158	THR
1	B	210	SER
1	B	458	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	119	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	453/504 (89%)	0.18	13 (2%)	51	41	29, 46, 92, 129	0
1	B	442/504 (87%)	0.23	15 (3%)	45	35	54, 80, 98, 135	0
All	All	895/1008 (88%)	0.21	28 (3%)	49	39	29, 72, 97, 135	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	391	ASP	5.1
1	A	393	GLN	4.5
1	A	394	LYS	4.1
1	A	159	VAL	4.0
1	B	251	ALA	3.7
1	B	385	GLY	3.3
1	B	248	LEU	3.1
1	B	229	LEU	2.7
1	A	392	GLY	2.7
1	A	390	LYS	2.7
1	B	436	CYS	2.7
1	A	401	MET	2.5
1	A	131	LEU	2.5
1	A	433	SER	2.5
1	B	131	LEU	2.4
1	B	126	GLY	2.4
1	A	135	CYS	2.4
1	B	398	ILE	2.4
1	B	270	ASN	2.3
1	A	385	GLY	2.3
1	B	433	SER	2.2
1	B	75	LEU	2.1
1	B	124	VAL	2.1
1	B	242	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	122	GLY	2.1
1	A	389	VAL	2.1
1	B	384	PRO	2.0
1	A	115	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.