



## wwPDB EM Validation Summary Report ⓘ

Nov 13, 2022 – 04:53 PM EST

PDB ID : 7JL2  
EMDB ID : EMD-22370  
Title : Cryo-EM structure of MDA5-dsRNA filament in complex with TRIM65 PSpry domain (Trimer)  
Authors : Kato, K.; Ahmad, S.; Hur, S.  
Deposited on : 2020-07-29  
Resolution : 4.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	44	
2	Y	44	
3	A	739	
3	C	739	
3	E	739	
4	B	191	
4	D	191	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	191	<div><div></div><div>72%</div><div>59%</div><div>36%</div><div>6%</div></div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 22128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (44-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	X	44	Total	C	N	O	P	0	0
			944	422	175	303	44		

- Molecule 2 is a RNA chain called RNA (44-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	Y	44	Total	C	N	O	P	0	0
			925	414	154	313	44		

- Molecule 3 is a protein called Interferon-induced helicase C domain-containing protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	A	662	Total	C	N	O	S	0	0
			5311	3375	921	978	37		
3	C	662	Total	C	N	O	S	0	0
			5311	3375	921	978	37		
3	E	662	Total	C	N	O	S	0	0
			5311	3375	921	978	37		

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	843	ARG	HIS	conflict	UNP Q9BYX4
A	944	LYS	ASN	conflict	UNP Q9BYX4
A	946	THR	ALA	conflict	UNP Q9BYX4
C	843	ARG	HIS	conflict	UNP Q9BYX4
C	944	LYS	ASN	conflict	UNP Q9BYX4
C	946	THR	ALA	conflict	UNP Q9BYX4
E	843	ARG	HIS	conflict	UNP Q9BYX4
E	944	LYS	ASN	conflict	UNP Q9BYX4
E	946	THR	ALA	conflict	UNP Q9BYX4

- Molecule 4 is a protein called Tripartite motif-containing protein 65.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	B	180	Total 1408	C 900	N 249	O 254	S 5	2	0
4	D	180	Total 1408	C 900	N 249	O 254	S 5	2	0
4	F	180	Total 1408	C 900	N 249	O 254	S 5	2	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	356	LEU	CYS	conflict	UNP Q6PJ69
B	407	SER	CYS	conflict	UNP Q6PJ69
B	421	SER	CYS	conflict	UNP Q6PJ69
D	356	LEU	CYS	conflict	UNP Q6PJ69
D	407	SER	CYS	conflict	UNP Q6PJ69
D	421	SER	CYS	conflict	UNP Q6PJ69
F	356	LEU	CYS	conflict	UNP Q6PJ69
F	407	SER	CYS	conflict	UNP Q6PJ69
F	421	SER	CYS	conflict	UNP Q6PJ69

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
5	A	1	Total 1	Zn 1	0
5	C	1	Total 1	Zn 1	0
5	E	1	Total 1	Zn 1	0

- Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					AltConf
6	A	1	Total	C	N	O	P	0
			27	10	5	10	2	
6	C	1	Total	C	N	O	P	0
			27	10	5	10	2	
6	E	1	Total	C	N	O	P	0
			27	10	5	10	2	

- Molecule 7 is TETRAFLUOROALUMINATE ION (three-letter code: ALF) (formula:  $\text{AlF}_4$ ).



Mol	Chain	Residues	Atoms			AltConf
7	A	1	Total	Al	F	0
			5	1	4	

Continued on next page...

*Continued from previous page...*


Mol	Chain	Residues	Atoms			AltConf
7	C	1	Total	Al	F	0
			5	1	4	
7	E	1	Total	Al	F	0
			5	1	4	

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
8	A	1	Total	Mg	0
			1	1	
8	C	1	Total	Mg	0
			1	1	
8	E	1	Total	Mg	0
			1	1	









• Molecule 3: Interferon-induced helicase C domain-containing protein 1






• Molecule 3: Interferon-induced helicase C domain-containing protein 1


Chain E: 13% 58% 31% 10%




• Molecule 4: Tripartite motif-containing protein 65

Chain B: 70% 58% 37% 6%






• Molecule 4: Tripartite motif-containing protein 65



• Molecule 4: Tripartite motif-containing protein 65



## 4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=86.9406°, rise=46.8426 Å, axial sym=C1	Depositor
Number of segments used	15306	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{Å}^2$ )	72.8	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.067	Depositor
Minimum map value	-0.028	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	265.20065, 265.20065, 265.20065	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.03594, 1.03594, 1.03594	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, ADP, ALF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	X	0.62	0/1057	1.12	5/1646 (0.3%)
2	Y	0.64	0/1030	1.21	2/1600 (0.1%)
3	A	0.36	0/5394	0.52	3/7248 (0.0%)
3	C	0.36	0/5394	0.52	3/7248 (0.0%)
3	E	0.36	0/5394	0.52	3/7248 (0.0%)
4	B	0.30	0/1460	0.48	0/1996
4	D	0.30	0/1460	0.48	0/1996
4	F	0.29	0/1460	0.46	0/1996
All	All	0.38	0/22649	0.61	16/30978 (0.1%)

There are no bond length outliers.

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	22	C	N3-C4-N4	7.33	123.13	118.00
1	X	22	C	C5-C4-N4	-7.20	115.16	120.20
3	A	314	GLU	CA-CB-CG	6.94	128.67	113.40
3	E	314	GLU	CA-CB-CG	6.93	128.66	113.40
3	C	314	GLU	CA-CB-CG	6.92	128.63	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	944	0	476	48	0
2	Y	925	0	471	30	0
3	A	5311	0	5409	193	0
3	C	5311	0	5409	186	0
3	E	5311	0	5409	188	0
4	B	1408	0	1304	56	0
4	D	1408	0	1304	49	0
4	F	1408	0	1304	54	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
5	E	1	0	0	0	0
6	A	27	0	12	7	0
6	C	27	0	12	6	0
6	E	27	0	12	7	0
7	A	5	0	0	1	0
7	C	5	0	0	1	0
7	E	5	0	0	1	0
8	A	1	0	0	0	0
8	C	1	0	0	0	0
8	E	1	0	0	0	0
All	All	22128	0	21122	783	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 783 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:355:HIS:O	4:D:356:LEU:HD12	1.41	1.18
3:A:1005:GLU:OE1	3:A:1005:GLU:N	2.01	0.92
3:E:1005:GLU:OE1	3:E:1005:GLU:N	2.01	0.92
3:C:1005:GLU:OE1	3:C:1005:GLU:N	2.01	0.91
3:E:458:MET:HE3	3:E:458:MET:HA	1.57	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	644/739 (87%)	605 (94%)	39 (6%)	0	100	100
3	C	644/739 (87%)	606 (94%)	38 (6%)	0	100	100
3	E	644/739 (87%)	605 (94%)	39 (6%)	0	100	100
4	B	178/191 (93%)	173 (97%)	5 (3%)	0	100	100
4	D	178/191 (93%)	173 (97%)	5 (3%)	0	100	100
4	F	178/191 (93%)	173 (97%)	5 (3%)	0	100	100
All	All	2466/2790 (88%)	2335 (95%)	131 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	584/659 (89%)	580 (99%)	4 (1%)	84	90
3	C	584/659 (89%)	580 (99%)	4 (1%)	84	90
3	E	584/659 (89%)	580 (99%)	4 (1%)	84	90
4	B	147/166 (89%)	146 (99%)	1 (1%)	84	90
4	D	147/166 (89%)	147 (100%)	0	100	100
4	F	147/166 (89%)	147 (100%)	0	100	100
All	All	2193/2475 (89%)	2180 (99%)	13 (1%)	86	92

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	784	ASN
3	C	1005	GLU
3	E	1005	GLU
3	E	498	LYS
3	E	784	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	428	GLN
3	E	446	HIS
3	E	835	HIS
3	C	446	HIS
3	C	835	HIS

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	X	43/44 (97%)	1 (2%)	1 (2%)
2	Y	43/44 (97%)	3 (6%)	0
All	All	86/88 (97%)	4 (4%)	1 (1%)

All (4) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	X	31	G
2	Y	14	C
2	Y	28	U
2	Y	29	C

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	X	30	A

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	ALF	E	1103	-	0,4,4	-	-	-		
7	ALF	A	1103	-	0,4,4	-	-	-		
6	ADP	C	1102	8	24,29,29	0.95	1 (4%)	29,45,45	1.51	4 (13%)
6	ADP	A	1102	8	24,29,29	0.95	1 (4%)	29,45,45	1.50	4 (13%)
6	ADP	E	1102	8	24,29,29	0.95	1 (4%)	29,45,45	1.51	4 (13%)
7	ALF	C	1103	-	0,4,4	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADP	A	1102	8	-	4/12/32/32	0/3/3/3
6	ADP	E	1102	8	-	4/12/32/32	0/3/3/3
6	ADP	C	1102	8	-	4/12/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	E	1102	ADP	C5-C4	2.39	1.47	1.40
6	C	1102	ADP	C5-C4	2.38	1.47	1.40
6	A	1102	ADP	C5-C4	2.37	1.47	1.40

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	1102	ADP	PA-O3A-PB	-4.10	118.77	132.83
6	C	1102	ADP	PA-O3A-PB	-4.09	118.78	132.83
6	A	1102	ADP	PA-O3A-PB	-4.09	118.79	132.83
6	E	1102	ADP	C3'-C2'-C1'	3.52	106.28	100.98
6	C	1102	ADP	C3'-C2'-C1'	3.50	106.24	100.98

There are no chirality outliers.

5 of 12 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	A	1102	ADP	C5'-O5'-PA-O2A
6	C	1102	ADP	C5'-O5'-PA-O2A
6	E	1102	ADP	C5'-O5'-PA-O2A
6	A	1102	ADP	PB-O3A-PA-O2A
6	C	1102	ADP	PB-O3A-PA-O2A


There are no ring outliers.

6 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	E	1103	ALF	1	0
7	A	1103	ALF	1	0
6	C	1102	ADP	6	0
6	A	1102	ADP	7	0
6	E	1102	ADP	7	0
7	C	1103	ALF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	A	1
3	C	1
3	E	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	1013:LEU	C	1014:ASP	N	3.05
1	C	1013:LEU	C	1014:ASP	N	3.05
1	E	1013:LEU	C	1014:ASP	N	3.05


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-22370. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)

#### 6.2.1 Primary map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 130



Y Index: 130



Z Index: 143

The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.




## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 136 nm<sup>3</sup>; this corresponds to an approximate mass of 123 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.233 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-22370 and PDB model 7JL2. Per-residue inclusion information can be found in section 3 on page 8.


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 77% of all backbone atoms, 57% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.5737	<div></div> 0.3620
A	<div></div> 0.6313	<div></div> 0.3860
B	<div></div> 0.2599	<div></div> 0.2680
C	<div></div> 0.6132	<div></div> 0.3840
D	<div></div> 0.2365	<div></div> 0.2700
E	<div></div> 0.6204	<div></div> 0.3830
F	<div></div> 0.2547	<div></div> 0.2660
X	<div></div> 0.8803	<div></div> 0.3750
Y	<div></div> 0.8800	<div></div> 0.3840

1.0

0.0

<0.0