



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 02:08 pm BST

PDB ID : 4JN6  
Title : Crystal Structure of the Aldolase-Dehydrogenase Complex from Mycobacterium tuberculosis HRv37  
Authors : Carere, J.; McKenna, S.E.; Kimber, M.S.; Seah, S.Y.K.  
Deposited on : 2013-03-14  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	<div> <div>3%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	C	346	<div> <div>92%</div> <div>6%</div> <div>.</div> </div>
2	B	306	<div> <div>15%</div> <div>85%</div> <div>12%</div> <div>.</div> </div>
2	D	306	<div> <div>18%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxy-2-oxovalerate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	0	0
			2499	1559	449	478	13			
1	C	339	Total	C	N	O	S	0	0	0
			2504	1562	450	479	13			


- Molecule 2 is a protein called Acetaldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	296	Total	C	N	O	S	0	0	0
			2205	1397	372	429	7			
2	D	296	Total	C	N	O	S	0	0	0
			2205	1397	372	429	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	EXPRESSION TAG	UNP P71866
B	-1	SER	-	EXPRESSION TAG	UNP P71866
B	0	HIS	-	EXPRESSION TAG	UNP P71866
D	-2	GLY	-	EXPRESSION TAG	UNP P71866
D	-1	SER	-	EXPRESSION TAG	UNP P71866
D	0	HIS	-	EXPRESSION TAG	UNP P71866

- Molecule 3 is OXALATE ION (three-letter code: OXL) (formula: C<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	2	4		
3	C	1	Total	C	O	0	0
			6	2	4		

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mn	0	0
			1	1		
4	C	1	Total	Mn	0	0
			1	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Na	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	101	Total	O	0	0
			101	101		
6	B	42	Total	O	0	0
			42	42		


*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	132	Total 132	O 132	0	0
6	D	78	Total 78	O 78	0	0



- Molecule 1: 4-hydroxy-2-oxovalerate aldolase





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.69Å 142.69Å 148.17Å 90.00° 95.08° 90.00°	Depositor
Resolution (Å)	45.89 – 1.93 45.89 – 1.93	Depositor EDS
% Data completeness (in resolution range)	98.1 (45.89-1.93) 98.1 (45.89-1.93)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 1.92Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.174 , 0.206 0.177 , 0.208	Depositor DCC
$R_{free}$ test set	5296 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9781	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OXL, NA, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/2536	0.47	0/3425
1	C	0.38	0/2541	0.51	0/3432
2	B	0.28	0/2250	0.48	0/3078
2	D	0.31	0/2250	0.51	0/3078
All	All	0.32	0/9577	0.49	0/13013

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2499	0	2486	17	0
1	C	2504	0	2491	10	0
2	B	2205	0	2211	20	0
2	D	2205	0	2211	16	0
3	A	6	0	0	0	0
3	C	6	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	C	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	101	0	0	2	0
6	B	42	0	0	3	0
6	C	132	0	0	2	0
6	D	78	0	0	6	1
All	All	9781	0	9399	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:265:VAL:O	6:C:565:HOH:O	1.89	0.90
1:A:122:GLN:NE2	6:A:576:HOH:O	2.15	0.79
2:B:49:LYS:NZ	6:B:440:HOH:O	2.24	0.70
1:A:82:ARG:NH1	1:A:133:GLU:OE1	2.27	0.67
2:D:242:GLU:OE1	6:D:455:HOH:O	2.12	0.66
2:D:164:ARG:O	6:D:416:HOH:O	2.13	0.66
2:B:7:VAL:HG12	2:B:71:LEU:HB3	1.78	0.65
2:B:186:ARG:NH2	6:B:417:HOH:O	2.31	0.64
2:D:293:ALA:HA	2:D:297:LEU:HB2	1.78	0.64
2:D:163:THR:N	6:D:475:HOH:O	2.31	0.62
1:A:101:ARG:NH2	1:A:102:ASP:OD1	2.33	0.61
1:A:94:LYS:HB3	1:A:126:LEU:HD22	1.83	0.59
2:D:108:ILE:HG22	2:D:110:PRO:HD2	1.84	0.58
2:B:107:VAL:HG11	2:B:113:LEU:HD13	1.85	0.57
2:B:238:ARG:NH1	2:B:263:GLU:OE2	2.37	0.56
2:B:108:ILE:HG22	2:B:110:PRO:HD2	1.87	0.56
2:D:39:PRO:HG3	2:D:56:HIS:CD2	2.42	0.55
2:B:35:VAL:HG21	2:B:62:LEU:HD22	1.89	0.54
1:C:17:GLY:HA2	1:C:229:GLY:HA2	1.89	0.54
1:A:58:PHE:HE1	2:B:252:SER:HB2	1.73	0.54
2:D:109:PRO:HA	2:D:113:LEU:HB2	1.89	0.54
1:C:94:LYS:HB3	1:C:126:LEU:HD22	1.89	0.53
1:A:236:PRO:HG2	1:A:239:ALA:HB3	1.90	0.53
2:D:21:TYR:OH	2:D:46:ARG:HD2	2.12	0.50
2:D:71:LEU:HD12	2:D:94:ARG:HB2	1.94	0.50
2:D:271:LEU:HD13	2:D:278:LEU:HB2	1.94	0.50
2:D:246:ASP:OD2	6:D:446:HOH:O	2.20	0.49
2:B:17:THR:O	2:B:20:LEU:HB3	2.12	0.49
1:A:295:HIS:O	1:A:299:GLN:HG2	2.13	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:35:VAL:HG21	2:D:62:LEU:HD22	1.95	0.48
1:A:114:CYS:O	1:A:152:GLN:HG3	2.14	0.48
1:A:295:HIS:CD2	1:A:324:GLU:HG2	2.48	0.48
1:C:237:VAL:O	1:C:241:ILE:HG12	2.14	0.47
2:B:109:PRO:HA	2:B:113:LEU:HB2	1.97	0.46
1:A:237:VAL:O	1:A:241:ILE:HG12	2.16	0.46
1:A:142:HIS:HD2	6:A:513:HOH:O	1.99	0.46
2:D:160:GLY:O	6:D:475:HOH:O	2.20	0.45
2:B:15:ILE:HD11	2:B:278:LEU:HD22	1.98	0.45
1:A:254:ASP:HB3	1:A:257:ASP:HB2	1.99	0.45
1:A:281:LEU:CD1	1:C:238:GLU:HG3	2.47	0.44
2:B:57:GLU:HB2	2:B:61:TRP:HB2	1.98	0.44
2:D:103:VAL:N	6:D:438:HOH:O	2.51	0.44
2:B:125:ILE:HG23	2:B:126:THR:H	1.82	0.44
1:A:246:LYS:HE3	6:C:544:HOH:O	2.17	0.44
2:B:20:LEU:HD21	2:B:50:LEU:HD12	1.99	0.43
1:C:114:CYS:O	1:C:152:GLN:HG3	2.18	0.43
2:D:28:TRP:CE2	2:D:294:LYS:HD3	2.54	0.43
2:D:240:LEU:HD21	2:D:263:GLU:HB2	2.01	0.43
1:C:295:HIS:CD2	1:C:324:GLU:HG2	2.54	0.43
2:B:268:GLY:HA2	2:B:271:LEU:O	2.20	0.42
1:A:309:ALA:O	1:A:313:ARG:HG3	2.20	0.42
1:A:281:LEU:HD11	1:C:238:GLU:HG3	2.02	0.41
2:B:199:PRO:HB2	2:B:274:TYR:CG	2.55	0.41
1:C:116:GLU:O	1:C:119:VAL:HG22	2.21	0.41
2:B:223:ILE:O	2:B:227:VAL:HG23	2.21	0.41
2:B:144:GLU:O	2:B:211:PRO:HD3	2.21	0.41
2:B:201:ILE:HD13	2:B:274:TYR:HA	2.03	0.41
1:A:238:GLU:HG3	1:C:281:LEU:CD1	2.51	0.41
2:B:46:ARG:NH1	6:B:435:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:D:448:HOH:O	6:D:448:HOH:O[2_557]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/346 (97%)	331 (98%)	3 (1%)	2 (1%)	25	13
1	C	337/346 (97%)	333 (99%)	2 (1%)	2 (1%)	25	13
2	B	294/306 (96%)	283 (96%)	10 (3%)	1 (0%)	41	32
2	D	294/306 (96%)	285 (97%)	8 (3%)	1 (0%)	41	32
All	All	1261/1304 (97%)	1232 (98%)	23 (2%)	6 (0%)	29	17

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	PHE
1	C	228	PHE
1	A	175	LEU
1	C	175	LEU
2	B	125	ILE
2	D	125	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/257 (98%)	248 (99%)	3 (1%)	71	64
1	C	251/257 (98%)	246 (98%)	5 (2%)	55	42
2	B	231/237 (98%)	227 (98%)	4 (2%)	60	49
2	D	231/237 (98%)	227 (98%)	4 (2%)	60	49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	964/988 (98%)	948 (98%)	16 (2%)	60	49

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	THR
1	A	228	PHE
1	A	301	GLU
2	B	56	HIS
2	B	63	LEU
2	B	79	TYR
2	B	297	LEU
1	C	4	MET
1	C	48	THR
1	C	101	ARG
1	C	228	PHE
1	C	301	GLU
2	D	114	ARG
2	D	226	VAL
2	D	257	LEU
2	D	297	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	OXL	A	401	4	0,5,5	0.00	-	0,6,6	0.00	-
3	OXL	C	401	4	0,5,5	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OXL	A	401	4	-	0/0/4/4	-
3	OXL	C	401	4	-	0/0/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	338/346 (97%)	0.35	11 (3%) 46 54	36, 49, 68, 95	0
1	C	339/346 (97%)	0.12	1 (0%) 94 96	28, 39, 54, 85	0
2	B	296/306 (96%)	1.00	47 (15%) 1 2	48, 69, 104, 120	0
2	D	296/306 (96%)	0.80	56 (18%) 1 1	27, 49, 100, 121	0
All	All	1269/1304 (97%)	0.55	115 (9%) 9 14	27, 49, 93, 121	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	63	LEU	10.4
2	B	61	TRP	10.0
2	B	64	ALA	7.8
2	D	62	LEU	7.5
2	B	37	ILE	6.7
2	B	57	GLU	6.6
2	D	297	LEU	6.5
2	B	66	PRO	6.2
2	D	52	LEU	6.0
2	B	79	TYR	5.9
2	B	60	ASP	5.6
2	D	64	ALA	5.2
1	A	341	GLY	5.2
2	B	88	TYR	5.0
2	B	39	PRO	4.7
2	B	91	ALA	4.6
2	D	39	PRO	4.6
2	D	61	TRP	4.6
2	D	55	THR	4.5
2	B	38	ASP	4.5
2	D	35	VAL	4.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	59	VAL	4.4
2	D	37	ILE	4.2
2	B	67	ASP	4.2
2	D	93	ILE	4.2
2	D	296	THR	4.2
2	B	58	GLY	4.1
2	B	298	VAL	4.0
2	D	41	SER	4.0
2	B	59	VAL	4.0
2	B	68	LYS	3.9
2	D	68	LYS	3.8
2	B	86	PRO	3.8
2	B	297	LEU	3.8
2	B	35	VAL	3.7
2	D	20	LEU	3.6
2	D	298	VAL	3.5
2	D	33	TRP	3.5
2	D	66	PRO	3.5
2	D	90	GLU	3.5
2	D	91	ALA	3.5
2	D	69	PRO	3.5
1	A	340	SER	3.4
2	D	67	ASP	3.4
2	B	296	THR	3.4
2	D	48	ALA	3.4
2	B	56	HIS	3.4
2	B	293	ALA	3.4
2	D	103	VAL	3.3
2	B	90	GLU	3.3
2	D	50	LEU	3.3
1	A	4	MET	3.3
2	B	33	TRP	3.3
2	D	292	ILE	3.2
2	B	40	GLU	3.2
1	A	338	LEU	3.2
2	D	47	ALA	3.1
2	D	63	LEU	3.1
2	D	88	TYR	3.1
2	B	93	ILE	3.1
2	D	34	MET	3.0
2	B	55	THR	3.0
2	B	216	ARG	3.0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	295	GLU	3.0
2	B	114	ARG	3.0
2	D	46	ARG	2.9
2	D	5	ALA	2.9
2	B	62	LEU	2.9
1	A	191	GLU	2.8
2	D	6	LYS	2.8
1	A	88	LEU	2.8
2	D	32	ARG	2.8
2	B	89	ALA	2.8
2	D	44	LEU	2.8
2	D	57	GLU	2.8
2	D	94	ARG	2.7
2	D	45	ALA	2.7
2	D	86	PRO	2.7
2	D	56	HIS	2.6
2	B	214	ALA	2.6
2	D	38	ASP	2.6
2	D	40	GLU	2.5
2	B	213	ASP	2.5
2	D	43	GLY	2.4
2	D	53	GLU	2.4
2	B	65	GLN	2.4
2	D	92	GLY	2.4
2	D	114	ARG	2.4
1	A	192	ASP	2.4
2	B	83	ASP	2.4
2	B	246	ASP	2.4
2	D	42	ASP	2.4
2	B	142	ILE	2.4
2	D	71	LEU	2.4
2	D	36	GLY	2.4
2	D	54	THR	2.3
2	B	220	ALA	2.3
2	B	82	ARG	2.3
2	D	28	TRP	2.3
1	A	170	ALA	2.2
2	D	95	ALA	2.2
1	A	339	ASP	2.2
2	D	58	GLY	2.2
2	D	60	ASP	2.2
2	B	71	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	240	LEU	2.1
1	A	129	GLU	2.1
2	B	190	ILE	2.1
1	C	205	LEU	2.1
2	D	118	ASP	2.1
2	B	245	PHE	2.1
2	D	102	ALA	2.1
2	B	94	ARG	2.0
2	B	84	ALA	2.0
1	A	336	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	OXL	A	401	6/6	0.94	0.21	27,38,43,52	6
3	OXL	C	401	6/6	0.96	0.18	22,31,33,42	6
5	NA	C	402	1/1	0.97	0.08	37,37,37,37	0
4	MN	A	402	1/1	0.98	0.17	41,41,41,41	0
4	MN	C	403	1/1	0.98	0.14	34,34,34,34	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.