



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 01:18 am BST

PDB ID : 4JQA
Title : AKR1C2 complex with mefenamic acid
Authors : Yosaatmadja, Y.; Flanagan, J.U.; Squire, C.J.
Deposited on : 2013-03-20
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>5% • 5%</div> </div> </div>
1	B	331	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>5% •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PO4	B	407	-	X	-	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aldo-keto reductase family 1 member C2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	2	8	0
			2588	1662	443	472	11			
1	B	317	Total	C	N	O	S	3	6	0
			2573	1651	446	465	11			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	324	LEU	-	EXPRESSION TAG	UNP P52895
A	325	GLU	-	EXPRESSION TAG	UNP P52895
A	326	HIS	-	EXPRESSION TAG	UNP P52895
A	327	HIS	-	EXPRESSION TAG	UNP P52895
A	328	HIS	-	EXPRESSION TAG	UNP P52895
A	329	HIS	-	EXPRESSION TAG	UNP P52895
A	330	HIS	-	EXPRESSION TAG	UNP P52895
A	331	HIS	-	EXPRESSION TAG	UNP P52895
B	324	LEU	-	EXPRESSION TAG	UNP P52895
B	325	GLU	-	EXPRESSION TAG	UNP P52895
B	326	HIS	-	EXPRESSION TAG	UNP P52895
B	327	HIS	-	EXPRESSION TAG	UNP P52895
B	328	HIS	-	EXPRESSION TAG	UNP P52895
B	329	HIS	-	EXPRESSION TAG	UNP P52895
B	330	HIS	-	EXPRESSION TAG	UNP P52895
B	331	HIS	-	EXPRESSION TAG	UNP P52895

- Molecule 2 is 2-[(2,3-DIMETHYLPHENYL)AMINO]BENZOIC ACID (three-letter code: ID8) (formula: C₁₅H₁₅NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			18	15	1	2		
2	B	1	Total	C	N	O	0	0
			18	15	1	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			10	4	6		

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	255	Total 255	O 255	0	0
8	B	246	Total 246	O 246	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldo-keto reductase family 1 member C2



- Molecule 1: Aldo-keto reductase family 1 member C2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	144.98Å 144.98Å 201.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.90 – 1.45 19.90 – 1.45	Depositor EDS
% Data completeness (in resolution range)	96.5 (19.90-1.45) 96.5 (19.90-1.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.144 , 0.184 0.153 , 0.185	Depositor DCC
R_{free} test set	6934 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5826	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ID8, NA, PO4, EDO, TLA, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	1/2657 (0.0%)	1.07	13/3599 (0.4%)
1	B	1.05	4/2641 (0.2%)	1.08	12/3577 (0.3%)
All	All	1.04	5/5298 (0.1%)	1.07	25/7176 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	291	GLU	CG-CD	5.78	1.60	1.51
1	B	171	ARG	CZ-NH1	5.44	1.40	1.33
1	B	110	TYR	CE1-CZ	-5.20	1.31	1.38
1	B	46	PHE	CG-CD1	5.13	1.46	1.38
1	A	101	ARG	CZ-NH1	5.04	1.39	1.33

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	ARG	NE-CZ-NH1	10.93	125.76	120.30
1	B	171	ARG	NE-CZ-NH1	10.92	125.76	120.30
1	A	101	ARG	NE-CZ-NH2	-10.88	114.86	120.30
1	A	200	ARG	NE-CZ-NH2	-10.16	115.22	120.30
1	A	200	ARG	NE-CZ-NH1	9.63	125.11	120.30
1	B	171	ARG	NE-CZ-NH2	-8.69	115.96	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	112	ASP	CB-CG-OD2	-7.82	111.26	118.30
1	A	276	ARG	NE-CZ-NH1	7.23	123.92	120.30
1	A	263	ARG	NE-CZ-NH2	-7.19	116.71	120.30
1	B	200	ARG	NE-CZ-NH1	6.61	123.60	120.30
1	B	200	ARG	NE-CZ-NH2	-6.47	117.06	120.30
1	A	301	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	B	91	ARG	NE-CZ-NH1	-6.35	117.12	120.30
1	B	96	ARG	NE-CZ-NH2	-6.26	117.17	120.30
1	A	263	ARG	NE-CZ-NH1	6.14	123.37	120.30
1	A	171	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	B	319	PHE	CB-CG-CD1	5.81	124.87	120.80
1	B	71	ASP	CB-CG-OD1	5.73	123.46	118.30
1	A	278	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	168	PHE	CB-CG-CD1	5.49	124.64	120.80
1	B	301[A]	ARG	NE-CZ-NH1	-5.33	117.63	120.30
1	B	301[B]	ARG	NE-CZ-NH1	-5.33	117.63	120.30
1	A	168	PHE	CB-CG-CD2	-5.32	117.08	120.80
1	B	50	ASP	CB-CG-OD1	-5.30	113.53	118.30
1	B	156	ASP	CB-CG-OD2	-5.14	113.68	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	301[A]	ARG	Mainchain
1	B	301[B]	ARG	Mainchain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2588	0	2604	12	0
1	B	2573	0	2595	10	0
2	A	18	0	14	1	0
2	B	18	0	14	1	0
3	A	48	0	24	3	0
3	B	48	0	25	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	4	0	6	0	0
4	B	12	0	18	0	0
5	A	1	0	0	0	0
6	B	10	0	5	0	0
7	B	5	0	0	0	0
8	A	255	0	0	4	0
8	B	246	0	0	2	0
All	All	5826	0	5305	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:402:NAP:O2B	3:B:402:NAP:C2B	1.71	1.37
3:B:402:NAP:P2B	3:B:402:NAP:C2B	2.74	0.76
1:A:301:ARG:HG3	1:A:303:VAL:HG23	1.74	0.70
1:A:222:HIS:CE1	1:A:270:LYS:HD2	2.31	0.65
1:A:200:ARG:HD3	1:A:200:ARG:O	1.99	0.63
1:A:291:GLU:HG2	8:A:733:HOH:O	1.98	0.63
3:B:402:NAP:H2B	3:B:402:NAP:P2B	2.41	0.59
3:B:402:NAP:O2B	3:B:402:NAP:H2B	1.93	0.59
1:B:270:LYS:O	3:B:402:NAP:H8A	2.04	0.57
1:A:270:LYS:O	3:A:402:NAP:H8A	2.06	0.55
1:B:301[A]:ARG:HB3	1:B:303:VAL:HG23	1.92	0.52
1:B:221:SER:C	3:B:402:NAP:H52A	2.31	0.51
2:A:401:ID8:H12	3:A:402:NAP:O7N	2.10	0.51
1:A:66:ARG:HD2	8:A:723:HOH:O	2.11	0.51
1:B:196:TYR:CD1	1:B:301[B]:ARG:HD2	2.45	0.50
2:B:401:ID8:H12	3:B:402:NAP:O7N	2.11	0.50
1:B:196:TYR:CG	1:B:301[B]:ARG:HD3	2.48	0.48
1:A:301:ARG:HG3	1:A:303:VAL:CG2	2.44	0.46
1:B:271:SER:OG	3:B:402:NAP:O3X	2.22	0.46
1:A:291:GLU:HG3	8:A:543:HOH:O	2.16	0.46
1:B:133:GLU:HG2	8:B:697:HOH:O	2.17	0.45
1:B:301[B]:ARG:HB3	1:B:303:VAL:HG23	1.98	0.44
1:A:200:ARG:HD2	1:A:204:ASP:OD2	2.18	0.43
1:B:200:ARG:HD2	8:B:572:HOH:O	2.19	0.42
1:A:248:HIS:HE1	8:A:715:HOH:O	2.02	0.41
1:A:58[B]:GLU:HG3	1:A:105:ASN:CB	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190:GLN:OE1	3:B:402:NAP:H2N	2.21	0.41
1:A:190:GLN:OE1	3:A:402:NAP:H2N	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/331 (97%)	315 (98%)	7 (2%)	0	100	100
1	B	321/331 (97%)	316 (98%)	5 (2%)	0	100	100
All	All	643/662 (97%)	631 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/293 (97%)	284 (100%)	1 (0%)	91	80
1	B	282/293 (96%)	279 (99%)	3 (1%)	73	48
All	All	567/586 (97%)	563 (99%)	4 (1%)	88	65

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	222	HIS
1	B	36	GLU
1	B	301[A]	ARG
1	B	301[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ID8	A	401	-	17,19,19	2.62	6 (35%)	22,26,26	1.67	3 (13%)
4	EDO	B	406	-	3,3,3	0.95	0	2,2,2	0.35	0
3	NAP	A	402	5	45,52,52	1.51	3 (6%)	56,80,80	1.75	12 (21%)
3	NAP	B	402	-	45,52,52	2.38	15 (33%)	56,80,80	2.14	18 (32%)
4	EDO	B	405	-	3,3,3	0.94	0	2,2,2	0.96	0
6	TLA	B	403	-	3,9,9	0.67	0	6,12,12	2.13	1 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ID8	B	401	-	17,19,19	1.83	4 (23%)	22,26,26	1.61	5 (22%)
4	EDO	B	404	-	3,3,3	0.68	0	2,2,2	0.16	0
7	PO4	B	407	-	4,4,4	2.94	2 (50%)	6,6,6	1.80	2 (33%)
4	EDO	A	403	-	3,3,3	0.71	0	2,2,2	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ID8	A	401	-	-	0/4/8/8	0/2/2/2
4	EDO	B	406	-	-	0/1/1/1	-
3	NAP	A	402	5	-	2/31/67/67	0/5/5/5
3	NAP	B	402	-	-	6/31/67/67	0/5/5/5
4	EDO	B	405	-	-	0/1/1/1	-
6	TLA	B	403	-	-	0/4/12/12	-
2	ID8	B	401	-	-	0/4/8/8	0/2/2/2
4	EDO	B	404	-	-	0/1/1/1	-
4	EDO	A	403	-	-	0/1/1/1	-

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	NAP	C2A-N3A	8.57	1.45	1.32
3	B	402	NAP	O2B-C2B	7.46	1.71	1.44
2	A	401	ID8	C17-C1	-7.09	1.36	1.51
2	B	401	ID8	C17-C1	-5.03	1.41	1.51
3	A	402	NAP	C3N-C7N	-4.84	1.43	1.50
7	B	407	PO4	P-O3	4.63	1.68	1.54
3	A	402	NAP	C2A-N3A	4.62	1.39	1.32
2	A	401	ID8	C13-C14	-4.41	1.43	1.47
3	B	402	NAP	O4D-C1D	4.15	1.46	1.41
2	A	401	ID8	C18-C2	-3.73	1.43	1.51
7	B	407	PO4	P-O2	-3.46	1.44	1.54
2	B	401	ID8	C18-C2	-3.36	1.44	1.51
3	B	402	NAP	C5B-C4B	3.23	1.61	1.51
3	B	402	NAP	C5N-C4N	3.06	1.45	1.38
2	A	401	ID8	C13-C8	2.98	1.44	1.40
3	B	402	NAP	C3N-C7N	-2.85	1.46	1.50
2	B	401	ID8	C13-C14	-2.84	1.44	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	ID8	C11-C12	2.82	1.44	1.38
3	A	402	NAP	O3B-C3B	2.81	1.49	1.43
2	A	401	ID8	C12-C13	2.78	1.44	1.40
3	B	402	NAP	C2D-C1D	-2.76	1.49	1.53
3	B	402	NAP	PA-O1A	-2.71	1.41	1.50
3	B	402	NAP	C3B-C2B	2.56	1.58	1.52
3	B	402	NAP	C3D-C4D	-2.39	1.46	1.53
3	B	402	NAP	O4B-C1B	2.31	1.44	1.41
2	B	401	ID8	C10-C9	-2.28	1.34	1.38
3	B	402	NAP	P2B-O2X	-2.24	1.46	1.54
3	B	402	NAP	PN-O5D	-2.24	1.50	1.59
3	B	402	NAP	PA-O2A	-2.22	1.44	1.55
3	B	402	NAP	C5A-C4A	-2.11	1.35	1.40

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	NAP	O2B-P2B-O1X	-5.78	87.09	109.39
3	A	402	NAP	O4B-C4B-C3B	5.45	115.89	105.11
6	B	403	TLA	O2-C2-C1	-4.77	99.62	111.10
3	B	402	NAP	O2B-C2B-C3B	4.68	128.66	111.68
3	A	402	NAP	C4A-C5A-N7A	-4.56	104.64	109.40
3	B	402	NAP	N3A-C2A-N1A	-4.33	121.91	128.68
3	B	402	NAP	O2A-PA-O1A	4.09	132.46	112.24
3	B	402	NAP	O3X-P2B-O2B	3.94	123.64	105.99
3	B	402	NAP	O2N-PN-O5D	3.80	125.39	107.75
3	B	402	NAP	O4B-C4B-C3B	-3.58	98.04	105.11
7	B	407	PO4	O3-P-O1	-3.37	98.56	110.89
3	B	402	NAP	C4A-C5A-N7A	-3.32	105.94	109.40
2	B	401	ID8	C10-C11-C12	-3.32	115.14	120.19
2	A	401	ID8	C18-C2-C1	3.31	126.33	121.17
3	B	402	NAP	PN-O3-PA	3.30	144.14	132.83
3	B	402	NAP	O2N-PN-O1N	3.19	128.01	112.24
3	B	402	NAP	C2D-C3D-C4D	-3.19	96.44	102.64
3	B	402	NAP	C3B-C2B-C1B	-3.17	96.93	102.89
2	B	401	ID8	C10-C9-C8	3.08	125.03	118.62
3	A	402	NAP	C2B-C3B-C4B	-3.08	95.30	101.99
3	A	402	NAP	O7N-C7N-N7N	-3.08	118.20	122.58
3	A	402	NAP	O2N-PN-O1N	3.06	127.35	112.24
3	A	402	NAP	C3N-C2N-N1N	3.04	123.40	120.43
2	A	401	ID8	C10-C9-C8	3.02	124.91	118.62
2	B	401	ID8	C11-C12-C13	2.96	125.32	120.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	ID8	C6-C1-C2	2.92	121.42	118.49
3	B	402	NAP	C2B-C3B-C4B	2.82	108.11	101.99
3	A	402	NAP	C2N-C3N-C4N	-2.75	115.14	118.26
3	A	402	NAP	C6N-N1N-C2N	-2.73	119.48	121.97
2	B	401	ID8	C9-C8-C13	-2.73	116.54	119.98
3	A	402	NAP	C2A-N1A-C6A	-2.72	114.11	118.75
3	B	402	NAP	C3D-C2D-C1D	2.62	104.92	100.98
3	A	402	NAP	O7N-C7N-C3N	2.54	122.67	119.63
2	B	401	ID8	C18-C2-C3	-2.51	115.41	120.31
3	A	402	NAP	C5N-C4N-C3N	2.37	123.15	120.34
3	A	402	NAP	O4D-C1D-C2D	2.34	110.35	106.93
7	B	407	PO4	O4-P-O2	2.29	115.31	107.97
3	B	402	NAP	C3N-C2N-N1N	-2.26	118.22	120.43
3	B	402	NAP	C6N-C5N-C4N	-2.14	116.33	119.44
3	B	402	NAP	O4D-C1D-C2D	-2.12	103.83	106.93
3	B	402	NAP	O3X-P2B-O1X	2.03	118.61	110.68

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	NAP	PA-O3-PN-O5D
3	B	402	NAP	PA-O3-PN-O5D
3	A	402	NAP	C4D-C5D-O5D-PN
3	B	402	NAP	C2B-O2B-P2B-O1X
3	B	402	NAP	C3B-C4B-C5B-O5B
3	B	402	NAP	PN-O3-PA-O1A
3	B	402	NAP	C4D-C5D-O5D-PN
3	B	402	NAP	C5D-O5D-PN-O2N


There are no ring outliers.


4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	ID8	1	0
3	A	402	NAP	3	0
3	B	402	NAP	9	0
2	B	401	ID8	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/331 (95%)	0.13	9 (2%) 53 55	13, 19, 33, 46	9 (2%)
1	B	317/331 (95%)	0.19	8 (2%) 57 60	10, 17, 34, 51	8 (2%)
All	All	633/662 (95%)	0.16	17 (2%) 54 56	10, 18, 33, 51	17 (2%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	ASN	3.8
1	B	3	SER	3.4
1	A	303	VAL	3.3
1	B	301[A]	ARG	3.0
1	B	49[A]	ILE	2.9
1	A	226	PRO	2.8
1	B	28	GLU	2.6
1	A	230	PRO	2.6
1	A	27	ALA	2.6
1	B	303	VAL	2.4
1	A	225	GLU	2.3
1	A	243	ALA	2.3
1	A	180	PRO	2.3
1	B	30	PRO	2.2
1	A	249	LYS	2.1
1	B	163	ILE	2.0
1	B	249	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	PO4	B	407	5/5	0.83	0.22	31,39,60,65	5
6	TLA	B	403	10/10	0.86	0.14	23,27,36,47	0
4	EDO	B	406	4/4	0.88	0.14	27,29,30,32	0
2	ID8	A	401	18/18	0.90	0.14	16,24,33,39	0
5	NA	A	404	1/1	0.91	0.16	42,42,42,42	0
4	EDO	B	405	4/4	0.93	0.12	21,25,25,28	0
2	ID8	B	401	18/18	0.93	0.13	15,25,33,43	0
3	NAP	B	402	48/48	0.96	0.09	12,20,34,41	0
4	EDO	B	404	4/4	0.96	0.10	15,15,18,21	0
3	NAP	A	402	48/48	0.97	0.07	12,19,26,28	0
4	EDO	A	403	4/4	0.97	0.08	14,16,19,20	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.