



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 11:54 am BST

PDB ID : 3JRU
Title : Crystal structure of Leucyl Aminopeptidase (pepA) from
Xoo0834, Xanthomonas oryzae pv. oryzae KACC10331
Authors : Natarajan, S.; Huynh, K.-H.; Kang, L.W.
Deposited on : 2009-09-08
Resolution : 2.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>12%</div> <div>.</div> </div> </div>
1	B	490	<div> <div></div> <div>85%</div> <div>12%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CO3	A	491	-	-	X	-
2	CO3	B	491	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable cytosol aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	490	Total	C	N	O	S	0	0	0
			3587	2257	625	690	15			
1	A	490	Total	C	N	O	S	0	0	0
			3587	2257	625	690	15			

- Molecule 2 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	1	3		
2	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total 2	Zn 2	0	0
3	A	2	Total 2	Zn 2	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Ca 1	0	0
4	A	1	Total 1	Ca 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	420	Total 420	O 420	0	0
5	A	276	Total 276	O 276	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable cytosol aminopeptidase



- Molecule 1: Probable cytosol aminopeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	152.13Å 152.13Å 152.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.02 – 2.60 19.02 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.02-2.60) 100.0 (19.02-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.98 (at 2.59Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.146 , 0.215 0.146 , 0.214	Depositor DCC
R_{free} test set	999 reflections (2.74%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.032 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7884	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, CO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	2/3647 (0.1%)	0.80	0/4964
1	B	0.87	1/3647 (0.0%)	0.87	9/4964 (0.2%)
All	All	0.84	3/7294 (0.0%)	0.84	9/9928 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	253	ASP	CB-CG	6.13	1.64	1.51
1	A	9	GLN	CG-CD	6.11	1.65	1.51
1	B	325	ARG	CZ-NH1	-5.26	1.26	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	488	ARG	NE-CZ-NH2	-6.73	116.93	120.30
1	B	415	TRP	CA-CB-CG	6.14	125.36	113.70
1	B	183	ARG	NE-CZ-NH1	-6.06	117.27	120.30
1	B	164	ASP	CB-CG-OD2	5.69	123.42	118.30
1	B	408	ARG	NE-CZ-NH1	5.49	123.04	120.30
1	B	242	ARG	NE-CZ-NH1	5.18	122.89	120.30
1	B	240	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	B	488	ARG	CG-CD-NE	-5.10	101.08	111.80
1	B	408	ARG	CD-NE-CZ	5.05	130.67	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3587	0	3595	48	0
1	B	3587	0	3595	60	0
2	A	4	0	0	2	0
2	B	4	0	0	3	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	276	0	0	8	3
5	B	420	0	0	7	4
All	All	7884	0	7190	107	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:488:ARG:HH11	1:A:488:ARG:HG3	1.19	1.01
1:B:408:ARG:HD2	5:A:621:HOH:O	1.65	0.95
1:B:408:ARG:CD	5:A:621:HOH:O	2.14	0.94
1:B:129:ARG:HD3	5:B:604:HOH:O	1.72	0.88
1:A:470:ARG:H	1:A:470:ARG:HH11	1.19	0.86
1:B:291:THR:HG21	1:B:477:PRO:HD2	1.56	0.85
1:B:287:CYS:O	1:B:291:THR:HB	1.77	0.84
1:A:488:ARG:NH1	1:A:488:ARG:HG3	1.89	0.82
1:B:244:ILE:HD11	1:B:246:LEU:HG	1.62	0.81
1:B:272:ASN:HD22	1:B:343:THR:H	1.30	0.79
1:A:30:LEU:HD21	1:A:49:LEU:HD11	1.67	0.75
1:A:272:ASN:HD22	1:A:343:THR:H	1.34	0.73
1:B:257:TYR:CE1	1:B:488:ARG:HG3	2.23	0.73
1:B:291:THR:CG2	1:B:477:PRO:HD2	2.19	0.72
1:B:342:ASN:HD22	1:B:344:ASP:H	1.36	0.72
1:A:257:TYR:CZ	1:A:488:ARG:HG2	2.28	0.68
1:B:4:GLN:NE2	5:B:649:HOH:O	2.25	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:408:ARG:H	1:B:408:ARG:NH1	1.93	0.66
1:B:272:ASN:ND2	1:B:343:THR:H	1.93	0.65
1:A:209:PHE:CZ	1:A:301:LYS:HE2	2.32	0.65
1:B:342:ASN:ND2	1:B:344:ASP:H	1.94	0.65
1:B:244:ILE:HD11	1:B:246:LEU:CG	2.27	0.64
1:A:1:MET:HA	1:A:180:GLU:OE1	1.98	0.64
1:B:257:TYR:CZ	1:B:488:ARG:HG3	2.32	0.64
1:B:434:GLY:N	5:B:663:HOH:O	2.31	0.62
1:B:408:ARG:HD3	5:A:621:HOH:O	1.91	0.62
1:A:9:GLN:NE2	1:A:164:ASP:OD2	2.33	0.61
1:A:272:ASN:ND2	1:A:343:THR:H	1.97	0.61
1:A:30:LEU:CD2	1:A:49:LEU:HD11	2.31	0.61
1:B:141:ARG:HD2	1:B:151:VAL:HG21	1.84	0.59
1:A:488:ARG:HH11	1:A:488:ARG:CG	2.01	0.59
1:A:257:TYR:CE1	1:A:488:ARG:HG2	2.37	0.58
1:A:229:SER:HA	1:A:330:ILE:HD12	1.85	0.57
1:B:408:ARG:HH11	1:B:408:ARG:H	1.52	0.57
1:A:89:ALA:HB3	1:A:90:PRO:HD3	1.85	0.57
1:A:61:THR:HG23	5:A:581:HOH:O	2.04	0.57
1:B:4:GLN:HG3	1:B:158:THR:HB	1.87	0.56
1:B:432:ILE:HG13	5:B:663:HOH:O	2.05	0.56
1:B:260:VAL:O	1:B:368:ASP:HA	2.06	0.56
1:B:410:TRP:CE2	1:A:413:PRO:HD3	2.41	0.56
1:B:419:GLN:HE21	1:B:448:ARG:HH21	1.55	0.55
1:A:460:ILE:HD12	1:A:464:ALA:HB2	1.88	0.55
1:B:419:GLN:NE2	1:B:448:ARG:HE	2.05	0.54
1:A:325:ARG:HD2	1:A:328:ASP:OD1	2.08	0.53
1:B:30:LEU:HD21	1:B:49:LEU:HD21	1.91	0.53
1:A:258:VAL:HG23	1:A:363:PRO:HB3	1.90	0.52
1:B:342:ASN:HD22	1:B:344:ASP:N	2.05	0.52
1:A:217:LEU:HA	1:A:221:GLN:OE1	2.10	0.52
1:B:264:ILE:HA	1:B:315:VAL:O	2.10	0.52
1:B:372:LEU:HB3	1:B:373:THR:HG23	1.93	0.50
1:A:255:ARG:NH1	1:A:362:ASN:OD1	2.44	0.50
1:B:383:GLN:H	1:B:383:GLN:HE21	1.60	0.50
1:A:22:VAL:HG11	1:A:37:LEU:HD22	1.94	0.50
1:B:151:VAL:HG22	1:B:152:ASP:N	2.26	0.49
1:B:209:PHE:CZ	1:B:301:LYS:HE2	2.48	0.49
1:B:342:ASN:ND2	1:B:344:ASP:HB3	2.28	0.49
1:B:383:GLN:NE2	1:B:383:GLN:H	2.11	0.49
1:B:28:LYS:HG2	1:B:49:LEU:HD12	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:LEU:HD21	1:A:130:GLN:HG2	1.95	0.49
1:B:91:TYR:CZ	1:B:95:ILE:HD11	2.48	0.49
1:B:348:ARG:NH2	2:B:491:CO3:O3	2.40	0.49
1:B:419:GLN:HE22	1:B:448:ARG:HE	1.60	0.49
1:B:342:ASN:HD21	1:B:344:ASP:HB3	1.78	0.48
1:B:35:GLN:NE2	5:B:694:HOH:O	2.47	0.48
1:B:348:ARG:N	2:B:491:CO3:O1	2.43	0.48
1:A:264:ILE:HD11	1:A:350:ILE:HD13	1.95	0.48
1:A:274:LYS:HB2	1:A:279:ILE:HD12	1.96	0.47
1:A:209:PHE:HZ	1:A:301:LYS:HE2	1.78	0.47
1:B:99:THR:OG1	1:B:135:SER:HB2	2.16	0.46
1:A:105:GLY:O	1:A:154:THR:HB	2.14	0.46
1:B:1:MET:HA	1:B:180:GLU:OE2	2.16	0.46
1:B:247:LYS:HD2	5:B:620:HOH:O	2.15	0.46
1:A:372:LEU:HB3	1:A:373:THR:HG23	1.98	0.46
1:A:165:ASP:HB2	5:A:731:HOH:O	2.16	0.45
1:A:348:ARG:NE	2:A:491:CO3:O2	2.46	0.45
1:B:51:ARG:HD2	1:B:53:ASP:OD2	2.17	0.45
1:A:366:LEU:O	1:A:455:TRP:HA	2.17	0.45
1:B:326:PRO:O	1:B:327:SER:CB	2.65	0.45
1:A:264:ILE:CD1	1:A:350:ILE:HD13	2.47	0.45
1:B:343:THR:O	1:B:346:GLU:HB3	2.17	0.44
1:A:391:HIS:HD2	5:A:744:HOH:O	1.99	0.44
1:B:126:TRP:CZ2	1:B:130:GLN:HG3	2.51	0.44
1:B:1:MET:SD	1:B:177:GLU:HG3	2.58	0.44
1:B:18:ASP:HB2	1:B:107:VAL:HG13	1.99	0.44
1:B:408:ARG:HH11	1:B:408:ARG:N	2.16	0.43
1:A:30:LEU:HD21	1:A:49:LEU:CD1	2.44	0.43
1:B:372:LEU:HD22	2:B:491:CO3:C	2.48	0.43
1:A:470:ARG:NH1	1:A:470:ARG:H	1.99	0.43
1:A:126:TRP:O	1:A:130:GLN:HB2	2.18	0.43
1:B:157:THR:OG1	1:B:158:THR:HG22	2.17	0.43
1:B:44:ARG:HG2	1:B:67:LEU:HD22	2.00	0.43
1:A:258:VAL:CG2	1:A:363:PRO:HB3	2.48	0.43
1:B:480:LEU:HD23	1:B:480:LEU:C	2.39	0.43
1:B:261:GLY:O	1:B:313:PRO:HD2	2.19	0.43
1:B:372:LEU:HA	1:B:372:LEU:HD23	1.86	0.43
1:B:25:PHE:O	1:B:56:GLY:HA3	2.18	0.42
1:A:11:ALA:HA	1:A:12:PRO:HD2	1.75	0.42
1:A:455:TRP:CZ2	1:A:457:HIS:HB2	2.55	0.42
1:A:463:VAL:HG22	1:A:476:ARG:NH1	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:408:ARG:NH2	5:A:665:HOH:O	2.44	0.41
1:A:348:ARG:NH2	2:A:491:CO3:O2	2.49	0.41
1:B:488:ARG:NH2	5:B:908:HOH:O	2.52	0.41
1:A:150:LYS:HA	5:A:509:HOH:O	2.21	0.41
1:A:45:LEU:HD23	1:A:45:LEU:HA	1.81	0.41
1:A:27:ASP:O	1:A:28:LYS:HB2	2.21	0.41
1:A:88:VAL:O	1:A:92:LEU:HG	2.22	0.40
1:A:343:THR:O	1:A:346:GLU:HB3	2.21	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:669:HOH:O	5:B:880:HOH:O[9_555]	1.98	0.22
5:B:574:HOH:O	5:B:635:HOH:O[11_455]	2.04	0.16
5:A:613:HOH:O	5:A:613:HOH:O[8_555]	2.10	0.10
5:A:611:HOH:O	5:A:615:HOH:O[8_555]	2.12	0.08
5:B:573:HOH:O	5:B:619:HOH:O[11_455]	2.15	0.05
5:B:574:HOH:O	5:B:619:HOH:O[11_455]	2.19	0.01
5:A:611:HOH:O	5:A:613:HOH:O[8_555]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	488/490 (100%)	472 (97%)	15 (3%)	1 (0%)	47	71
1	B	488/490 (100%)	468 (96%)	20 (4%)	0	100	100
All	All	976/980 (100%)	940 (96%)	35 (4%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	252	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/357 (100%)	334 (94%)	23 (6%)	17	35
1	B	357/357 (100%)	341 (96%)	16 (4%)	27	52
All	All	714/714 (100%)	675 (94%)	39 (6%)	21	43

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	4	GLN
1	B	49	LEU
1	B	77	VAL
1	B	141	ARG
1	B	158	THR
1	B	244	ILE
1	B	247	LYS
1	B	280	GLU
1	B	291	THR
1	B	324	TYR
1	B	342	ASN
1	B	372	LEU
1	B	383	GLN
1	B	408	ARG
1	B	470	ARG
1	B	488	ARG
1	A	8	ASN
1	A	14	SER
1	A	28	LYS
1	A	37	LEU
1	A	39	SER
1	A	51	ARG
1	A	114	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	120	LYS
1	A	130	GLN
1	A	132	VAL
1	A	150	LYS
1	A	213	GLU
1	A	253	ASP
1	A	322	ASN
1	A	324	TYR
1	A	325	ARG
1	A	346	GLU
1	A	362	ASN
1	A	372	LEU
1	A	421	LEU
1	A	467	GLU
1	A	470	ARG
1	A	488	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	272	ASN
1	B	342	ASN
1	B	383	GLN
1	B	419	GLN
1	A	8	ASN
1	A	249	ASN
1	A	272	ASN
1	A	322	ASN
1	A	391	HIS
1	A	404	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CO3	B	491	-	0,3,3	0.00	-	0,3,3	0.00	-
2	CO3	A	491	-	0,3,3	0.00	-	0,3,3	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	491	CO3	3	0
2	A	491	CO3	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/490 (100%)	-0.76	4 (0%) 86 84	18, 26, 43, 65	0
1	B	490/490 (100%)	-0.92	2 (0%) 92 91	12, 20, 37, 54	0
All	All	980/980 (100%)	-0.84	6 (0%) 89 88	12, 24, 41, 65	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	151	VAL	3.1
1	A	151	VAL	3.0
1	A	153	GLU	2.8
1	A	253	ASP	2.7
1	B	150	LYS	2.1
1	A	470	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CO3	A	491	4/4	0.96	0.21	34,34,35,35	0
3	ZN	B	631	1/1	0.97	0.07	66,66,66,66	0
3	ZN	A	630	1/1	0.97	0.06	72,72,72,72	0
4	CA	B	634	1/1	0.98	0.03	23,23,23,23	1
3	ZN	A	629	1/1	0.98	0.03	44,44,44,44	0
3	ZN	B	632	1/1	0.98	0.05	39,39,39,39	0
2	CO3	B	491	4/4	0.99	0.16	27,28,28,28	0
4	CA	A	633	1/1	1.00	0.05	25,25,25,25	1

6.5 Other polymers [i](#)

There are no such residues in this entry.