



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 07:24 AM EDT

PDB ID : 1JTM
Title : Alternative Structures of a Sequence Extended T4 Lysozyme Show that the Highly Conserved Beta-Sheet has Weak Intrinsic Folding Propensity
Authors : Sagermann, M.; Matthews, B.W.
Deposited on : 2001-08-21
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BME	A	200	-	X	-	X

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	170	Total	C	N	O	S	0	0	0
			1338	842	245	246	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	THR	CYS	engineered mutation	UNP P00720
A	97	ALA	CYS	engineered mutation	UNP P00720
A	165	GLU	-	insertion	UNP P00720
A	166	THR	-	insertion	UNP P00720
A	167	TYR	-	insertion	UNP P00720
A	168	TYR	-	insertion	UNP P00720
A	169	THR	-	insertion	UNP P00720
A	170	ILE	-	insertion	UNP P00720
A	171	GLY	-	insertion	UNP P00720
A	172	ILE	-	insertion	UNP P00720
A	173	GLY	-	insertion	UNP P00720
A	174	HIS	-	insertion	UNP P00720
A	175	LEU	-	insertion	UNP P00720
A	176	LEU	-	insertion	UNP P00720
A	177	THR	-	insertion	UNP P00720
A	178	LYS	-	insertion	UNP P00720

- Molecule 2 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
2	A	1	4	2	1	1	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LYSOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	60.97Å 60.97Å 97.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	17.00 – 1.90 19.96 – 1.90	Depositor EDS
% Data completeness (in resolution range)	(Not available) (17.00-1.90) 97.1 (19.96-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.90Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.214 , 0.265 0.216 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.044 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1342	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	6/1358 (0.4%)	1.10	20/1828 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	2	0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	128	GLU	CD-OE2	6.66	1.32	1.25
1	A	11	GLU	CD-OE2	6.09	1.32	1.25
1	A	22	GLU	CD-OE2	5.86	1.32	1.25
1	A	108	GLU	CD-OE2	5.78	1.32	1.25
1	A	64	GLU	CD-OE2	5.37	1.31	1.25
1	A	5	GLU	CD-OE2	5.20	1.31	1.25

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	89	ASP	CB-CG-OD2	-7.32	111.71	118.30
1	A	127	ASP	CB-CG-OD2	-6.88	112.11	118.30
1	A	127	ASP	CB-CG-OD1	6.62	124.26	118.30
1	A	72	ASP	CB-CG-OD2	-6.46	112.49	118.30
1	A	47	ASP	CB-CG-OD2	-6.29	112.64	118.30
1	A	10	ASP	CB-CG-OD2	-6.25	112.67	118.30
1	A	61	ASP	CB-CG-OD2	-6.03	112.87	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	ASP	CB-CG-OD2	-5.81	113.07	118.30
1	A	89	ASP	CB-CG-OD1	5.70	123.43	118.30
1	A	175	LEU	N-CA-C	5.70	126.38	111.00
1	A	159	ASP	CB-CG-OD2	-5.69	113.18	118.30
1	A	20	ASP	CB-CG-OD1	5.52	123.27	118.30
1	A	70	ASP	CB-CG-OD2	-5.40	113.44	118.30
1	A	72	ASP	CB-CG-OD1	5.36	123.12	118.30
1	A	70	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	61	ASP	CB-CG-OD1	5.24	123.02	118.30
1	A	20	ASP	CB-CG-OD2	-5.22	113.60	118.30
1	A	10	ASP	CB-CG-OD1	5.19	122.97	118.30
1	A	159	ASP	CB-CG-OD1	5.16	122.94	118.30
1	A	47	ASP	CB-CG-OD1	5.08	122.88	118.30

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	175	LEU	CA
1	A	177	THR	CA

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1338	0	1350	37	0
2	A	4	0	6	0	0
All	All	1342	0	1356	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:ARG:HG3	1:A:8:ARG:HH11	1.29	0.97
1:A:145:ARG:NH1	1:A:172:ILE:HA	1.93	0.83
1:A:119:ARG:HG2	1:A:123:GLN:NE2	1.98	0.78
1:A:143:PRO:O	1:A:147:LYS:HG3	1.90	0.71
1:A:145:ARG:HH11	1:A:172:ILE:HA	1.55	0.70
1:A:1:MET:HG2	1:A:158:TRP:CE3	2.31	0.65
1:A:22:GLU:HG3	1:A:174:HIS:HE1	1.61	0.64
1:A:70:ASP:HB3	1:A:175:LEU:HD12	1.80	0.62
1:A:119:ARG:HG2	1:A:123:GLN:HE21	1.65	0.61
1:A:8:ARG:HG3	1:A:8:ARG:NH1	2.03	0.58
1:A:131:VAL:O	1:A:135:LYS:HD2	2.04	0.57
1:A:1:MET:HG3	1:A:2:ASN:N	2.20	0.56
1:A:105:GLN:O	1:A:174:HIS:HB2	2.04	0.56
1:A:32:LEU:O	1:A:176:LEU:O	2.24	0.55
1:A:104:PHE:O	1:A:175:LEU:HD23	2.07	0.55
1:A:22:GLU:HG3	1:A:174:HIS:CE1	2.42	0.54
1:A:175:LEU:C	1:A:175:LEU:HD13	2.28	0.54
1:A:16:LYS:HD2	1:A:57:VAL:HG22	1.89	0.53
1:A:145:ARG:HD2	1:A:171:GLY:O	2.09	0.51
1:A:174:HIS:O	1:A:175:LEU:HB2	2.11	0.51
1:A:104:PHE:O	1:A:175:LEU:HB2	2.10	0.51
1:A:24:TYR:CZ	1:A:35:LYS:HD3	2.49	0.48
1:A:19:LYS:HD3	1:A:23:GLY:O	2.14	0.47
1:A:114:PHE:O	1:A:118:LEU:HG	2.14	0.47
1:A:21:THR:HG23	1:A:172:ILE:O	2.14	0.47
1:A:1:MET:HG2	1:A:158:TRP:CD2	2.51	0.45
1:A:8:ARG:NH1	1:A:8:ARG:CG	2.78	0.45
1:A:8:ARG:NH2	1:A:13:LEU:HB3	2.32	0.45
1:A:137:ARG:CZ	1:A:141:GLN:HG3	2.49	0.42
1:A:85:LYS:N	1:A:86:PRO:CD	2.82	0.42
1:A:1:MET:HE2	1:A:1:MET:HB2	1.94	0.42
1:A:8:ARG:HG3	1:A:8:ARG:O	2.20	0.41
1:A:27:ILE:HG12	1:A:28:GLY:H	1.86	0.41
1:A:36:SER:HA	1:A:37:PRO:HD3	1.93	0.40
1:A:85:LYS:N	1:A:86:PRO:HD2	2.36	0.40
1:A:116:ASN:HA	1:A:119:ARG:NH2	2.36	0.40
1:A:81:ASN:HB3	1:A:84:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	166/178 (93%)	158 (95%)	6 (4%)	2 (1%)	13 4

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	175	LEU
1	A	177	THR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	137/149 (92%)	131 (96%)	6 (4%)	28 19

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ARG
1	A	72	ASP
1	A	85	LYS
1	A	108	GLU
1	A	154	ARG
1	A	175	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN
1	A	123	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BME	A	200	-	3,3,3	16.71	1 (33%)	1,2,2	6.17	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BME	A	200	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	200	BME	O1-C1	28.93	2.91	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	200	BME	O1-C1-C2	-6.17	86.50	110.83

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	200	BME	O1-C1-C2-S2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	170/178 (95%)	0.82	12 (7%) 16 17	18, 29, 60, 79	8 (4%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	177	THR	17.9
1	A	175	LEU	16.0
1	A	171	GLY	13.4
1	A	176	LEU	13.2
1	A	174	HIS	12.5
1	A	173	GLY	12.3
1	A	172	ILE	10.5
1	A	178	LYS	10.4
1	A	36	SER	3.2
1	A	99	LEU	2.7
1	A	37	PRO	2.6
1	A	109	THR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BME	A	200	4/4	0.50	0.44	51,60,92,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.