



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 27, 2021 – 12:23 PM EDT

PDB ID : 7JTO
Title : Crystal structure of Protac MS33 in complex with the WD repeat-containing protein 5 and pVHL:ElonginC:ElonginB
Authors : Kottur, J.; Jain, R.; Aggarwal, A.K.
Deposited on : 2020-08-18
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	308	<div> <div>2%</div> <div>86% 12% .</div> </div>
2	J	104	<div> <div>90% 9% .</div> </div>
3	K	97	<div> <div>84% 6% 10%</div> </div>
4	L	162	<div> <div>2%</div> <div>77% 14% . 9%</div> </div>

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 5804 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WD repeat-containing protein 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	303	Total	C	N	O	S	0	2	0
			2332	1485	384	453	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	27	GLU	-	expression tag	UNP P61964
B	28	PHE	-	expression tag	UNP P61964
B	29	HIS	-	expression tag	UNP P61964
B	30	MET	-	expression tag	UNP P61964

- Molecule 2 is a protein called Elongin-B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	J	103	Total	C	N	O	S	0	0	0
			800	508	133	155	4			

- Molecule 3 is a protein called Elongin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	K	87	Total	C	N	O	S	0	0	0
			681	442	107	127	5			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	16	MET	-	initiating methionine	UNP Q15369


- Molecule 4 is a protein called von Hippel-Lindau disease tumor suppressor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	L	148	Total	C	N	O	S	0	6	0
			1242	793	225	222	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	52	GLY	-	expression tag	UNP P40337
L	53	SER	-	expression tag	UNP P40337

- Molecule 5 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	S	0	0
			3	1	1	1		
5	B	1	Total	C	N	S	0	0
			3	1	1	1		
5	B	1	Total	C	N	S	0	0
			3	1	1	1		
5	J	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	K	1	Total	C	O	0	0
			4	2	2		
6	L	1	Total	C	O	0	0
			4	2	2		
6	L	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is 3-methyl-N-(11-{[2-(4-{[4'-(4-methylpiperazin-1-yl)-3'-{[6-oxo-4-(trifluoromethyl)-5,6-dihydropyridine-3-carbonyl]amino}[1,1'-biphenyl]-3-yl]methyl}piperazin-1-yl)ethyl]amino}-11-oxoundecanoyl)-L-valyl-(4R)-4-hydroxy-N-{[4-(4-methyl-1,3-thiazol-5-yl)phenyl]methyl}-L-prolinamide (three-letter code: VKA) (formula: C₆₄H₈₄F₃N₁₁O₇S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	B	1	Total	C	F	N	O	S	0	0
			86	64	3	11	7	1		

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	K	1	6	3	3	0	0

- Molecule 9 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	L	1	Total 1	Na 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	B	273	Total 273	O 273	0	0
10	J	105	Total 105	O 105	0	0
10	K	82	Total 82	O 82	0	0
10	L	164	Total 164	O 164	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: WD repeat-containing protein 5




- Molecule 2: Elongin-B



- Molecule 3: Elongin-C



- Molecule 4: von Hippel-Lindau disease tumor suppressor



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.38Å 187.83Å 49.15Å 90.00° 116.96° 90.00°	Depositor
Resolution (Å)	46.96 – 1.70 93.92 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.96-1.70) 99.9 (93.92-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 1.70Å)	Xtriage
Refinement program	PHENIX 1.13	Depositor
R, R_{free}	0.200 , 0.225 0.198 , 0.224	Depositor DCC
R_{free} test set	4162 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.570	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5804	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VKA, SCN, NA, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.46	1/2388 (0.0%)	0.60	0/3247
2	J	0.51	0/816	0.60	0/1105
3	K	0.37	0/695	0.51	0/939
4	L	0.41	0/1277	0.59	0/1744
All	All	0.45	1/5176 (0.0%)	0.59	0/7035

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	228	TYR	C-O	-5.92	1.12	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2332	0	2257	28	0
2	J	800	0	787	7	0
3	K	681	0	669	4	0
4	L	1242	0	1211	26	0
5	B	9	0	0	0	0
5	J	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	8	0	12	3	0
6	K	4	0	6	0	0
6	L	8	0	12	3	0
7	B	86	0	0	0	0
8	K	6	0	8	3	0
9	L	1	0	0	0	0
10	B	273	0	0	6	1
10	J	105	0	0	2	0
10	K	82	0	0	4	0
10	L	164	0	0	11	1
All	All	5804	0	4962	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:211[A]:ASP:OD1	10:B:501:HOH:O	1.85	0.94
4:L:69[A]:ARG:HA	10:L:922:HOH:O	1.78	0.82
8:K:301:GOL:H11	10:K:404:HOH:O	1.80	0.81
4:L:69[B]:ARG:HA	10:L:922:HOH:O	1.81	0.80
8:K:301:GOL:C1	10:K:404:HOH:O	2.32	0.77
4:L:58:ARG:N	10:L:804:HOH:O	2.20	0.75
1:B:211[A]:ASP:OD2	1:B:239:LYS:NZ	2.21	0.71
4:L:69[A]:ARG:N	10:L:801:HOH:O	2.26	0.69
3:K:105:MET:HG2	6:L:701:EDO:H12	1.76	0.68
4:L:182:ARG:NE	10:L:807:HOH:O	2.27	0.68
1:B:172:ASP:OD2	10:B:503:HOH:O	2.15	0.65
4:L:184:LEU:HD12	6:L:701:EDO:H11	1.81	0.63
2:J:63:THR:HG23	2:J:66:THR:H	1.64	0.62
1:B:225:ASN:ND2	10:B:502:HOH:O	1.95	0.61
1:B:211[A]:ASP:OD1	1:B:211[A]:ASP:N	2.25	0.59
4:L:71:PRO:HB3	4:L:110[B]:HIS:CE1	2.38	0.59
4:L:109:ILE:HA	6:L:702:EDO:H12	1.85	0.59
1:B:33:PRO:HA	1:B:334:CYS:SG	2.42	0.59
4:L:69[B]:ARG:N	10:L:801:HOH:O	2.35	0.58
4:L:61:PRO:O	10:L:803:HOH:O	2.16	0.57
1:B:168:PRO:O	10:B:504:HOH:O	2.18	0.56
2:J:103:MET:O	10:J:301:HOH:O	2.17	0.56
1:B:38:LYS:CE	6:B:405:EDO:H11	2.37	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:64[A]:ARG:NE	4:L:91[A]:PHE:O	2.23	0.54
4:L:60:ARG:NH1	4:L:199:GLU:OE2	2.42	0.53
1:B:32:LYS:N	6:B:403:EDO:H21	2.24	0.52
4:L:66:VAL:HG23	4:L:66:VAL:O	2.08	0.52
1:B:38:LYS:HE3	6:B:405:EDO:H11	1.92	0.51
1:B:125:LEU:HB3	1:B:156:TRP:CE3	2.45	0.51
2:J:43:ARG:NH2	2:J:86:GLU:O	2.34	0.51
1:B:214:ASN:O	4:L:69[B]:ARG:NH1	2.43	0.51
1:B:265:ASN:ND2	1:B:307:THR:O	2.45	0.49
4:L:69[A]:ARG:NE	10:L:811:HOH:O	2.31	0.49
4:L:176:ARG:HA	4:L:185:TYR:CE1	2.47	0.48
1:B:47:ALA:HB2	1:B:322[B]:GLU:HG3	1.95	0.48
4:L:81:PRO:HD2	4:L:153:LEU:HG	1.93	0.48
8:K:301:GOL:H12	10:K:404:HOH:O	2.05	0.47
4:L:73:GLN:NE2	4:L:108:ARG:HE	2.13	0.46
1:B:168:PRO:HD2	10:B:699:HOH:O	2.16	0.45
2:J:23:THR:OG1	2:J:26:GLU:HG3	2.16	0.45
1:B:115:ASP:OD1	1:B:118:SER:OG	2.32	0.45
4:L:64[A]:ARG:HH21	4:L:91[A]:PHE:HB3	1.81	0.45
4:L:110[A]:HIS:CE1	10:L:816:HOH:O	2.69	0.45
1:B:63:SER:HB3	1:B:93:VAL:HG13	1.98	0.45
1:B:81:LYS:NZ	1:B:117:SER:O	2.49	0.45
3:K:101:LEU:O	3:K:105:MET:HG3	2.17	0.44
1:B:69:ILE:HD11	1:B:104:SER:HB3	2.00	0.44
2:J:100:PRO:HG2	2:J:103:MET:SD	2.58	0.44
1:B:315:ILE:HG12	1:B:331:LYS:HG2	2.00	0.44
1:B:218:SER:HB2	1:B:261:CYS:HA	1.98	0.44
4:L:73:GLN:OE1	4:L:110[A]:HIS:CE1	2.71	0.43
1:B:213:ASP:C	1:B:215:PRO:HD3	2.39	0.42
4:L:66:VAL:CG2	4:L:114:GLY:HA3	2.49	0.42
2:J:23:THR:HA	2:J:56:THR:HA	2.01	0.42
1:B:310:HIS:CG	1:B:311:PRO:HD2	2.54	0.42
2:J:86:GLU:OE1	10:J:302:HOH:O	2.21	0.42
1:B:43:GLY:O	1:B:70:LYS:NZ	2.50	0.41
3:K:86:SER:OG	3:K:88:THR:HG22	2.21	0.41
1:B:117:SER:OG	10:B:505:HOH:O	2.21	0.41
3:K:63:ARG:HD3	10:K:439:HOH:O	2.21	0.41
4:L:112:TYR:HB3	10:L:801:HOH:O	2.20	0.41
4:L:106:GLY:O	10:L:806:HOH:O	2.22	0.40
1:B:242:ASP:HB2	1:B:249:LEU:HD11	2.02	0.40
1:B:310:HIS:HB2	1:B:315:ILE:HB	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:66:VAL:O	4:L:66:VAL:CG2	2.70	0.40
4:L:173:GLU:CD	4:L:173:GLU:H	2.25	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:B:662:HOH:O	10:L:951:HOH:O[1_455]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	303/308 (98%)	288 (95%)	15 (5%)	0	100	100
2	J	101/104 (97%)	96 (95%)	5 (5%)	0	100	100
3	K	83/97 (86%)	83 (100%)	0	0	100	100
4	L	152/162 (94%)	150 (99%)	2 (1%)	0	100	100
All	All	639/671 (95%)	617 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	258/270 (96%)	256 (99%)	2 (1%)	81	74
2	J	87/92 (95%)	87 (100%)	0	100	100
3	K	74/86 (86%)	73 (99%)	1 (1%)	67	53
4	L	138/148 (93%)	137 (99%)	1 (1%)	84	77
All	All	557/596 (94%)	553 (99%)	4 (1%)	88	77

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	211[A]	ASP
1	B	211[B]	ASP
3	K	80	LYS
4	L	73	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SCN	B	401	-	1,2,2	0.85	0	0,1,1	-	-
5	SCN	B	404	-	1,2,2	0.75	0	0,1,1	-	-
6	EDO	L	702	-	3,3,3	0.26	0	2,2,2	0.45	0
5	SCN	J	201	-	1,2,2	0.89	0	0,1,1	-	-
5	SCN	B	402	-	1,2,2	0.73	0	0,1,1	-	-
6	EDO	B	405	-	3,3,3	0.22	0	2,2,2	0.88	0
7	VKA	B	406	-	86,93,93	3.70	34 (39%)	111,131,131	1.30	12 (10%)
6	EDO	B	403	-	3,3,3	0.51	0	2,2,2	0.05	0
8	GOL	K	301	-	5,5,5	0.96	0	5,5,5	0.86	0
6	EDO	K	302	-	3,3,3	0.49	0	2,2,2	0.34	0
6	EDO	L	701	-	3,3,3	0.60	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	L	702	-	-	1/1/1/1	-
6	EDO	B	405	-	-	1/1/1/1	-
7	VKA	B	406	-	-	15/75/120/120	0/8/8/8
6	EDO	B	403	-	-	0/1/1/1	-
8	GOL	K	301	-	-	2/4/4/4	-
6	EDO	K	302	-	-	1/1/1/1	-
6	EDO	L	701	-	-	0/1/1/1	-

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	406	VKA	C20-C16	-13.22	1.28	1.53
7	B	406	VKA	O86-C77	12.38	1.39	1.22
7	B	406	VKA	C18-N17	-10.24	1.31	1.47
7	B	406	VKA	C52-N49	-9.69	1.28	1.47
7	B	406	VKA	C30-N26	7.92	1.50	1.34
7	B	406	VKA	C44-N45	-7.72	1.29	1.47
7	B	406	VKA	C16-N17	7.37	1.62	1.47
7	B	406	VKA	C71-N68	-7.27	1.29	1.46
7	B	406	VKA	C40-N41	6.49	1.48	1.33
7	B	406	VKA	C21-N17	6.04	1.48	1.34
7	B	406	VKA	C6-C1	5.98	1.54	1.48
7	B	406	VKA	C69-N68	-4.98	1.35	1.46
7	B	406	VKA	C73-N72	4.74	1.45	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	406	VKA	C52-C53	4.56	1.59	1.51
7	B	406	VKA	C77-N76	-4.55	1.33	1.37
7	B	406	VKA	C50-N49	-4.01	1.35	1.46
7	B	406	VKA	C14-N13	3.92	1.42	1.33
7	B	406	VKA	C47-N45	-3.58	1.37	1.46
7	B	406	VKA	C48-N49	-3.53	1.37	1.46
7	B	406	VKA	C39-C40	3.46	1.57	1.51
7	B	406	VKA	C78-C79	-3.44	1.44	1.50
7	B	406	VKA	C51-N45	-3.38	1.37	1.46
7	B	406	VKA	C67-N68	-3.06	1.39	1.46
7	B	406	VKA	C62-N65	2.94	1.47	1.41
7	B	406	VKA	O42-C40	-2.76	1.17	1.23
7	B	406	VKA	O85-C73	-2.70	1.18	1.23
7	B	406	VKA	O23-C21	-2.55	1.17	1.22
7	B	406	VKA	C58-C53	2.38	1.44	1.38
7	B	406	VKA	C32-C31	2.31	1.60	1.52
7	B	406	VKA	C74-C73	2.30	1.57	1.49
7	B	406	VKA	C12-C11	2.24	1.56	1.51
7	B	406	VKA	C59-C55	2.11	1.54	1.49
7	B	406	VKA	O22-C19	-2.10	1.37	1.43
7	B	406	VKA	C24-N26	-2.10	1.42	1.45

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	406	VKA	C11-C12-N13	-4.25	103.95	113.05
7	B	406	VKA	C25-C24-C21	-3.73	108.61	113.40
7	B	406	VKA	C62-C61-N72	3.41	124.97	118.58
7	B	406	VKA	C25-C24-N26	-3.29	107.72	111.84
7	B	406	VKA	C60-C61-N72	-2.78	114.55	121.90
7	B	406	VKA	C60-C59-C55	-2.53	116.68	120.86
7	B	406	VKA	C74-C75-N76	-2.52	121.52	125.31
7	B	406	VKA	C64-C59-C60	2.19	121.26	118.16
7	B	406	VKA	C51-N45-C47	2.12	113.61	108.83
7	B	406	VKA	C39-C40-N41	2.12	119.99	116.42
7	B	406	VKA	C20-C16-N17	2.10	105.87	103.10
7	B	406	VKA	C69-N68-C67	2.09	112.45	109.52

There are no chirality outliers.

All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
8	K	301	GOL	O1-C1-C2-C3
7	B	406	VKA	C37-C38-C39-C40
8	K	301	GOL	O1-C1-C2-O2
7	B	406	VKA	N26-C30-C31-C32
7	B	406	VKA	C33-C34-C35-C36
7	B	406	VKA	O32-C30-C31-C32
7	B	406	VKA	C32-C33-C34-C35
7	B	406	VKA	C30-C31-C32-C33
6	B	405	EDO	O1-C1-C2-O2
6	K	302	EDO	O1-C1-C2-O2
6	L	702	EDO	O1-C1-C2-O2
7	B	406	VKA	C31-C32-C33-C34
7	B	406	VKA	N72-C73-C74-C75
7	B	406	VKA	O85-C73-C74-C75
7	B	406	VKA	N72-C73-C74-C79
7	B	406	VKA	O85-C73-C74-C79
7	B	406	VKA	C35-C36-C37-C38
7	B	406	VKA	C36-C37-C38-C39
7	B	406	VKA	C60-C61-N72-C73
7	B	406	VKA	C34-C35-C36-C37

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	L	702	EDO	1	0
6	B	405	EDO	2	0
6	B	403	EDO	1	0
8	K	301	GOL	3	0
6	L	701	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	303/308 (98%)	-0.27	4 (1%) 77 81	15, 25, 41, 59	0
2	J	103/104 (99%)	-0.33	0 100 100	18, 27, 41, 47	2 (1%)
3	K	87/97 (89%)	-0.28	0 100 100	16, 23, 45, 55	0
4	L	148/162 (91%)	-0.09	3 (2%) 65 69	16, 27, 43, 55	0
All	All	641/671 (95%)	-0.24	7 (1%) 80 83	15, 26, 42, 59	2 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	212	ASP	2.7
1	B	210	ILE	2.5
4	L	91[A]	PHE	2.4
1	B	228	TYR	2.2
4	L	142	VAL	2.1
1	B	211[A]	ASP	2.1
4	L	66	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	B	403	4/4	0.37	0.18	51,52,55,57	0
6	EDO	B	405	4/4	0.65	0.26	44,46,47,48	0
5	SCN	B	402	3/3	0.76	0.21	39,39,40,54	0
8	GOL	K	301	6/6	0.80	0.24	32,43,47,47	0
6	EDO	K	302	4/4	0.82	0.09	45,51,54,58	0
5	SCN	B	401	3/3	0.82	0.15	35,35,40,47	0
5	SCN	B	404	3/3	0.84	0.26	44,44,52,61	0
6	EDO	L	701	4/4	0.90	0.15	22,27,33,38	0
7	VKA	B	406	86/86	0.94	0.11	12,24,44,49	0
5	SCN	J	201	3/3	0.95	0.12	29,29,34,43	0
6	EDO	L	702	4/4	0.95	0.16	21,28,28,30	0
9	NA	L	703	1/1	0.97	0.05	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.