



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 2, 2020 – 06:08 PM EST

PDB ID : 7JTQ
Title : Human Complement Factor B Inhibited by a Slow Off-Rate Modified Aptamer of 31 Bases
Authors : Xu, X.; Geisbrecht, B.V.
Deposited on : 2020-08-18
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.14.6
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	764	<div> <div>%</div> <div> <div></div> <div>72%</div> <div>18%</div> <div>• 8%</div> </div> </div>
1	C	764	<div> <div>71%</div> <div>19%</div> <div>• 8%</div> </div>
2	B	32	<div> <div>6%</div> <div>50%</div> <div>41%</div> <div>•</div> </div>
2	D	32	<div> <div>6%</div> <div>63%</div> <div>28%</div> <div>•</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12610 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement factor B.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	703	Total	C	N	O	S	0	0	0
			5549	3494	963	1059	33			
1	C	703	Total	C	N	O	S	0	0	0
			5549	3494	963	1059	33			

- Molecule 2 is a DNA chain called DNA (32-MER).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	31	Total	C	N	O	P	0	0	0
			756	392	133	200	31			
2	D	31	Total	C	N	O	P	0	0	0
			756	392	133	200	31			



● Molecule 2: DNA (32-MER)



● Molecule 2: DNA (32-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.04Å 145.75Å 87.06Å 90.00° 109.62° 90.00°	Depositor
Resolution (Å)	42.88 – 3.50 42.88 – 3.50	Depositor EDS
% Data completeness (in resolution range)	96.7 (42.88-3.50) 96.6 (42.88-3.50)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 3.48Å)	Xtriage
Refinement program	PHENIX v1.17	Depositor
R, R_{free}	0.227 , 0.257 0.226 , 0.256	Depositor DCC
R_{free} test set	1998 reflections (8.01%)	wwPDB-VP
Wilson B-factor (Å ²)	82.4	Xtriage
Anisotropy	0.339	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.349 for l,-k,h	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	12610	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, OMG, 85Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	5/5673 (0.1%)	0.52	3/7678 (0.0%)
1	C	0.43	2/5673 (0.0%)	0.50	0/7678
2	B	1.54	1/358 (0.3%)	1.30	1/544 (0.2%)
2	D	0.92	0/358	0.86	0/544
All	All	0.53	8/12062 (0.1%)	0.57	4/16444 (0.0%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	481	LYS	C-N	10.68	1.52	1.33
1	C	479	PRO	N-CD	-9.37	1.34	1.47
1	A	479	PRO	N-CD	-9.36	1.34	1.47
1	A	348	PRO	N-CD	7.92	1.58	1.47
1	A	482	GLY	C-N	6.15	1.48	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	481	LYS	C-N-CA	-8.72	103.98	122.30
1	A	481	LYS	O-C-N	7.39	135.77	123.20
1	A	481	LYS	CA-C-N	-6.83	102.53	116.20
2	B	5	OMG	P-O3'-C3'	5.07	125.79	119.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5549	0	5419	103	0
1	C	5549	0	5419	118	0
2	B	756	0	283	38	0
2	D	756	0	283	33	0
All	All	12610	0	11404	259	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 259 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:478:ARG:NH1	1:A:507:ASP:OD2	1.75	1.19
1:C:478:ARG:NH1	1:C:507:ASP:OD2	1.75	1.18
1:C:477:ILE:HD13	1:C:483:HIS:ND1	1.68	1.08
1:C:347:VAL:HB	1:C:348:PRO:HD3	1.37	1.04
1:C:50:THR:HB	2:D:22:85Y:C27	1.87	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	693/764 (91%)	643 (93%)	47 (7%)	3 (0%)	34 72
1	C	693/764 (91%)	646 (93%)	45 (6%)	2 (0%)	41 75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1386/1528 (91%)	1289 (93%)	92 (7%)	5 (0%)	34 72

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	344	PRO
1	A	483	HIS
1	C	483	HIS
1	C	347	VAL
1	A	349	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	614/665 (92%)	592 (96%)	22 (4%)	35 66
1	C	614/665 (92%)	591 (96%)	23 (4%)	34 65
All	All	1228/1330 (92%)	1183 (96%)	45 (4%)	34 65

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	702	ASN
1	C	90	TRP
1	C	698	ASP
1	C	20	LYS
1	C	112	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	468	GLN
1	A	636	GLN
1	C	468	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	636	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

32 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OMG	D	18	2	18,26,27	3.88	6 (33%)	20,38,41	2.62	6 (30%)
2	OMG	B	18	2	18,26,27	4.34	6 (33%)	20,38,41	3.07	11 (55%)
2	OMG	B	2	2	18,26,27	3.99	6 (33%)	20,38,41	2.94	6 (30%)
2	85Y	D	10	2	30,36,37	3.34	9 (30%)	36,51,54	2.27	5 (13%)
2	OMG	D	5	2	18,26,27	3.80	7 (38%)	20,38,41	2.57	6 (30%)
2	OMC	D	30	2	15,22,23	4.09	6 (40%)	17,31,34	1.40	1 (5%)
2	85Y	B	28	2	30,36,37	3.72	11 (36%)	36,51,54	3.11	10 (27%)
2	OMG	D	2	2	18,26,27	3.71	7 (38%)	20,38,41	2.62	5 (25%)
2	85Y	B	4	2	30,36,37	3.65	10 (33%)	36,51,54	2.49	10 (27%)
2	85Y	B	24	2	30,36,37	3.44	11 (36%)	36,51,54	2.43	7 (19%)
2	OMG	B	19	2	18,26,27	2.96	7 (38%)	20,38,41	2.64	5 (25%)
2	85Y	B	10	2	30,36,37	3.26	9 (30%)	36,51,54	2.74	11 (30%)
2	85Y	D	22	2	30,36,37	3.43	8 (26%)	36,51,54	2.24	4 (11%)
2	OMG	B	5	2	18,26,27	3.79	6 (33%)	20,38,41	2.86	7 (35%)
2	OMC	B	30	2	15,22,23	4.30	6 (40%)	17,31,34	1.48	1 (5%)
2	85Y	B	16	2	30,36,37	3.31	12 (40%)	36,51,54	2.93	10 (27%)
2	85Y	D	16	2	30,36,37	3.48	11 (36%)	36,51,54	2.44	10 (27%)
2	OMG	D	31	2	18,26,27	3.89	7 (38%)	20,38,41	2.72	6 (30%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	OMC	D	3	2	15,22,23	3.84	6 (40%)	17,31,34	1.41	2 (11%)
2	OMG	B	29	2	18,26,27	3.75	7 (38%)	20,38,41	2.90	8 (40%)
2	85Y	D	24	2	30,36,37	3.50	11 (36%)	36,51,54	2.13	5 (13%)
2	OMG	D	19	2	18,26,27	3.29	7 (38%)	20,38,41	2.50	4 (20%)
2	OMC	D	1	2	15,22,23	4.05	6 (40%)	17,31,34	1.87	4 (23%)
2	85Y	D	27	2	30,36,37	3.55	9 (30%)	36,51,54	2.43	10 (27%)
2	85Y	D	28	2	30,36,37	3.59	10 (33%)	36,51,54	2.55	6 (16%)
2	OMG	B	31	2	18,26,27	4.25	6 (33%)	20,38,41	2.75	7 (35%)
2	OMC	B	1	2	15,22,23	4.22	6 (40%)	17,31,34	2.30	5 (29%)
2	OMC	B	3	2	15,22,23	3.60	6 (40%)	17,31,34	1.38	4 (23%)
2	OMG	D	29	2	18,26,27	3.66	7 (38%)	20,38,41	2.60	7 (35%)
2	85Y	D	4	2	30,36,37	3.57	10 (33%)	36,51,54	2.57	6 (16%)
2	85Y	B	22	2	30,36,37	3.51	9 (30%)	36,51,54	2.82	8 (22%)
2	85Y	B	27	2	30,36,37	3.72	11 (36%)	36,51,54	2.72	10 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OMG	D	18	2	-	4/5/27/28	0/3/3/3
2	OMG	B	18	2	-	4/5/27/28	0/3/3/3
2	OMG	B	2	2	-	2/5/27/28	0/3/3/3
2	85Y	D	10	2	-	6/13/30/31	0/4/4/4
2	OMG	D	5	2	-	2/5/27/28	0/3/3/3
2	OMC	D	30	2	-	4/7/27/28	0/2/2/2
2	85Y	B	28	2	-	7/13/30/31	0/4/4/4
2	OMG	D	2	2	-	3/5/27/28	0/3/3/3
2	85Y	B	4	2	-	2/13/30/31	0/4/4/4
2	85Y	B	24	2	-	2/13/30/31	0/4/4/4
2	OMG	B	19	2	-	4/5/27/28	0/3/3/3
2	85Y	B	10	2	-	6/13/30/31	0/4/4/4
2	85Y	D	22	2	-	6/13/30/31	0/4/4/4
2	OMG	B	5	2	-	3/5/27/28	0/3/3/3
2	OMC	B	30	2	-	4/7/27/28	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	85Y	B	16	2	-	4/13/30/31	0/4/4/4
2	85Y	D	16	2	-	5/13/30/31	0/4/4/4
2	OMG	D	31	2	-	3/5/27/28	0/3/3/3
2	OMC	D	3	2	-	0/7/27/28	0/2/2/2
2	OMG	B	29	2	-	2/5/27/28	0/3/3/3
2	85Y	D	24	2	-	3/13/30/31	0/4/4/4
2	OMG	D	19	2	-	3/5/27/28	0/3/3/3
2	OMC	D	1	2	-	4/7/27/28	0/2/2/2
2	85Y	D	27	2	-	2/13/30/31	0/4/4/4
2	85Y	D	28	2	-	6/13/30/31	0/4/4/4
2	OMG	B	31	2	-	3/5/27/28	0/3/3/3
2	OMC	B	1	2	-	4/7/27/28	0/2/2/2
2	OMC	B	3	2	-	2/7/27/28	0/2/2/2
2	OMG	D	29	2	-	3/5/27/28	0/3/3/3
2	85Y	D	4	2	-	2/13/30/31	0/4/4/4
2	85Y	B	22	2	-	5/13/30/31	0/4/4/4
2	85Y	B	27	2	-	2/13/30/31	0/4/4/4

The worst 5 of 256 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	30	OMC	C6-N1	10.98	1.49	1.35
2	B	31	OMG	C4-N3	10.39	1.51	1.35
2	D	30	OMC	C6-N1	10.30	1.48	1.35
2	B	2	OMG	C4-N3	10.16	1.51	1.35
2	B	1	OMC	C6-N1	10.02	1.48	1.35

The worst 5 of 207 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	28	85Y	C23-C24-C25	10.33	140.01	121.51
2	B	10	85Y	C23-C24-C29	-9.43	101.34	120.91
2	D	4	85Y	C24-C23-N22	-9.27	93.20	113.05
2	D	28	85Y	C23-C24-C25	9.02	137.66	121.51
2	B	2	OMG	C1'-N9-C4	9.02	142.49	126.64

There are no chirality outliers.

5 of 112 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	18	OMG	C1'-C2'-O2'-CM2
2	B	18	OMG	C1'-C2'-O2'-CM2
2	B	2	OMG	C1'-C2'-O2'-CM2
2	D	10	85Y	C3'-C4'-C5'-O5'
2	D	30	OMC	C1'-C2'-O2'-CM2

There are no ring outliers.

21 monomers are involved in 40 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	18	OMG	4	0
2	B	18	OMG	3	0
2	B	2	OMG	2	0
2	D	10	85Y	1	0
2	D	5	OMG	1	0
2	D	2	OMG	1	0
2	B	4	85Y	1	0
2	B	19	OMG	5	0
2	B	10	85Y	3	0
2	D	22	85Y	3	0
2	B	5	OMG	1	0
2	B	30	OMC	1	0
2	B	16	85Y	2	0
2	D	16	85Y	2	0
2	D	3	OMC	2	0
2	B	29	OMG	1	0
2	D	19	OMG	4	0
2	B	1	OMC	1	0
2	B	3	OMC	3	0
2	D	29	OMG	2	0
2	B	27	85Y	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	703/764 (92%)	-0.05	4 (0%) 89 86	43, 78, 122, 157	0
1	C	703/764 (92%)	-0.06	3 (0%) 92 90	45, 80, 124, 162	0
2	B	15/32 (46%)	-0.45	0 100 100	61, 69, 81, 85	0
2	D	15/32 (46%)	-0.39	0 100 100	67, 73, 80, 86	0
All	All	1436/1592 (90%)	-0.06	7 (0%) 91 88	43, 79, 122, 162	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	707	LYS	2.9
1	C	707	LYS	2.7
1	A	217	ILE	2.5
1	A	137	ALA	2.2
1	C	346	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OMG	D	31	24/25	0.79	0.22	104,139,152,157	0
2	OMC	D	1	21/22	0.81	0.25	118,138,161,186	0
2	OMG	B	31	24/25	0.83	0.18	116,130,143,147	0
2	OMC	B	1	21/22	0.85	0.29	125,149,165,173	0
2	OMG	D	18	24/25	0.86	0.27	83,102,118,119	0
2	85Y	D	28	33/34	0.87	0.32	62,77,91,112	0
2	OMG	D	29	24/25	0.87	0.24	101,109,129,138	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	OMG	B	29	24/25	0.88	0.20	89,110,129,138	0
2	85Y	B	28	33/34	0.88	0.27	54,70,97,119	0
2	OMC	D	30	21/22	0.89	0.17	102,118,130,143	0
2	85Y	D	27	33/34	0.89	0.24	64,74,102,105	0
2	OMG	B	2	24/25	0.89	0.17	100,122,134,157	0
2	OMG	B	18	24/25	0.90	0.26	74,89,112,119	0
2	OMG	D	2	24/25	0.90	0.18	102,121,143,187	0
2	OMC	B	30	21/22	0.91	0.12	95,112,138,159	0
2	85Y	D	16	33/34	0.91	0.24	59,75,143,154	0
2	85Y	B	16	33/34	0.92	0.18	55,77,108,111	0
2	OMG	D	5	24/25	0.92	0.22	59,74,91,102	0
2	OMG	D	19	24/25	0.92	0.22	62,86,98,101	0
2	85Y	D	24	33/34	0.93	0.28	61,71,77,86	0
2	85Y	D	22	33/34	0.93	0.32	56,74,84,94	0
2	85Y	D	4	33/34	0.93	0.27	61,84,100,114	0
2	85Y	B	27	33/34	0.93	0.24	57,71,90,92	0
2	OMC	D	3	21/22	0.94	0.20	89,99,115,140	0
2	OMC	B	3	21/22	0.94	0.22	82,92,109,120	0
2	85Y	B	4	33/34	0.94	0.24	62,76,101,108	0
2	OMG	B	5	24/25	0.94	0.21	66,76,89,96	0
2	OMG	B	19	24/25	0.94	0.18	61,82,93,95	0
2	85Y	D	10	33/34	0.95	0.29	50,71,110,115	0
2	85Y	B	10	33/34	0.95	0.26	55,68,108,119	0
2	85Y	B	22	33/34	0.95	0.28	52,71,89,96	0
2	85Y	B	24	33/34	0.95	0.21	63,70,82,89	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.