



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 03:33 AM BST

PDB ID : 3JUV  
Title : Crystal structure of human lanosterol 14alpha-demethylase (CYP51)  
Authors : Strushkevich, N.; MacKenzie, F.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Weigelt, J.; Usanov, S.A.; Park, H.; Structural Genomics Consortium (SGC)  
Deposited on : 2009-09-15  
Resolution : 3.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1292 (3.14-3.10)
Clashscore	141614	1389 (3.14-3.10)
Ramachandran outliers	138981	1337 (3.14-3.10)
Sidechain outliers	138945	1337 (3.14-3.10)
RSRZ outliers	127900	1260 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>21%</div> <div>5%</div> </div> </div>
2	B	7	<div> <div>29%</div> <div>71%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	B	7	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3736 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lanosterol 14-alpha demethylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	448	3605	2320	614	655	16	0	0	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	49	MET	-	expression tag	UNP Q16850
A	50	ALA	-	expression tag	UNP Q16850
A	51	LYS	-	expression tag	UNP Q16850
A	52	LYS	-	expression tag	UNP Q16850
A	53	THR	-	expression tag	UNP Q16850
A	503	THR	-	expression tag	UNP Q16850
A	504	HIS	-	expression tag	UNP Q16850
A	505	HIS	-	expression tag	UNP Q16850
A	506	HIS	-	expression tag	UNP Q16850
A	507	HIS	-	expression tag	UNP Q16850
A	508	HIS	-	expression tag	UNP Q16850
A	509	HIS	-	expression tag	UNP Q16850

- Molecule 2 is an oligosaccharide called Cycloheptakis-(1-4)-(alpha-D-glucopyranose).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	7	77	42	35	0	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total	O	0	0
			11	11		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lanosterol 14-alpha demethylase



- Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	174.14Å 174.14Å 244.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.67 – 3.12 24.67 – 3.12	Depositor EDS
% Data completeness (in resolution range)	99.5 (24.67-3.12) 99.5 (24.67-3.12)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.05 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, $R_{free}$	0.198 , 0.227 0.198 , 0.227	Depositor DCC
$R_{free}$ test set	1300 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	80.7	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3736	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/3699	0.61	1/5011 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	252	ARG	NE-CZ-NH2	-5.37	117.62	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3605	0	3597	82	0
2	B	77	0	63	6	0
3	A	43	0	30	1	0
4	A	11	0	0	0	0
All	All	3736	0	3690	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:GLN:HE22	1:A:472:PHE:H	1.10	0.91
1:A:116:LEU:HD22	1:A:390:VAL:HG22	1.51	0.90
1:A:300:GLU:O	1:A:304:MET:HG2	1.76	0.85
1:A:353:THR:H	1:A:356:GLN:NE2	1.72	0.85
1:A:290:TYR:HB2	1:A:292:ASP:HB2	1.62	0.81
1:A:292:ASP:HB3	1:A:294:ARG:H	1.45	0.79
1:A:139:PHE:CD2	1:A:307:GLY:HA3	2.17	0.79
1:A:418:ARG:HG3	1:A:419:LEU:HG	1.67	0.77
1:A:289:THR:HG22	1:A:295:PRO:HA	1.69	0.73
1:A:369:GLU:OE1	1:A:372:ARG:NH1	2.22	0.72
2:B:5:GLC:H61	2:B:6:GLC:O6	1.89	0.72
2:B:3:GLC:HO3	2:B:4:GLC:HO2	1.29	0.70
1:A:471:GLU:OE1	1:A:499:LYS:HE2	1.94	0.68
1:A:328:ALA:HB1	1:A:473:ASP:HA	1.75	0.68
1:A:200:GLU:O	1:A:203:ILE:HG22	1.94	0.67
1:A:144:ALA:HA	1:A:152:PHE:CD2	2.30	0.67
1:A:203:ILE:O	1:A:207:SER:HB2	1.95	0.67
1:A:213:LYS:NZ	1:A:278:ILE:HD12	2.11	0.65
1:A:456:TYR:O	1:A:460:LYS:HB2	1.97	0.64
2:B:5:GLC:C6	2:B:6:GLC:O6	2.46	0.63
1:A:116:LEU:HD22	1:A:390:VAL:CG2	2.28	0.62
1:A:372:ARG:HG2	1:A:409:GLN:HB3	1.82	0.62
1:A:372:ARG:HD2	1:A:415:TRP:CZ3	2.35	0.62
1:A:68:ILE:HD12	1:A:68:ILE:N	2.16	0.61
1:A:484:TYR:HD2	1:A:489:HIS:CE1	2.18	0.61
1:A:352:LEU:HA	1:A:356:GLN:HE22	1.67	0.60
1:A:226:LEU:HD21	1:A:259:GLU:HG2	1.81	0.60
1:A:381:MET:HG3	1:A:402:CYS:SG	2.42	0.59
1:A:426:TYR:HA	1:A:430:ASN:OD1	2.02	0.59
1:A:242:PRO:HD2	1:A:245:LEU:HD11	1.83	0.58
1:A:338:TYR:O	1:A:341:GLN:HB2	2.02	0.58
1:A:334:GLN:HE21	1:A:501:ARG:HH12	1.53	0.57
1:A:208:HIS:CA	1:A:216:ARG:HG3	2.35	0.56
1:A:164:ASN:ND2	1:A:166:ALA:HB3	2.20	0.56
1:A:159:LEU:HD11	1:A:304:MET:HB2	1.87	0.56
1:A:422:ASN:HD21	1:A:424:ASP:HB2	1.73	0.54
1:A:204:LEU:HD23	1:A:219:LEU:HD23	1.90	0.54
1:A:213:LYS:HZ3	1:A:278:ILE:HD12	1.71	0.53
1:A:483:ASN:HD21	1:A:485:THR:HB	1.74	0.53
1:A:471:GLU:HG3	1:A:501:ARG:NH2	2.23	0.53
1:A:372:ARG:HA	1:A:409:GLN:NE2	2.24	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:LYS:HG3	1:A:446:ARG:HH22	1.74	0.52
1:A:484:TYR:CD2	1:A:489:HIS:CE1	2.98	0.52
1:A:422:ASN:O	1:A:425:ARG:HG2	2.11	0.51
1:A:430:ASN:ND2	1:A:432:ALA:HB3	2.25	0.51
1:A:369:GLU:CD	1:A:372:ARG:NH1	2.65	0.50
1:A:337:CYS:HB3	1:A:362:LEU:HD23	1.94	0.49
1:A:289:THR:HG22	1:A:295:PRO:CA	2.42	0.48
1:A:353:THR:H	1:A:356:GLN:HE22	1.58	0.48
1:A:315:THR:HB	3:A:601:HEM:HBB2	1.95	0.48
1:A:160:LYS:HG2	1:A:450:ILE:HB	1.96	0.48
1:A:369:GLU:CD	1:A:372:ARG:HH12	2.16	0.48
1:A:463:TRP:O	1:A:467:LEU:HB2	2.12	0.48
1:A:334:GLN:NE2	1:A:472:PHE:H	1.94	0.47
1:A:483:ASN:ND2	1:A:485:THR:H	2.13	0.47
1:A:191:GLU:CG	1:A:497:ARG:HG3	2.44	0.47
2:B:2:GLC:H61	2:B:3:GLC:C1	2.46	0.46
1:A:165:ILE:O	1:A:169:LYS:HG3	2.16	0.45
1:A:194:VAL:HG13	1:A:496:ILE:HG21	1.99	0.45
1:A:390:VAL:O	1:A:392:GLY:N	2.43	0.45
2:B:3:GLC:O3	2:B:4:GLC:O2	2.07	0.45
1:A:213:LYS:HZ1	1:A:278:ILE:HD12	1.80	0.45
1:A:369:GLU:HG3	1:A:421:PHE:CE1	2.52	0.45
1:A:121:LYS:HG3	1:A:446:ARG:NH2	2.32	0.45
1:A:230:LEU:HD12	1:A:230:LEU:HA	1.71	0.45
1:A:164:ASN:ND2	1:A:167:HIS:H	2.15	0.45
1:A:165:ILE:HG13	1:A:165:ILE:H	1.58	0.44
1:A:372:ARG:HG3	1:A:439:TYR:CD1	2.52	0.44
1:A:214:GLU:O	1:A:217:SER:OG	2.35	0.44
1:A:369:GLU:HG3	1:A:421:PHE:CD1	2.52	0.44
2:B:4:GLC:O6	2:B:5:GLC:C1	2.67	0.43
1:A:149:ASN:N	1:A:150:PRO:HD2	2.34	0.42
1:A:356:GLN:HB2	1:A:356:GLN:HE21	1.66	0.42
1:A:101:VAL:HG11	1:A:242:PRO:HD3	2.01	0.42
1:A:139:PHE:HB3	1:A:143:VAL:HG22	2.01	0.42
1:A:124:ASP:HB3	1:A:388:GLN:NE2	2.34	0.42
1:A:120:SER:HB3	1:A:125:LEU:HD12	2.01	0.42
1:A:381:MET:O	1:A:382:ARG:HD3	2.20	0.42
1:A:417:GLU:O	1:A:420:ASP:HB2	2.19	0.42
1:A:336:LYS:HE3	1:A:336:LYS:HB3	1.85	0.41
1:A:353:THR:N	1:A:356:GLN:NE2	2.53	0.41
1:A:66:SER:HB2	1:A:73:HIS:CE1	2.56	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:ASN:HD21	1:A:166:ALA:HB3	1.83	0.41
1:A:372:ARG:HD2	1:A:415:TRP:CH2	2.56	0.41
1:A:446:ARG:H	1:A:446:ARG:HG2	1.59	0.41
1:A:208:HIS:N	1:A:216:ARG:HG3	2.36	0.40
1:A:390:VAL:HB	1:A:391:ALA:H	1.75	0.40
1:A:194:VAL:HG13	1:A:496:ILE:CG2	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/461 (97%)	409 (92%)	32 (7%)	5 (1%)	14	45

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	144	ALA
1	A	273	GLN
1	A	347	GLU
1	A	428	GLN
1	A	390	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	391/403 (97%)	353 (90%)	38 (10%)	8 30

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	LYS
1	A	60	SER
1	A	123	GLU
1	A	143	VAL
1	A	160	LYS
1	A	164	ASN
1	A	180	LYS
1	A	194	VAL
1	A	207	SER
1	A	213	LYS
1	A	216	ARG
1	A	221	GLU
1	A	225	GLN
1	A	245	LEU
1	A	249	SER
1	A	271	ARG
1	A	274	SER
1	A	292	ASP
1	A	297	THR
1	A	313	GLN
1	A	317	SER
1	A	333	LEU
1	A	345	CYS
1	A	352	LEU
1	A	353	THR
1	A	381	MET
1	A	389	THR
1	A	390	VAL
1	A	418	ARG
1	A	420	ASP
1	A	422	ASN
1	A	430	ASN
1	A	446	ARG
1	A	473	ASP
1	A	483	ASN
1	A	485	THR
1	A	487	MET
1	A	501	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	164	ASN
1	A	334	GLN
1	A	356	GLN
1	A	400	GLN
1	A	422	ASN
1	A	430	ASN
1	A	483	ASN
1	A	489	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	B	1	2	11,11,12	0.61	0	15,15,17	1.58	3 (20%)
2	GLC	B	2	2	11,11,12	0.47	0	15,15,17	1.82	5 (33%)
2	GLC	B	3	2	11,11,12	0.63	0	15,15,17	1.10	1 (6%)
2	GLC	B	4	2	11,11,12	0.69	0	15,15,17	1.91	5 (33%)
2	GLC	B	5	2	11,11,12	0.71	0	15,15,17	1.71	3 (20%)
2	GLC	B	6	2	11,11,12	0.75	0	15,15,17	1.94	3 (20%)
2	GLC	B	7	2	11,11,12	0.60	0	15,15,17	2.67	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/19/22	0/1/1/1
2	GLC	B	2	2	-	2/2/19/22	0/1/1/1
2	GLC	B	3	2	-	2/2/19/22	0/1/1/1
2	GLC	B	4	2	-	1/2/19/22	0/1/1/1
2	GLC	B	5	2	-	2/2/19/22	0/1/1/1
2	GLC	B	6	2	-	0/2/19/22	0/1/1/1
2	GLC	B	7	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	7	GLC	C1-C2-C3	6.29	117.39	109.67
2	B	7	GLC	C1-O5-C5	5.15	119.16	112.19
2	B	6	GLC	C1-C2-C3	5.06	115.88	109.67
2	B	5	GLC	C1-C2-C3	4.39	115.06	109.67
2	B	7	GLC	C2-C3-C4	4.19	118.14	110.89
2	B	2	GLC	O5-C5-C6	3.80	113.16	107.20
2	B	4	GLC	C3-C4-C5	3.78	116.97	110.24
2	B	6	GLC	C2-C3-C4	3.67	117.24	110.89
2	B	4	GLC	C1-C2-C3	3.20	113.60	109.67
2	B	1	GLC	O5-C1-C2	-3.14	105.92	110.77
2	B	2	GLC	C1-O5-C5	3.13	116.44	112.19
2	B	4	GLC	C2-C3-C4	2.89	115.89	110.89
2	B	1	GLC	C1-C2-C3	2.84	113.16	109.67
2	B	6	GLC	C1-O5-C5	-2.83	108.36	112.19
2	B	7	GLC	C3-C4-C5	2.80	115.23	110.24
2	B	5	GLC	C1-O5-C5	2.80	115.98	112.19
2	B	2	GLC	C3-C4-C5	2.73	115.11	110.24
2	B	5	GLC	O5-C5-C6	2.69	111.42	107.20
2	B	3	GLC	C1-O5-C5	2.60	115.72	112.19
2	B	2	GLC	C6-C5-C4	-2.39	107.40	113.00
2	B	4	GLC	O5-C5-C4	2.37	116.59	110.83
2	B	4	GLC	O5-C1-C2	-2.37	107.12	110.77
2	B	1	GLC	C2-C3-C4	2.33	114.94	110.89
2	B	2	GLC	O3-C3-C4	-2.24	105.16	110.35

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2	GLC	O5-C5-C6-O6
2	B	3	GLC	O5-C5-C6-O6
2	B	2	GLC	C4-C5-C6-O6
2	B	3	GLC	C4-C5-C6-O6
2	B	7	GLC	O5-C5-C6-O6
2	B	5	GLC	C4-C5-C6-O6
2	B	5	GLC	O5-C5-C6-O6
2	B	4	GLC	O5-C5-C6-O6
2	B	7	GLC	C4-C5-C6-O6

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	5	GLC	3	0
2	B	4	GLC	3	0
2	B	2	GLC	1	0
2	B	3	GLC	3	0
2	B	6	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection.



RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HEM	A	601	1	27,50,50	2.12	5 (18%)	17,82,82	1.95	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	601	1	-	0/6/54/54	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	HEM	C3B-C2B	-4.87	1.33	1.40
3	A	601	HEM	C3D-C2D	4.77	1.51	1.37
3	A	601	HEM	C3C-C2C	-4.38	1.34	1.40
3	A	601	HEM	C3C-CAC	3.64	1.55	1.47
3	A	601	HEM	C3B-CAB	3.30	1.54	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	HEM	CBD-CAD-C3D	-5.36	102.60	112.48
3	A	601	HEM	C1D-C2D-C3D	-4.49	103.87	107.00
3	A	601	HEM	CAA-CBA-CGA	-2.76	108.04	112.67

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	HEM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	448/461 (97%)	-0.25	8 (1%) 68 48	53, 75, 97, 119	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	55	PRO	3.7
1	A	277	LYS	3.3
1	A	502	SER	2.9
1	A	275	GLN	2.3
1	A	273	GLN	2.2
1	A	56	ALA	2.1
1	A	293	GLY	2.1
1	A	274	SER	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	B	7	11/12	0.67	0.43	66,68,69,70	11
2	GLC	B	2	11/12	0.76	0.30	64,65,67,68	11
2	GLC	B	1	11/12	0.78	0.34	65,66,67,67	11
2	GLC	B	3	11/12	0.83	0.23	69,70,71,71	11

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	B	6	11/12	0.87	0.21	70,72,74,75	11
2	GLC	B	5	11/12	0.93	0.14	70,72,72,72	11
2	GLC	B	4	11/12	0.94	0.14	69,71,71,72	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	HEM	A	601	43/43	0.99	0.15	53,59,67,71	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers ⓘ

There are no such residues in this entry.