



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 02:26 pm BST

PDB ID : 3JV2
Title : Crystal Structure of B. subtilis SecA with bound peptide
Authors : Zimmer, J.
Deposited on : 2009-09-15
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	783	<div> <div>2%</div> <div> <div></div> <div>66%</div> <div>25%</div> <div>5%</div> <div>• •</div> </div> </div>
1	B	783	<div> <div>%</div> <div> <div></div> <div>69%</div> <div>21%</div> <div>6%</div> <div>• •</div> </div> </div>
2	C	3	<div> <div></div> <div> <div>67%</div> <div>33%</div> </div> </div>
2	D	3	<div> <div></div> <div> <div>67%</div> <div>33%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12364 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein translocase subunit secA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	758	Total	C	N	O	S	0	0	0
			6043	3782	1053	1175	33			
1	B	762	Total	C	N	O	S	0	0	0
			6083	3804	1063	1183	33			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P28366
A	-1	PRO	-	EXPRESSION TAG	UNP P28366
A	0	HIS	-	EXPRESSION TAG	UNP P28366
B	-2	GLY	-	EXPRESSION TAG	UNP P28366
B	-1	PRO	-	EXPRESSION TAG	UNP P28366
B	0	HIS	-	EXPRESSION TAG	UNP P28366

- Molecule 2 is a protein called peptide.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			15	9	3	3			
2	D	3	Total	C	N	O	0	0	0
			15	9	3	3			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:

C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	81	Total	O	0	0
			81	81		
5	B	71	Total	O	0	0
			71	71		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:

Position	Amino Acid	Frequency (bits)
1	T694	0.01
2	K695	0.01
3	Y696	0.01
4	N697	0.01
5	E698	0.01
6	K699	0.01
7	Q702	0.01
8	F703	0.01
9	G704	0.01
10	K705	0.01
11	E706	0.01
12	Q707	0.01
13	E712	0.01
14	K713	0.01
15	N725	0.01
16	L740	0.01
17	R741	0.01
18	A742	0.01
19	Y743	0.01
20	A744	0.01
21	Q745	0.01
22	T746	0.01
23	N747	0.01
24	P748	0.01
25	E751	0.01
26	M754	0.01
27	E755	0.01
28	E761	0.01
29	E765	0.01
30	T766	0.01
31	I767	0.01
32	E768	0.01
33	D769	0.01
34	E770	0.01
35	V771	0.01
36	F774	0.01
37	V775	0.01
38	K776	0.01
39	K777	0.01
40	A778	0.01
41	E779	0.01
42	I780	0.01
43	R466	0.01
44	E467	0.01
45	M496	0.01
46	R489	0.01
47	I493	0.01
48	V507	0.01
49	E511	0.01
50	R512	0.01
51	H513	0.01
52	R516	0.01
53	R517	0.01
54	Q521	0.01
55	R528	0.01
56	T535	0.01
57	L539	0.01
58	M541	0.01
59	E544	0.01
60	L545	0.01
61	M546	0.01
62	R547	0.01
63	L560	0.01
64	I561	0.01
65	M562	0.01
66	T563	0.01
67	L666	0.01
68	D667	0.01
69	E668	0.01
70	L671	0.01
71	E672	0.01
72	S674	0.01
73	D675	0.01
74	I676	0.01
75	G677	0.01
76	G678	0.01
77	K679	0.01
78	D682	0.01
79	E683	0.01
80	L684	0.01
81	L685	0.01
82	E686	0.01
83	L687	0.01
84	D690	0.01
85	R466	0.01
86	E467	0.01
87	M496	0.01
88	R489	0.01
89	I493	0.01
90	V507	0.01
91	E511	0.01
92	R512	0.01
93	H513	0.01
94	R516	0.01
95	R517	0.01
96	Q521	0.01
97	R528	0.01
98	T535	0.01
99	L539	0.01
100	M541	0.01
101	E544	0.01
102	L545	0.01
103	M546	0.01
104	R547	0.01
105	L560	0.01
106	I561	0.01
107	M562	0.01
108	T563	0.01
109	L666	0.01
110	D667	0.01
111	E668	0.01
112	L671	0.01
113	E672	0.01
114	S674	0.01
115	D675	0.01
116	I676	0.01
117	G677	0.01
118	G678	0.01
119	K679	0.01
120	D682	0.01
121	E683	0.01
122	L684	0.01
123	L685	0.01
124	E686	0.01
125	L687	0.01
126	D690	0.01
127	R466	0.01
128	E467	0.01
129	M496	0.01
130	R489	0.01
131	I493	0.01
132	V507	0.01
133	E511	

- Chain B:
-
- 69% 21% 6%
- GLY PRO HIS MET LEU GLY ILE LEU ASN LYS MET PHE ASP PRO THR LYS R14 M17 E20 G31 D32 Y33 E34 N35 L36 A40 L41 K42 L62 V69 R70 P81 F82 K83 V84 Q85 L86 H93 N96 K101 K106 L112 P113 N116 M117 Q123



● Molecule 2: peptide



● Molecule 2: peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	106.72Å 106.72Å 175.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50 32.31 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.50) 99.7 (32.31-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.208 , 0.258 0.210 , 0.259	Depositor DCC
R_{free} test set	3443 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtriage
Anisotropy	0.196	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 12.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.488 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12364	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/6127	0.60	5/8239 (0.1%)
1	B	0.50	0/6167	1.10	21/8291 (0.3%)
All	All	0.49	0/12294	0.89	26/16530 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	5
All	All	0	6

There are no bond length outliers.

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	341	GLU	CA-C-N	29.99	176.18	116.20
1	B	325	LYS	CA-C-N	27.76	171.72	116.20
1	B	318	SER	CA-C-N	26.39	175.26	117.20
1	B	255	LYS	CA-C-N	24.04	170.09	117.20
1	B	319	PHE	O-C-N	-23.70	84.78	122.70

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	742	ALA	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	255	LYS	Mainchain
1	B	318	SER	Mainchain
1	B	325	LYS	Mainchain
1	B	341	GLU	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6043	0	6031	292	0
1	B	6083	0	6069	281	0
2	C	15	0	5	1	0
2	D	15	0	6	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	27	0	12	0	0
4	B	27	0	12	1	0
5	A	81	0	0	8	0
5	B	71	0	0	6	0
All	All	12364	0	12135	573	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 573 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:316:VAL:HG12	1:A:320:THR:CG2	1.37	1.50
1:A:316:VAL:CG1	1:A:320:THR:HG21	1.41	1.49
1:A:673:LYS:HA	1:A:676:ILE:CD1	1.41	1.45
1:A:677:PHE:CE1	1:A:679:LYS:HD3	1.60	1.36
1:A:317:ASP:HB3	1:A:322:ARG:CG	1.55	1.35

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	754/783 (96%)	724 (96%)	28 (4%)	2 (0%)	41	61
1	B	758/783 (97%)	732 (97%)	24 (3%)	2 (0%)	41	61
All	All	1512/1566 (97%)	1456 (96%)	52 (3%)	4 (0%)	41	61

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	550	GLY
1	A	677	PHE
1	B	317	ASP
1	A	318	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	649/672 (97%)	570 (88%)	79 (12%)	5	9
1	B	653/672 (97%)	571 (87%)	82 (13%)	4	8
All	All	1302/1344 (97%)	1141 (88%)	161 (12%)	4	9

5 of 161 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	703	PHE
1	B	248	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	646	GLU
1	A	713	LYS
1	B	69	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	521	GLN
1	B	85	GLN
1	B	521	GLN
1	A	582	GLN
1	A	588	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ADP	A	873	3	24,29,29	0.97	1 (4%)	29,45,45	1.31	4 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ADP	B	873	3	24,29,29	0.99	1 (4%)	29,45,45	1.41	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	873	3	-	6/12/32/32	0/3/3/3
4	ADP	B	873	3	-	5/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	873	ADP	C5-C4	2.58	1.47	1.40
4	A	873	ADP	C5-C4	2.50	1.47	1.40

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	873	ADP	N3-C2-N1	-3.36	123.43	128.68
4	B	873	ADP	N3-C2-N1	-3.22	123.64	128.68
4	B	873	ADP	C4-C5-N7	-3.08	106.19	109.40
4	B	873	ADP	PA-O3A-PB	-2.83	123.13	132.83
4	A	873	ADP	PA-O3A-PB	-2.72	123.48	132.83

There are no chirality outliers.

5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	873	ADP	C5'-O5'-PA-O1A
4	A	873	ADP	C5'-O5'-PA-O2A
4	A	873	ADP	C5'-O5'-PA-O3A
4	A	873	ADP	C3'-C4'-C5'-O5'
4	A	873	ADP	O4'-C4'-C5'-O5'


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	873	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	758/783 (96%)	-0.30	15 (1%) 65 68	21, 48, 106, 178	0
1	B	762/783 (97%)	-0.34	11 (1%) 75 77	21, 49, 104, 171	0
2	C	0/3	-	-	-	-
2	D	0/3	-	-	-	-
All	All	1520/1572 (96%)	-0.32	26 (1%) 70 72	21, 49, 106, 178	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	319	PHE	5.7
1	A	256	THR	5.6
1	A	254	ILE	4.3
1	A	257	LYS	4.2
1	A	253	ASP	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	B	872	1/1	0.84	0.24	69,69,69,69	0
3	MG	A	872	1/1	0.95	0.18	60,60,60,60	0
4	ADP	B	873	27/27	0.96	0.14	49,66,76,78	0
4	ADP	A	873	27/27	0.97	0.15	50,73,78,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.