



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 6, 2021 – 04:09 PM EST

PDB ID : 7JZ0
Title : Crystal Structure of SARS-CoV-2 Nsp16/10 Heterodimer in Complex with (m7GpppA2m)pUpUpApApA (Cap-1) and S-Adenosyl-L-homocysteine (SAH).
Authors : Minasov, G.; Shuvalova, L.; Rosas-Lemus, M.; Kiryukhina, O.; Brunzelle, J.S.; Satchell, K.J.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2020-09-01
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.24
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)
RNA backbone	3102	1060 (2.60-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	<div> <div>13%</div> <div>93%</div> <div>7%</div> </div>
1	C	300	<div> <div>11%</div> <div>93%</div> <div>6%</div> </div>
2	B	141	<div> <div>31%</div> <div>89%</div> <div>9%</div> </div>
2	D	141	<div> <div>22%</div> <div>79%</div> <div>18%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	7	<div><div></div><div>14%</div><div>29%</div><div>14%</div><div>14%</div><div>43%</div></div>
3	F	7	<div><div></div><div>29%</div><div>29%</div><div>14%</div><div>29%</div></div>

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2'-O-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	10	0
			2427	1549	406	454	18			
1	C	298	Total	C	N	O	S	0	6	0
			2386	1521	403	445	17			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	6797	SER	-	expression tag	UNP P0DTD1
A	6798	ASN	-	expression tag	UNP P0DTD1
C	6797	SER	-	expression tag	UNP P0DTD1
C	6798	ASN	-	expression tag	UNP P0DTD1

- Molecule 2 is a protein called Non-structural protein 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	129	Total	C	N	O	S	0	0	0
			960	598	161	184	17			
2	D	115	Total	C	N	O	S	0	1	0
			855	531	144	165	15			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	4252	SER	-	expression tag	UNP P0DTD1
B	4253	ASN	-	expression tag	UNP P0DTD1
D	4252	SER	-	expression tag	UNP P0DTD1
D	4253	ASN	-	expression tag	UNP P0DTD1

- Molecule 3 is a RNA chain called RNA (5'-D*(M7G))-R(P*(A2M)P*UP*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	P	0	0	0
			92	40	14	33	5			
3	F	5	Total	C	N	O	P	0	0	0
			96	40	14	36	6			

- Molecule 4 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: $C_{14}H_{20}N_6O_5S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
4	C	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Na	0	0
			2	2		
5	C	2	Total	Na	0	0
			2	2		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Cl	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	5	Total	Cl	0	0
			5	5		

- Molecule 7 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			3	1	2		
7	A	1	Total	C	O	0	0
			3	1	2		
7	A	1	Total	C	O	0	0
			3	1	2		
7	B	1	Total	C	O	0	0
			3	1	2		
7	B	1	Total	C	O	0	0
			3	1	2		
7	B	1	Total	C	O	0	0
			3	1	2		
7	C	1	Total	C	O	0	0
			3	1	2		
7	C	1	Total	C	O	0	0
			3	1	2		

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	2	Total 2	Zn 2	0	0
8	D	2	Total 2	Zn 2	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	151	Total 154	O 154	0	6
9	B	22	Total 22	O 22	0	0
9	C	165	Total 165	O 165	0	0
9	D	41	Total 43	O 43	0	2
9	E	7	Total 7	O 7	0	0
9	F	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2'-O-methyltransferase




- Molecule 1: 2'-O-methyltransferase




- Molecule 2: Non-structural protein 10




- Molecule 2: Non-structural protein 10







● Molecule 3: RNA (5'-D*(M7G))-R(P*(A2M)P*UP*U)-3')



● Molecule 3: RNA (5'-D*(M7G))-R(P*(A2M)P*UP*U)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	167.24Å 167.24Å 98.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.15 30.00 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.15) 99.9 (30.00-2.15)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.8.0266	Depositor
R, R_{free}	0.173 , 0.198 0.179 , 0.202	Depositor DCC
R_{free} test set	4338 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	42.2	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7305	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M7G, ZN, CL, FMT, SAH, NA, A2M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2479	0.75	0/3362
1	C	0.63	0/2438	0.75	0/3306
2	B	0.73	0/981	0.74	0/1329
2	D	0.72	0/874	0.72	0/1186
3	E	0.29	0/44	0.69	0/68
3	F	0.31	0/48	0.77	0/75
All	All	0.66	0/6864	0.75	0/9326

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2427	0	2410	15	0
1	C	2386	0	2367	13	0
2	B	960	0	921	2	0
2	D	855	0	811	3	0
3	E	92	0	50	1	0
3	F	96	0	49	2	0
4	A	26	0	19	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	26	0	19	0	0
5	A	2	0	0	0	0
5	C	2	0	0	0	0
6	A	2	0	0	0	0
6	C	5	0	0	0	0
7	A	9	0	3	0	0
7	B	9	0	3	0	0
7	C	6	0	2	0	0
8	B	2	0	0	0	0
8	D	2	0	0	0	0
9	A	154	0	0	5	0
9	B	22	0	0	0	0
9	C	165	0	0	0	0
9	D	43	0	0	0	0
9	E	7	0	0	0	0
9	F	7	0	0	0	0
All	All	7305	0	6654	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6940[B]:GLU:OE2	9:A:7201:HOH:O	1.96	0.82
1:A:6890[A]:GLY:O	1:C:7096:ASN:ND2	2.35	0.58
1:C:6800:SER:HA	1:C:6803:TRP:CZ3	2.39	0.57
1:A:7014:ARG:NH1	9:A:7203:HOH:O	2.39	0.54
1:A:7033:ASN:ND2	9:A:7204:HOH:O	2.42	0.52
1:C:6930:TYR:CD1	3:F:1:A2M:C4	2.94	0.51
1:A:6930:TYR:CD1	3:E:1:A2M:C4	2.95	0.50
1:A:6915:THR:HB	1:A:7091:ASP:HB2	1.94	0.50
1:A:7087:VAL:HG13	1:C:7091:ASP:HB3	1.93	0.49
1:A:6823:CYS:HB2	1:A:7027[A]:ILE:HD12	1.96	0.47
1:C:6884:ARG:HH22	1:C:6906:ASP:CG	2.17	0.47
1:A:6961:LEU:HB2	1:A:7080:ILE:HB	1.98	0.46
1:C:6902:VAL:CG1	2:D:4324:ALA:HB1	2.46	0.45
1:A:6902:VAL:HG12	2:B:4324:ALA:HB1	1.99	0.44
1:A:7076:GLY:HA2	9:A:7255:HOH:O	2.17	0.44
1:A:6884:ARG:HH12	1:A:6906:ASP:CG	2.20	0.44
1:A:7009:TYR:CZ	1:A:7011:GLY:HA2	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:4328:LEU:HD22	2:B:4365:LEU:HD11	2.01	0.43
1:C:6997:ALA:HB1	1:C:7035:ILE:HB	2.01	0.43
1:A:6946:GLY:N	9:A:7209:HOH:O	2.49	0.43
1:C:6961:LEU:HB2	1:C:7080:ILE:HB	1.99	0.43
2:D:4325:SER:HA	2:D:4331:ARG:HD3	2.02	0.42
1:C:6841:ASN:ND2	3:F:2:U:H4'	2.35	0.41
1:A:6931:ASP:OD1	1:A:6933:LYS:HB3	2.21	0.41
1:C:6902:VAL:HG12	2:D:4324:ALA:HB1	2.02	0.41
1:C:6937:VAL:HG13	1:C:6939:LYS:HG2	2.03	0.41
1:C:7009:TYR:CZ	1:C:7011:GLY:HA2	2.56	0.40
1:C:7058:MET:O	1:C:7080:ILE:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/300 (102%)	295 (96%)	12 (4%)	0	100	100
1	C	302/300 (101%)	291 (96%)	11 (4%)	0	100	100
2	B	127/141 (90%)	119 (94%)	8 (6%)	0	100	100
2	D	114/141 (81%)	107 (94%)	7 (6%)	0	100	100
All	All	850/882 (96%)	812 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/260 (103%)	268 (100%)	0	100	100
1	C	263/260 (101%)	263 (100%)	0	100	100
2	B	107/115 (93%)	107 (100%)	0	100	100
2	D	94/115 (82%)	94 (100%)	0	100	100
All	All	732/750 (98%)	732 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	6972	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	E	2/7 (28%)	1 (50%)	0
3	F	2/7 (28%)	1 (50%)	0
All	All	4/14 (28%)	2 (50%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	E	3	U
3	F	3	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	A2M	F	1	3	18,25,26	1.11	2 (11%)	18,36,39	1.42	3 (16%)
3	A2M	E	1	3	18,25,26	1.20	3 (16%)	18,36,39	1.39	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A2M	F	1	3	-	0/5/27/28	0/3/3/3
3	A2M	E	1	3	-	0/5/27/28	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1	A2M	C5-C4	2.64	1.47	1.40
3	F	1	A2M	C5-C4	2.63	1.47	1.40
3	E	1	A2M	O4'-C1'	2.49	1.44	1.41
3	F	1	A2M	C2-N3	2.18	1.35	1.32
3	E	1	A2M	C2-N3	2.12	1.35	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1	A2M	N3-C2-N1	-4.14	122.21	128.68
3	E	1	A2M	N3-C2-N1	-3.99	122.45	128.68
3	E	1	A2M	C4-C5-N7	-2.58	106.72	109.40
3	F	1	A2M	C2-N1-C6	2.28	122.65	118.75
3	F	1	A2M	C4-C5-N7	-2.18	107.12	109.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	1	A2M	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1	A2M	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 15 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SAH	A	7101	-	21,28,28	0.62	0	20,40,40	0.94	1 (5%)
4	SAH	C	7101	-	21,28,28	0.65	0	20,40,40	0.90	2 (10%)
7	FMT	B	4404	-	0,2,2	-	-	0,1,1	-	-
7	FMT	B	4405	-	0,2,2	-	-	0,1,1	-	-
7	FMT	C	7109	-	0,2,2	-	-	0,1,1	-	-
7	FMT	A	7107	-	0,2,2	-	-	0,1,1	-	-
7	FMT	C	7110	-	0,2,2	-	-	0,1,1	-	-
7	FMT	B	4403	-	0,2,2	-	-	0,1,1	-	-
7	FMT	A	7106	-	0,2,2	-	-	0,1,1	-	-
7	FMT	A	7108	-	0,2,2	-	-	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAH	C	7101	-	-	0/7/31/31	0/3/3/3
4	SAH	A	7101	-	-	0/7/31/31	0/3/3/3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	7101	SAH	C5-C6-N6	2.51	124.17	120.35
4	C	7101	SAH	C5-C6-N6	2.05	123.47	120.35
4	C	7101	SAH	O4'-C1'-C2'	-2.01	103.98	106.93


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/300 (99%)	0.68	39 (13%) 3 4	30, 47, 93, 137	0
1	C	298/300 (99%)	0.49	33 (11%) 5 7	30, 45, 82, 137	0
2	B	129/141 (91%)	1.80	44 (34%) 0 0	45, 79, 130, 139	0
2	D	115/141 (81%)	1.36	31 (26%) 0 0	38, 73, 131, 161	0
3	E	2/7 (28%)	1.27	1 (50%) 0 0	118, 118, 118, 137	0
3	F	3/7 (42%)	0.62	0 100 100	80, 80, 106, 123	0
All	All	846/896 (94%)	0.88	148 (17%) 1 1	30, 52, 113, 161	0

All (148) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	4273	ALA	14.3
1	A	6936	ASN	11.4
2	D	4339	PRO	9.8
2	D	4274	VAL	9.3
2	B	4285	ALA	8.2
1	C	6937	VAL	8.1
2	B	4339	PRO	8.1
2	B	4384	ASP	7.9
2	D	4383	CYS	7.7
2	B	4383	CYS	7.4
2	D	4385	GLN	7.2
2	B	4272	PHE	7.2
2	B	4287	GLY	7.1
2	B	4273	ALA	6.8
2	D	4372	VAL	6.8
2	B	4386	LEU	6.7
2	D	4384	ASP	6.6
2	B	4284	LEU	6.2
2	B	4385	GLN	6.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	4340	LYS	6.1
2	D	4272	PHE	6.0
1	A	6798	ASN	6.0
1	C	7062	GLU	5.8
1	A	6799	SER	5.8
2	D	4287	GLY	5.8
2	B	4271	ALA	5.8
2	B	4265	THR	5.8
2	B	4342	PHE	5.8
2	D	4342	PHE	5.6
1	C	6936	ASN	5.5
2	B	4382	SER	5.5
2	B	4372	VAL	5.4
2	B	4288	GLY	5.4
2	B	4373	CYS	5.4
1	A	6937	VAL	5.4
2	B	4381	CYS	5.4
1	A	7061	LYS	5.3
1	C	6799	SER	5.3
2	D	4285	ALA	5.3
2	B	4286	SER	5.2
2	B	4380	GLY	5.1
2	D	4286	SER	5.1
2	D	4382	SER	5.0
2	B	4268	SER	5.0
2	D	4371	THR	5.0
2	B	4274	VAL	4.9
1	C	6938	THR	4.7
2	D	4288	GLY	4.7
2	D	4284	LEU	4.6
2	B	4371	THR	4.6
2	B	4388	GLU	4.5
2	B	4387	ARG	4.5
1	C	7004	LEU	4.5
1	A	6933	LYS	4.4
1	C	7065	ILE	4.4
2	D	4377	LYS	4.4
2	B	4267	LEU	4.4
1	A	7003	PHE	4.3
1	A	7062	GLU	4.2
2	B	4341	GLY	4.2
1	C	6967	ILE	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	4278	LYS	4.1
2	D	4340	LYS	4.1
2	B	4289	GLN	4.1
2	D	4341	GLY	4.1
1	A	7005	ILE	4.0
1	C	7093	LEU	4.0
1	C	7003	PHE	3.9
2	B	4269	PHE	3.9
2	D	4275	ASP	3.9
2	D	4373	CYS	3.9
2	B	4338	ASN	3.9
1	C	7096	ASN	3.8
2	D	4282	ASP	3.7
1	C	6803	TRP	3.7
1	A	6966	ALA	3.7
1	C	7061	LYS	3.7
1	C	7063	GLY	3.7
1	C	6800	SER	3.6
1	A	6926	ILE	3.6
1	A	6969	ILE	3.6
1	C	6933	LYS	3.6
2	D	4338	ASN	3.6
2	B	4282	ASP	3.5
1	C	6969	ILE	3.5
1	A	6967	ILE	3.5
2	B	4266	VAL	3.5
2	B	4278	LYS	3.4
1	A	6829	GLY	3.4
2	B	4264	SER	3.3
2	D	4381	CYS	3.3
2	B	4283	TYR	3.3
2	D	4283	TYR	3.2
1	C	6925	ILE	3.2
1	C	6966	ALA	3.1
1	A	6932	PRO	3.1
1	C	7002	ALA	3.1
1	C	6848	LEU	3.1
2	D	4289	GLN	3.1
1	A	7085	ARG	3.1
1	C	6926	ILE	3.1
1	C	6965	VAL	2.9
1	A	6831	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	7064	GLN	2.9
1	A	6924	LEU	2.9
1	A	7004	LEU	2.8
1	A	6832	ALA	2.8
1	A	7096	ASN	2.8
1	A	7002	ALA	2.7
1	A	7067	ASP	2.7
1	A	6848	LEU	2.7
2	B	4370	CYS	2.7
2	D	4370	CYS	2.7
1	A	7063	GLY	2.6
2	D	4337	PRO	2.6
1	A	6935	LYS	2.6
1	C	6826	GLN	2.6
2	B	4337	PRO	2.6
2	B	4379	TYR	2.5
1	A	6845	TYR	2.5
1	C	6828	TYR	2.5
1	A	6925	ILE	2.5
2	B	4344	ASP	2.5
1	A	6938	THR	2.4
1	A	7064	GLN	2.4
1	C	7005	ILE	2.4
2	B	4369	VAL	2.4
2	D	4271	ALA	2.4
1	C	6934	THR	2.3
1	C	7095	ASN	2.3
1	A	6830	ASP	2.3
1	A	6991	PHE	2.2
1	A	7066	ASN	2.2
1	C	6883	LEU	2.2
1	A	7084	ASN	2.2
2	D	4280	TYR	2.2
2	B	4392	GLN	2.1
2	B	4378	GLY	2.1
2	B	4304	THR	2.1
1	C	6866	ILE	2.1
1	A	6970	THR	2.1
3	E	3	U	2.1
1	A	6879	GLY	2.0
1	A	6939	LYS	2.0
1	C	6968	LYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	6881	ALA	2.0
1	A	6846	THR	2.0
1	C	6990	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A2M	E	1	23/24	0.87	0.29	80,90,105,111	0
3	A2M	F	1	23/24	0.93	0.16	56,62,75,76	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	FMT	B	4404	3/3	0.51	0.26	88,88,91,97	0
7	FMT	B	4403	3/3	0.59	0.32	82,82,84,85	0
6	CL	A	7105	1/1	0.70	0.34	81,81,81,81	0
7	FMT	B	4405	3/3	0.77	0.22	94,94,95,97	0
6	CL	C	7105	1/1	0.81	0.10	79,79,79,79	0
7	FMT	C	7110	3/3	0.81	0.30	100,100,100,101	0
5	NA	A	7103	1/1	0.84	0.07	67,67,67,67	0
6	CL	A	7104	1/1	0.84	0.08	85,85,85,85	0
6	CL	C	7108	1/1	0.85	0.26	93,93,93,93	0
6	CL	C	7107	1/1	0.86	0.14	89,89,89,89	0
5	NA	C	7102	1/1	0.88	0.16	61,61,61,61	0
5	NA	A	7102	1/1	0.90	0.17	69,69,69,69	0
6	CL	C	7106	1/1	0.92	0.35	94,94,94,94	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	FMT	A	7107	3/3	0.92	0.22	75,75,78,82	0
6	CL	C	7104	1/1	0.92	0.17	73,73,73,73	0
7	FMT	A	7108	3/3	0.93	0.41	83,83,89,90	0
8	ZN	B	4401	1/1	0.94	0.07	63,63,63,63	0
5	NA	C	7103	1/1	0.95	0.18	58,58,58,58	0
4	SAH	C	7101	26/26	0.95	0.10	41,44,49,50	0
4	SAH	A	7101	26/26	0.95	0.08	40,46,53,54	0
7	FMT	A	7106	3/3	0.96	0.08	71,71,75,81	0
8	ZN	B	4402	1/1	0.96	0.11	71,71,71,71	1
8	ZN	D	4402	1/1	0.96	0.14	79,79,79,79	1
8	ZN	D	4401	1/1	0.97	0.08	57,57,57,57	0
7	FMT	C	7109	3/3	0.99	0.08	47,47,52,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN B 4401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SAH C 7101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around SAH A 7101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 4402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 4402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN D 4401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.