



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 31, 2021 – 12:04 PM EDT

PDB ID : 7K29  
Title : Kelch domain of human KEAP1 bound to Nrf2 peptide, LDEETGEAL  
Authors : Muellers, S.N.; Allen, K.N.  
Deposited on : 2020-09-08  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.18  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.18


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	<div> <div>18%</div> <div> <div></div> <div>72%</div> <div>21%</div> <div>• 6%</div> </div> </div>
1	B	301	<div> <div>10%</div> <div> <div></div> <div>79%</div> <div>16%</div> <div>• •</div> </div> </div>
2	P	11	<div> <div></div> <div> <div></div> <div>82%</div> <div>18%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4554 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kelch-like ECH-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2170	1351	392	412	15			
1	B	290	Total	C	N	O	S	0	1	0
			2229	1385	405	423	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	540	ALA	GLU	engineered mutation	UNP Q14145
A	542	ALA	GLU	engineered mutation	UNP Q14145
A	613	SER	CYS	conflict	UNP Q14145
B	540	ALA	GLU	engineered mutation	UNP Q14145
B	542	ALA	GLU	engineered mutation	UNP Q14145
B	613	SER	CYS	conflict	UNP Q14145

- Molecule 2 is a protein called ACE-LEU-ASP-GLU-GLU-THR-GLY-GLU-ALA-LEU-NH2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	P	11	Total	C	N	O	0	0	1
			71	42	10	19			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	12	Total	O	0	0
			12	12		
4	B	59	Total	O	0	0
			59	59		
4	P	3	Total	O	0	0
			3	3		



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	162.55Å 68.92Å 77.07Å 90.00° 117.70° 90.00°	Depositor
Resolution (Å)	29.37 – 2.20 29.37 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.37-2.20) 99.4 (29.37-2.20)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.80 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.229 , 0.269 0.229 , 0.269	Depositor DCC
$R_{free}$ test set	2013 reflections (5.26%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.7	Xtriage
Anisotropy	0.556	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 38.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.006 for -h-2*k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4554	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NH2, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/2223	0.61	0/3029
1	B	0.48	0/2286	0.67	2/3112 (0.1%)
2	P	0.68	0/67	0.76	0/90
All	All	0.44	0/4576	0.65	2/6231 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	498	ARG	NE-CZ-NH1	-6.95	116.82	120.30
1	B	336	ARG	NE-CZ-NH1	-5.71	117.45	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2170	0	2066	41	1
1	B	2229	0	2125	28	1
2	P	71	0	62	2	0
3	A	10	0	0	0	0
4	A	12	0	0	0	0
4	B	59	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	P	3	0	0	0	0
All	All	4554	0	4253	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:466:ALA:HB1	1:B:514:VAL:HG23	1.70	0.74
1:A:573:ASP:HB3	1:A:578:LEU:HD11	1.70	0.72
1:A:522:ALA:HB1	1:A:550:MET:HE3	1.80	0.64
1:A:441:GLU:HB3	1:A:452:LEU:HD23	1.78	0.64
1:A:346:ASN:HB3	1:A:349:ASP:OD1	2.01	0.59
1:A:447:ARG:HB3	1:A:449:GLU:OE2	2.02	0.59
1:A:565:ARG:HH11	1:A:565:ARG:HG3	1.69	0.57
1:A:525:TYR:CE2	2:P:79:GLU:HG3	2.40	0.57
1:A:329:TYR:HE1	1:A:355:LEU:HD11	1.69	0.56
1:A:365:LEU:HD23	1:A:365:LEU:H	1.71	0.56
1:A:373:LEU:HD22	1:A:397:ASN:HA	1.88	0.55
1:B:557:LEU:H	1:B:557:LEU:HD23	1.71	0.55
1:A:327:LEU:N	1:A:609:THR:O	2.39	0.55
1:A:472:LEU:HB3	1:A:490:TYR:HB3	1.89	0.54
1:A:515:LEU:HD22	1:A:566:ILE:HG13	1.90	0.54
1:A:566:ILE:HB	1:A:584:TYR:HB3	1.91	0.52
1:A:366:ALA:HB3	1:A:418:VAL:HG13	1.91	0.51
1:B:426:TYR:CZ	1:B:442:ARG:HD3	2.45	0.51
1:B:421:ILE:HD11	1:B:472:LEU:HB2	1.93	0.51
1:A:565:ARG:HG3	1:A:565:ARG:NH1	2.26	0.51
1:A:554:ARG:HG3	1:A:557:LEU:HD22	1.93	0.50
1:B:614:ARG:NE	1:B:614:ARG:HA	2.27	0.50
1:B:515:LEU:HD22	1:B:566:ILE:HG13	1.93	0.50
1:A:562:HIS:HB3	1:A:567:TYR:CE1	2.47	0.49
1:B:512:VAL:HA	1:B:520:TYR:O	2.12	0.49
1:A:467:VAL:O	1:A:514:VAL:HG21	2.12	0.49
1:B:519:ILE:O	1:B:536:ARG:HA	2.13	0.49
1:A:354:ARG:O	1:A:355:LEU:HD23	2.13	0.48
1:B:397:ASN:OD1	1:B:399:MET:HB2	2.12	0.48
1:B:436:HIS:ND1	1:B:461:ILE:HD11	2.27	0.48
1:B:335:PHE:O	1:B:336:ARG:HB2	2.13	0.48
1:A:397:ASN:OD1	1:A:399:MET:HG2	2.14	0.48

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:566:ILE:O	1:A:583:CYS:HA	2.13	0.47
1:B:446:GLU:H	1:B:446:GLU:CD	2.17	0.47
1:A:339:LEU:HD22	1:A:341:TYR:CE2	2.49	0.47
1:B:444:GLU:HB3	1:B:446:GLU:OE1	2.15	0.47
1:A:519:ILE:O	1:A:536:ARG:HA	2.15	0.46
1:A:550:MET:CE	1:A:568:VAL:HG11	2.45	0.46
1:B:566:ILE:HB	1:B:584:TYR:HB3	1.97	0.46
1:B:467:VAL:O	1:B:514:VAL:HG21	2.15	0.46
1:A:425:ILE:HB	1:A:443:TYR:HB3	1.97	0.46
1:B:416:ILE:HD11	1:B:427:ALA:HB1	1.98	0.46
1:B:365:LEU:HD12	1:B:376:ALA:HB1	1.99	0.45
1:A:447:ARG:O	1:A:449:GLU:HG3	2.17	0.45
1:B:491:TYR:O	1:B:495:ASN:N	2.50	0.45
1:B:579:ASP:OD1	1:B:596:ARG:HD2	2.18	0.43
1:A:478:PHE:CD1	1:A:483:ARG:HG3	2.53	0.43
1:A:354:ARG:NH1	1:A:357:ASP:OD1	2.52	0.43
1:A:541:THR:O	1:A:543:THR:HG23	2.18	0.43
1:A:334:TYR:CG	2:P:83:GLU:HG2	2.54	0.43
1:B:581:VAL:HB	1:B:595:THR:HG22	2.01	0.43
1:A:369:VAL:HG23	1:A:607:ALA:HB1	2.02	0.42
1:A:369:VAL:HG21	1:A:608:VAL:O	2.19	0.42
1:B:584:TYR:HB2	1:B:591:TRP:CZ3	2.55	0.42
1:A:460:ARG:HB3	1:A:463:VAL:HB	2.02	0.42
1:A:483:ARG:HG2	1:A:508:SER:HB2	2.01	0.42
1:A:570:GLY:HA2	1:A:578:LEU:O	2.20	0.42
1:A:333:GLY:HA3	1:A:604:VAL:HG12	2.01	0.42
1:B:347:PRO:HG2	1:B:562:HIS:CD2	2.54	0.41
1:B:447:ARG:O	1:B:449:GLU:HG3	2.19	0.41
1:B:381:ASN:O	1:B:387:ASN:HA	2.21	0.41
1:A:584:TYR:HB2	1:A:591:TRP:CD2	2.56	0.41
1:B:490:TYR:HB2	1:B:497:TRP:CZ2	2.55	0.41
1:B:424:HIS:HD2	4:B:749:HOH:O	2.03	0.41
1:B:409:MET:SD	1:B:413:ARG:HD2	2.61	0.40
1:B:369:VAL:HG21	1:B:609:THR:HB	2.01	0.40
1:A:362:ARG:NH2	1:A:394:ASP:OD1	2.55	0.40
1:A:491:TYR:CE1	1:A:498:ARG:HG3	2.56	0.40
1:A:512:VAL:HA	1:A:520:TYR:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359:GLN:O	1:B:336:ARG:NH1[1_556]	1.90	0.30

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/301 (93%)	265 (94%)	15 (5%)	1 (0%)	34	37
1	B	289/301 (96%)	282 (98%)	7 (2%)	0	100	100
2	P	9/11 (82%)	9 (100%)	0	0	100	100
All	All	579/613 (94%)	556 (96%)	22 (4%)	1 (0%)	47	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	359	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/245 (93%)	226 (99%)	2 (1%)	78	88
1	B	235/245 (96%)	230 (98%)	5 (2%)	53	67
2	P	7/7 (100%)	7 (100%)	0	100	100
All	All	470/497 (95%)	463 (98%)	7 (2%)	65	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	354	ARG
1	A	557	LEU
1	B	470	ARG
1	B	528	GLN
1	B	557	LEU
1	B	613	SER
1	B	614	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	401	ASN
1	B	402	GLN
1	B	528	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	702	-	4,4,4	0.28	0	6,6,6	0.39	0
3	SO4	A	701	-	4,4,4	0.11	0	6,6,6	0.50	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.









## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	283/301 (94%)	0.89	53 (18%)  	28, 56, 92, 110	0
1	B	290/301 (96%)	0.62	30 (10%)  	26, 37, 60, 92	0
2	P	9/11 (81%)	0.18	0  	36, 41, 49, 55	0
All	All	582/613 (94%)	0.75	83 (14%)  	26, 43, 86, 110	0

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	385	ASP	5.5
1	A	353	LEU	5.3
1	B	366	ALA	5.1
1	B	384	PRO	4.9
1	A	447	ARG	4.9
1	A	400	THR	4.8
1	A	349	ASP	4.6
1	A	351	THR	4.6
1	A	563	GLN	4.3
1	B	512	VAL	4.2
1	A	402	GLN	4.0
1	B	447	ARG	4.0
1	A	399	MET	3.9
1	B	365	LEU	3.8
1	A	423	GLY	3.8
1	A	365	LEU	3.7
1	A	446	GLU	3.7
1	A	348	SER	3.5
1	A	452	LEU	3.5
1	A	604	VAL	3.5
1	B	465	VAL	3.5
1	A	416	ILE	3.5
1	A	605	GLY	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	399	MET	3.4
1	A	588	THR	3.3
1	A	371	GLY	3.3
1	A	347	PRO	3.3
1	A	449	GLU	3.2
1	A	397	ASN	3.2
1	B	605	GLY	3.2
1	A	567	TYR	3.1
1	A	557	LEU	3.1
1	B	386	GLY	3.1
1	B	464	GLY	3.0
1	B	416	ILE	2.9
1	A	494	ARG	2.9
1	A	463	VAL	2.9
1	A	367	GLY	2.9
1	A	590	THR	2.8
1	A	401	ASN	2.8
1	B	463	VAL	2.8
1	B	606	VAL	2.7
1	A	451	HIS	2.7
1	A	516	HIS	2.7
1	B	377	VAL	2.7
1	B	336	ARG	2.6
1	A	510	ALA	2.6
1	B	418	VAL	2.6
1	A	512	VAL	2.6
1	B	604	VAL	2.6
1	B	422	ASP	2.5
1	A	559	ILE	2.5
1	A	421	ILE	2.5
1	A	366	ALA	2.5
1	A	422	ASP	2.5
1	A	398	PRO	2.4
1	A	408	PRO	2.4
1	B	379	GLY	2.4
1	B	367	GLY	2.4
1	A	569	LEU	2.4
1	B	434	CYS	2.4
1	A	418	VAL	2.4
1	B	588	THR	2.4
1	B	587	ASP	2.4
1	A	331	ALA	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	417	GLY	2.4
1	A	424	HIS	2.4
1	A	561	VAL	2.4
1	A	377	VAL	2.4
1	B	576	THR	2.3
1	B	419	GLY	2.3
1	A	575	HIS	2.3
1	B	326	ARG	2.3
1	A	576	THR	2.3
1	A	329	TYR	2.2
1	B	417	GLY	2.2
1	A	493	GLU	2.2
1	A	327	LEU	2.2
1	A	330	THR	2.1
1	A	350	GLY	2.1
1	B	393	LEU	2.1
1	A	556	ALA	2.1
1	B	378	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	702	5/5	0.95	0.46	39,42,52,55	0
3	SO4	A	701	5/5	0.96	0.34	51,52,54,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.