



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 17, 2021 – 03:36 AM EDT

PDB ID : 1K3Z
Title : X-ray crystal structure of the I κ B β /NF- κ B p65 homodimer complex
Authors : Shiva, M.; Huang, D.B.; Chen, Y.; Huxford, T.; Ghosh, S.; Ghosh, G.
Deposited on : 2001-10-04
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	136	<div> <div>54%</div> <div>39%</div> <div>.</div> <div>.</div> </div>
1	B	136	<div> <div>54%</div> <div>29%</div> <div>.</div> <div>13%</div> </div>
2	D	282	<div> <div>42%</div> <div>31%</div> <div>5%</div> <div>22%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3769 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription factor p65.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	130	Total	C	N	O	S	0	0	0
			1056	660	191	199	6			
1	B	118	Total	C	N	O	S	0	0	0
			953	590	175	183	5			

- Molecule 2 is a protein called transcription factor inhibitor I-kappa-B-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	221	Total	C	N	O	S	0	0	0
			1661	1041	305	309	6			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	312	GLU	SER	engineered mutation	UNP Q60778
D	313	GLU	SER	engineered mutation	UNP Q60778
D	314	GLU	SER	engineered mutation	UNP Q60778
D	316	GLU	SER	engineered mutation	UNP Q60778
D	318	GLU	SER	engineered mutation	UNP Q60778

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total	O	0	0
			27	27		
3	B	23	Total	O	0	0
			23	23		
3	D	49	Total	O	0	0
			49	49		

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.40 Å 48.92 Å 59.47 Å 95.17° 91.80° 105.45°	Depositor
Resolution (Å)	30.00 – 2.50	Depositor
% Data completeness (in resolution range)	80.0 (30.00-2.50)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.181 , 0.256	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3769	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/1079	0.62	0/1453
1	B	0.37	0/972	0.65	0/1312
2	D	0.35	0/1696	0.62	1/2309 (0.0%)
All	All	0.36	0/3747	0.63	1/5074 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	238	PRO	N-CA-C	5.08	125.30	112.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1056	0	1037	82	0
1	B	953	0	927	55	0
2	D	1661	0	1642	98	0
3	A	27	0	0	1	0
3	B	23	0	0	4	0
3	D	49	0	0	3	0
All	All	3769	0	3606	209	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 209 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:ARG:HG3	1:B:248:VAL:HG21	1.34	1.07
2:D:55:THR:HG22	2:D:59:ASP:H	1.18	1.02
1:A:218:LYS:NZ	1:B:246:ARG:HE	1.63	0.95
1:A:278:ARG:HH11	1:A:278:ARG:HB3	1.30	0.94
2:D:55:THR:HG23	2:D:57:ASP:H	1.35	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	128/136 (94%)	117 (91%)	10 (8%)	1 (1%)	19	35
1	B	116/136 (85%)	101 (87%)	14 (12%)	1 (1%)	17	31
2	D	217/282 (77%)	195 (90%)	15 (7%)	7 (3%)	4	5
All	All	461/554 (83%)	413 (90%)	39 (8%)	9 (2%)	7	12

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	303	ASP
2	D	305	GLY
2	D	308	LEU
1	B	292	THR
2	D	306	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/123 (95%)	109 (93%)	8 (7%)	16	30
1	B	105/123 (85%)	95 (90%)	10 (10%)	8	17
2	D	168/225 (75%)	156 (93%)	12 (7%)	14	28
All	All	390/471 (83%)	360 (92%)	30 (8%)	13	25

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	296	HIS
2	D	278	LEU
1	B	304	ARG
2	D	308	LEU
2	D	155	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	287	GLN
2	D	69	GLN
2	D	286	ASN
2	D	95	GLN
2	D	140	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.