



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 03:26 pm BST

PDB ID : 4K30
Title : Structure of the N-acetyltransferase domain of human N-acetylglutamate synthase
Authors : Shi, D.; Zhao, G.; Jin, Z.; Allewell, N.M.; Tuchman, M.
Deposited on : 2013-04-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>• •</div> </div> </div>
1	B	160	<div> <div>5%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div>• 5%</div> </div> </div>
1	X	160	<div> <div>6%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div>• •</div> </div> </div>
1	Y	160	<div> <div>18%</div> <div> <div></div> <div>76%</div> <div>16%</div> <div>• 7%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N-acetylglutamate synthase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	0	0
			1251	802	224	221	4			
1	B	152	Total	C	N	O	S	0	0	0
			1245	799	223	219	4			
1	X	153	Total	C	N	O	S	0	0	0
			1251	802	224	221	4			
1	Y	149	Total	C	N	O	S	0	0	0
			1216	781	215	216	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	375	SER	-	EXPRESSION TAG	UNP Q8N159
A	376	HIS	-	EXPRESSION TAG	UNP Q8N159
B	375	SER	-	EXPRESSION TAG	UNP Q8N159
B	376	HIS	-	EXPRESSION TAG	UNP Q8N159
X	375	SER	-	EXPRESSION TAG	UNP Q8N159
X	376	HIS	-	EXPRESSION TAG	UNP Q8N159
Y	375	SER	-	EXPRESSION TAG	UNP Q8N159
Y	376	HIS	-	EXPRESSION TAG	UNP Q8N159

- Molecule 2 is N-ACETYL-L-GLUTAMATE (three-letter code: NLG) (formula: C₇H₁₁NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 13	C 7	N 1	O 5	0	0
2	B	1	Total 13	C 7	N 1	O 5	0	0
2	X	1	Total 13	C 7	N 1	O 5	0	0
2	Y	1	Total 13	C 7	N 1	O 5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	119	Total O 119 119	0	0
3	B	60	Total O 60 60	0	0
3	X	70	Total O 70 70	0	0
3	Y	17	Total O 17 17	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: N-acetylglutamate synthase, mitochondrial





- Molecule 1: N-acetylglutamate synthase, mitochondrial



- Molecule 1: N-acetylglutamate synthase, mitochondrial



- Molecule 1: N-acetylglutamate synthase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	116.14Å 116.14Å 109.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.46 – 2.10 39.88 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.46-2.10) 95.7 (39.88-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.185 , 0.244 0.186 , 0.245	Depositor DCC
R_{free} test set	1999 reflections (4.54%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5281	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NLG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/1286	0.63	2/1738 (0.1%)
1	B	0.36	0/1280	0.56	1/1730 (0.1%)
1	X	0.42	0/1286	0.59	0/1738
1	Y	0.29	0/1249	0.46	0/1689
All	All	0.40	0/5101	0.56	3/6895 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	414	ARG	NE-CZ-NH2	-5.85	117.38	120.30
1	B	509	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	A	414	ARG	NE-CZ-NH1	5.67	123.13	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	423	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1251	0	1226	18	0
1	B	1245	0	1221	17	0
1	X	1251	0	1226	17	0
1	Y	1216	0	1194	15	0
2	A	13	0	9	0	0
2	B	13	0	9	0	0
2	X	13	0	9	2	0
2	Y	13	0	9	0	0
3	A	119	0	0	2	0
3	B	60	0	0	3	0
3	X	70	0	0	5	0
3	Y	17	0	0	2	0
All	All	5281	0	4903	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:508:ILE:H	1:B:508:ILE:HD12	1.45	0.79
1:B:463:CYS:SG	3:B:732:HOH:O	2.43	0.76
1:A:508:ILE:H	1:A:508:ILE:HD12	1.51	0.73
1:Y:377:MET:N	3:Y:717:HOH:O	2.21	0.73
1:Y:451:ARG:NE	3:Y:706:HOH:O	1.95	0.71
1:Y:436:LEU:HD21	1:Y:510:ASP:HB3	1.71	0.71
1:B:495:ASN:OD1	3:B:751:HOH:O	2.11	0.69
1:X:390:ARG:NH1	3:X:758:HOH:O	2.25	0.69
1:X:405:ASP:H	1:X:527:LYS:HD2	1.57	0.68
1:A:422:GLU:O	3:A:778:HOH:O	2.11	0.67
1:X:401:LYS:NZ	2:X:600:NLG:OE2	2.28	0.67
1:X:477:VAL:O	3:X:746:HOH:O	2.13	0.66
1:A:402:LYS:H	1:A:527:LYS:NZ	1.95	0.65
1:X:458:GLN:OE1	1:X:484:TRP:NE1	2.31	0.62
1:X:390:ARG:CZ	3:X:758:HOH:O	2.46	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Y:442:LEU:HB3	1:Y:473:TRP:HB3	1.82	0.60
1:B:442:LEU:HB3	1:B:473:TRP:HB3	1.83	0.60
1:A:436:LEU:HD22	1:A:514:LEU:HD21	1.85	0.57
1:B:519:LYS:HD2	3:B:756:HOH:O	2.05	0.57
2:X:600:NLG:OE1	3:X:718:HOH:O	2.18	0.56
1:A:402:LYS:H	1:A:527:LYS:HZ3	1.56	0.53
1:A:442:LEU:HB3	1:A:473:TRP:HB3	1.90	0.52
1:B:392:VAL:HG13	1:B:403:LEU:HD12	1.89	0.52
1:Y:393:ASP:O	1:Y:397:ALA:N	2.42	0.50
1:Y:430:LEU:HD21	1:Y:468:LEU:HD11	1.93	0.50
1:A:393:ASP:OD2	1:X:385:LYS:NZ	2.42	0.49
1:A:527:LYS:HZ2	1:X:376:HIS:HB3	1.78	0.49
1:X:436:LEU:O	3:X:732:HOH:O	2.19	0.48
1:Y:436:LEU:HD22	1:Y:514:LEU:HD21	1.95	0.48
1:Y:430:LEU:CD2	1:Y:468:LEU:HD11	2.43	0.48
1:Y:414:ARG:HG2	1:Y:431:THR:HB	1.95	0.47
1:A:419:TYR:HB2	1:A:460:LEU:HD21	1.95	0.47
1:B:430:LEU:HD23	1:B:442:LEU:HA	1.96	0.47
1:A:525:PHE:O	1:A:527:LYS:HE3	2.15	0.47
1:B:507:ASP:OD1	1:B:509:ARG:HG2	2.14	0.47
1:A:523:ASP:HB3	1:A:526:HIS:HB2	1.96	0.47
1:A:512:TYR:HE1	1:B:508:ILE:HD13	1.80	0.47
1:A:508:ILE:H	1:A:508:ILE:CD1	2.16	0.46
1:Y:391:LEU:HD12	1:Y:424:TYR:HE2	1.79	0.46
1:Y:507:ASP:OD1	1:Y:509:ARG:HD2	2.17	0.45
1:Y:392:VAL:HG13	1:Y:403:LEU:HD12	1.99	0.44
1:X:479:ASN:CG	1:X:481:ILE:HG22	2.37	0.44
1:B:394:LEU:HD22	1:B:424:TYR:O	2.18	0.44
1:Y:456:SER:O	1:Y:460:LEU:HB2	2.17	0.44
1:X:380:VAL:HG21	1:X:386:LEU:HG	2.00	0.43
1:X:442:LEU:HB3	1:X:473:TRP:HB3	1.99	0.43
1:B:509:ARG:H	1:B:509:ARG:HG2	1.69	0.43
1:A:385:LYS:HD3	1:A:385:LYS:HA	1.85	0.43
1:A:527:LYS:NZ	1:X:376:HIS:HB3	2.34	0.43
1:Y:394:LEU:HG	1:Y:446:VAL:HG11	2.01	0.42
1:A:449:SER:HB3	1:X:390:ARG:HG3	2.02	0.42
1:B:508:ILE:H	1:B:508:ILE:CD1	2.14	0.41
1:A:502:TRP:CD1	1:A:505:LEU:HD22	2.55	0.41
1:X:441:TYR:CE2	1:X:443:ASP:HA	2.55	0.41
1:B:404:ARG:HG3	1:B:524:SER:O	2.21	0.41
1:B:500:PHE:CZ	1:B:519:LYS:HG3	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:423:GLY:HA3	1:X:425:ASN:ND2	2.37	0.40
1:B:495:ASN:HB3	1:B:497:GLN:H	1.86	0.40
1:Y:433:GLU:HB3	1:Y:517:HIS:HE2	1.85	0.40
1:A:376:HIS:HA	3:A:812:HOH:O	2.20	0.40
1:B:380:VAL:HG11	1:B:386:LEU:HG	2.03	0.40
1:B:384:ASP:OD1	1:B:412:ARG:NH2	2.55	0.40
1:X:526:HIS:O	1:X:526:HIS:ND1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/160 (94%)	147 (97%)	3 (2%)	1 (1%)	22	18
1	B	150/160 (94%)	149 (99%)	1 (1%)	0	100	100
1	X	151/160 (94%)	147 (97%)	4 (3%)	0	100	100
1	Y	147/160 (92%)	143 (97%)	4 (3%)	0	100	100
All	All	599/640 (94%)	586 (98%)	12 (2%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	422	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/140 (96%)	131 (97%)	4 (3%)	41	44
1	B	134/140 (96%)	134 (100%)	0	100	100
1	X	135/140 (96%)	133 (98%)	2 (2%)	65	71
1	Y	131/140 (94%)	126 (96%)	5 (4%)	33	34
All	All	535/560 (96%)	524 (98%)	11 (2%)	53	59

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	382	SER
1	A	436	LEU
1	A	505	LEU
1	A	508	ILE
1	X	403	LEU
1	X	458	GLN
1	Y	377	MET
1	Y	378	LEU
1	Y	460	LEU
1	Y	472	PHE
1	Y	481	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NLG	X	600	-	6,12,12	1.74	1 (16%)	7,15,15	1.17	0
2	NLG	B	600	-	6,12,12	2.01	3 (50%)	7,15,15	1.59	1 (14%)
2	NLG	A	600	-	6,12,12	2.11	2 (33%)	7,15,15	1.67	2 (28%)
2	NLG	Y	600	-	6,12,12	1.84	1 (16%)	7,15,15	1.93	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NLG	X	600	-	-	2/7/13/13	-
2	NLG	B	600	-	-	0/7/13/13	-
2	NLG	A	600	-	-	0/7/13/13	-
2	NLG	Y	600	-	-	3/7/13/13	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Y	600	NLG	C7-N2	3.70	1.47	1.34
2	A	600	NLG	C7-N2	3.62	1.46	1.34
2	B	600	NLG	C7-N2	3.43	1.46	1.34
2	X	600	NLG	C7-N2	3.31	1.45	1.34
2	A	600	NLG	CB-CA	-2.79	1.49	1.53
2	B	600	NLG	CB-CA	-2.45	1.50	1.53
2	B	600	NLG	CA-N2	-2.25	1.43	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	600	NLG	CA-N2-C7	3.32	126.09	122.44
2	Y	600	NLG	CG-CB-CA	3.19	119.48	113.04
2	B	600	NLG	CG-CB-CA	3.14	119.38	113.04
2	A	600	NLG	CG-CB-CA	2.81	118.71	113.04
2	Y	600	NLG	CA-N2-C7	-2.67	119.50	122.44

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Y	600	NLG	C-CA-CB-CG
2	Y	600	NLG	N2-CA-CB-CG
2	Y	600	NLG	C-CA-N2-C7
2	X	600	NLG	C-CA-CB-CG
2	X	600	NLG	N2-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	X	600	NLG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	153/160 (95%)	0.02	4 (2%) 56 61	23, 35, 61, 114	0
1	B	152/160 (95%)	0.32	8 (5%) 26 32	27, 54, 86, 108	0
1	X	153/160 (95%)	0.40	10 (6%) 18 23	30, 45, 77, 94	0
1	Y	149/160 (93%)	1.07	29 (19%) 1 1	44, 77, 114, 128	0
All	All	607/640 (94%)	0.45	51 (8%) 11 14	23, 50, 100, 128	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Y	378	LEU	6.3
1	Y	393	ASP	6.0
1	Y	383	LEU	5.7
1	B	376	HIS	4.5
1	B	452	GLN	4.2
1	Y	380	VAL	4.1
1	Y	386	LEU	4.1
1	Y	379	ARG	4.0
1	Y	412	ARG	3.7
1	B	386	LEU	3.7
1	X	451	ARG	3.7
1	X	453	GLY	3.7
1	Y	413	PRO	3.5
1	X	526	HIS	3.4
1	X	508	ILE	3.3
1	Y	496	LYS	3.1
1	Y	497	GLN	3.0
1	B	456	SER	2.9
1	Y	414	ARG	2.9
1	Y	498	TRP	2.9
1	Y	389	GLY	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	Y	415	LEU	2.9
1	Y	384	ASP	2.9
1	X	454	GLN	2.9
1	Y	420	VAL	2.8
1	Y	411	LEU	2.8
1	Y	381	ARG	2.7
1	A	375	SER	2.7
1	Y	404	ARG	2.6
1	Y	509	ARG	2.6
1	B	526	HIS	2.5
1	A	527	LYS	2.5
1	X	505	LEU	2.5
1	Y	418	ILE	2.4
1	Y	382	SER	2.4
1	B	405	ASP	2.4
1	Y	377	MET	2.3
1	X	511	SER	2.3
1	X	452	GLN	2.3
1	X	514	LEU	2.2
1	A	376	HIS	2.2
1	A	526	HIS	2.2
1	B	412	ARG	2.2
1	Y	409	ALA	2.2
1	B	459	MET	2.2
1	Y	406	ASP	2.2
1	Y	407	TYR	2.1
1	X	512	TYR	2.1
1	Y	424	TYR	2.1
1	Y	388	GLN	2.1
1	Y	426	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NLG	B	600	13/13	0.82	0.21	44,60,71,73	0
2	NLG	Y	600	13/13	0.82	0.22	55,81,90,92	0
2	NLG	X	600	13/13	0.84	0.18	36,44,48,51	0
2	NLG	A	600	13/13	0.90	0.16	25,29,38,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.