



wwPDB NMR Structure Validation Summary Report ⓘ

Jun 7, 2020 – 04:33 am BST

PDB ID : 6K3J
Title : Solution structure of APOBEC3G-CD2 with ssDNA, Product A
Authors : Cao, C.; Yan, X.; Lan, W.; Wang, C.
Deposited on : 2019-05-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 77%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	188	
2	B	10	

2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:197-A:384 (188)	0.71	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 8, 9, 12, 15, 17
2	2, 5, 10, 16, 19
3	3, 7, 11, 14, 20
4	4, 6
Single-model clusters	13; 18

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3332 atoms, of which 1576 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DNA dC->dU-editing enzyme APOBEC-3G.

Mol	Chain	Residues	Atoms							Trace
1	A	188	Total	C	H	N	O	S		0
			3020	985	1463	275	283	14		

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*TP*TP*CP*UP*(IUR)P*AP*AP*TP*T)-3').

Mol	Chain	Residues	Atoms								Trace
2	B	10	Total	C	H	I	N	O	P		0
			311	97	113	1	30	61	9		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
3	A	1	Total	Zn
			1	1

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3G




- Molecule 2: DNA (5'-D(*AP*TP*TP*CP*UP*(IUR)P*AP*AP*TP*T)-3')



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3G



- Molecule 2: DNA (5'-D(*AP*TP*TP*CP*UP*(IUR)P*AP*AP*TP*T)-3')

Chain B: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	
X-PLOR NIH	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	2221
Number of shifts mapped to atoms	2221
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	77%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1557	1463	1452	80±8
2	B	198	113	113	15±5
All	All	35120	31520	31300	1642

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

5 of 702 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:253:LEU:HD22	1:A:253:LEU:H	0.93	1.24	5	1
1:A:281:CYS:SG	1:A:309:ILE:HG23	0.86	2.09	6	6
1:A:299:ILE:HD12	1:A:331:ALA:O	0.84	1.73	12	14
1:A:208:ASN:O	1:A:212:VAL:HG13	0.83	1.73	14	3
1:A:209:GLU:O	1:A:212:VAL:HG13	0.81	1.75	15	3

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/188 (99%)	172±1 (92±1%)	10±2 (5±1%)	5±1 (2±0%)	9	45
All	All	3720/3760 (99%)	3430 (92%)	198 (5%)	92 (2%)	9	45

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	200	PRO	20
1	A	199	PRO	20
1	A	382	GLN	17
1	A	320	ARG	13
1	A	212	VAL	9

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/169 (100%)	157±3 (93±2%)	12±3 (7±2%)	18	66
All	All	3380/3380 (100%)	3132 (93%)	248 (7%)	18	66

5 of 52 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	198	ASP	20
1	A	286	SER	18
1	A	221	CYS	17
1	A	361	TRP	15
1	A	197	MET	15

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

MODRES-GEOMETRY INFOmissingINFO

5.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.4 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

5.5 Other polymers [i](#)

There are no such molecules in this entry.

5.6 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 77% for the well-defined parts and 77% for the entire structure.

6.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *data_A3G_PrC.str*

6.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2221
Number of shifts mapped to atoms	2221
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	18

6.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	175	-0.28 ± 0.19	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	165	0.68 ± 0.10	Should be applied
$^{13}\text{C}'$	160	-0.09 ± 0.10	None needed (< 0.5 ppm)
^{15}N	171	-0.23 ± 0.46	None needed (< 0.5 ppm)

6.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 77%, i.e. 1988 atoms were assigned a chemical shift out of a possible 2589. 21 out of 23 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	855/924 (93%)	349/368 (95%)	335/376 (89%)	171/180 (95%)
Sidechain	865/1186 (73%)	569/703 (81%)	285/420 (68%)	11/63 (17%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	202/309 (65%)	115/162 (71%)	81/124 (65%)	6/23 (26%)
Overall	1988/2589 (77%)	1099/1331 (83%)	701/983 (71%)	188/275 (68%)

6.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	206	PHE	CD2	122.14	137.34 – 125.84	-8.2
1	A	340	TYR	CE1	127.67	124.14 – 111.74	7.8
1	A	310	PHE	CD2	122.67	137.34 – 125.84	-7.8
1	A	206	PHE	CD1	122.14	137.63 – 125.43	-7.7
1	A	310	PHE	CD1	122.67	137.63 – 125.43	-7.3
1	A	340	TYR	CE2	127.67	124.68 – 111.18	7.2
1	A	247	PRO	CD	43.58	55.31 – 45.41	-6.9
1	A	314	ILE	CG2	26.07	24.63 – 10.43	6.0
1	A	219	TYR	CD1	125.40	139.11 – 126.41	-5.8
1	A	222	TYR	CE1	125.05	124.14 – 111.74	5.7
1	A	249	LYS	CD	22.29	34.86 – 23.06	-5.7
1	A	358	PHE	CD2	125.21	137.34 – 125.84	-5.5
1	A	325	LEU	CG	33.09	32.55 – 21.05	5.5
1	A	277	TYR	CD1	125.96	139.11 – 126.41	-5.4
1	A	222	TYR	CE2	125.05	124.68 – 111.18	5.3
1	A	358	PHE	CD1	125.21	137.63 – 125.43	-5.2
1	A	222	TYR	CD1	126.30	139.11 – 126.41	-5.1
1	A	355	GLY	H	11.65	11.63 – 5.03	5.0

6.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

