



wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 10:54 pm BST

PDB ID : 1K5G
Title : Crystal structure of Ran-GDP-AlFx-RanBP1-RanGAP complex
Authors : Seewald, M.J.; Koerner, C.; Wittinghofer, A.; Vetter, I.R.
Deposited on : 2001-10-10
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	<div> <div>2%</div> <div> <div>38%</div> <div>49%</div> <div>8%</div> <div>5%</div> </div> </div>
1	D	216	<div> <div>4%</div> <div> <div>34%</div> <div>52%</div> <div>9%</div> <div>5%</div> </div> </div>
1	G	216	<div> <div>35%</div> <div>51%</div> <div>8%</div> <div>5%</div> </div>
1	J	216	<div> <div>%</div> <div> <div>34%</div> <div>51%</div> <div>9%</div> <div>5%</div> </div> </div>
2	B	201	<div> <div>%</div> <div> <div>24%</div> <div>40%</div> <div>8%</div> <div>27%</div> </div> </div>
2	E	201	<div> <div>3%</div> <div> <div>22%</div> <div>43%</div> <div>7%</div> <div>27%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	201	 2% 25% 40% 7% 27%
2	K	201	 23% 43% 5% 27%
3	C	386	 45% 37% 6% 11%
3	F	386	 49% 34% 6% 11%
3	I	386	 47% 36% 6% 11%
3	L	386	 44% 39% 6% 11%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 22396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein RAN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	206	Total	C	N	O	S	0	0	0
			1652	1067	284	295	6			
1	D	206	Total	C	N	O	S	0	0	0
			1652	1067	284	295	6			
1	G	206	Total	C	N	O	S	0	0	0
			1652	1067	284	295	6			
1	J	206	Total	C	N	O	S	0	0	0
			1652	1067	284	295	6			

- Molecule 2 is a protein called Ran-specific GTPase-activating protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	0	0
			1216	769	214	226	7			
2	E	146	Total	C	N	O	S	0	0	0
			1216	769	214	226	7			
2	H	146	Total	C	N	O	S	0	0	0
			1216	769	214	226	7			
2	K	146	Total	C	N	O	S	0	0	0
			1216	769	214	226	7			

- Molecule 3 is a protein called Ran GTPase activating protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	344	Total	C	N	O	S	0	0	0
			2698	1699	469	522	8			
3	F	344	Total	C	N	O	S	0	0	0
			2698	1699	469	522	8			
3	I	344	Total	C	N	O	S	0	0	0
			2698	1699	469	522	8			
3	L	344	Total	C	N	O	S	0	0	0
			2698	1699	469	522	8			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	ALA	SER	SEE REMARK 999	UNP P41391
F	2	ALA	SER	SEE REMARK 999	UNP P41391
I	2	ALA	SER	SEE REMARK 999	UNP P41391
L	2	ALA	SER	SEE REMARK 999	UNP P41391

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	Total Mg 1 1	0	0
4	J	1	Total Mg 1 1	0	0
4	A	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0

- Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O P 28 10 5 11 2	0	0
5	D	1	Total C N O P 28 10 5 11 2	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	G	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
5	J	1	Total	C	N	O	P	0	0
			28	10	5	11	2		


- Molecule 6 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula: AlF_3).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	Al	F	0	0
			4	1	3		
6	D	1	Total	Al	F	0	0
			4	1	3		
6	G	1	Total	Al	F	0	0
			4	1	3		
6	J	1	Total	Al	F	0	0
			4	1	3		




• Molecule 1: GTP-binding nuclear protein RAN




• Molecule 2: Ran-specific GTPase-activating protein





• Molecule 2: Ran-specific GTPase-activating protein







• Molecule 2: Ran-specific GTPase-activating protein




• Molecule 2: Ran-specific GTPase-activating protein




• Molecule 3: Ran GTPase activating protein 1





• Molecule 3: Ran GTPase activating protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	99.93Å 102.57Å 118.85Å 71.67° 79.09° 67.81°	Depositor
Resolution (Å)	31.00 – 3.10 31.22 – 2.88	Depositor EDS
% Data completeness (in resolution range)	96.2 (31.00-3.10) 94.2 (31.22-2.88)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 2.90Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.250 , 0.269 0.224 , 0.263	Depositor DCC
R_{free} test set	8869 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	63.8	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 61.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	22396	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG, AF3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1693	0.67	0/2296
1	D	0.45	0/1693	0.68	0/2296
1	G	0.43	0/1693	0.67	0/2296
1	J	0.46	0/1693	0.69	0/2296
2	B	0.37	0/1242	0.63	0/1666
2	E	0.36	0/1242	0.62	0/1666
2	H	0.38	0/1242	0.62	0/1666
2	K	0.38	0/1242	0.63	0/1666
3	C	0.41	0/2737	0.68	0/3697
3	F	0.41	0/2737	0.68	0/3697
3	I	0.41	0/2737	0.67	0/3697
3	L	0.42	0/2737	0.69	0/3697
All	All	0.41	0/22688	0.67	0/30636

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1652	0	1660	129	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1652	0	1659	141	0
1	G	1652	0	1659	140	0
1	J	1652	0	1660	145	0
2	B	1216	0	1208	117	0
2	E	1216	0	1208	118	0
2	H	1216	0	1208	107	0
2	K	1216	0	1208	110	0
3	C	2698	0	2733	166	0
3	F	2698	0	2733	143	0
3	I	2698	0	2733	162	0
3	L	2698	0	2733	164	0
4	A	1	0	0	0	0
4	D	1	0	0	0	0
4	G	1	0	0	0	0
4	J	1	0	0	0	0
5	A	28	0	12	5	0
5	D	28	0	12	5	0
5	G	28	0	12	5	0
5	J	28	0	12	4	0
6	A	4	0	0	0	0
6	D	4	0	0	0	0
6	G	4	0	0	0	0
6	J	4	0	0	0	0
All	All	22396	0	22450	1526	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 1526 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:259:ARG:NH1	3:I:259:ARG:HB3	1.68	1.07
3:F:259:ARG:HB3	3:F:259:ARG:NH1	1.69	1.06
3:L:259:ARG:HB3	3:L:259:ARG:NH1	1.71	1.04
3:C:259:ARG:HB3	3:C:259:ARG:NH1	1.73	1.04
3:I:259:ARG:HH11	3:I:259:ARG:HB3	1.20	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/216 (94%)	156 (76%)	27 (13%)	21 (10%)	0	3
1	D	204/216 (94%)	153 (75%)	29 (14%)	22 (11%)	0	2
1	G	204/216 (94%)	156 (76%)	28 (14%)	20 (10%)	0	3
1	J	204/216 (94%)	154 (76%)	28 (14%)	22 (11%)	0	2
2	B	144/201 (72%)	118 (82%)	14 (10%)	12 (8%)	1	5
2	E	144/201 (72%)	117 (81%)	15 (10%)	12 (8%)	1	5
2	H	144/201 (72%)	119 (83%)	14 (10%)	11 (8%)	1	5
2	K	144/201 (72%)	117 (81%)	18 (12%)	9 (6%)	1	8
3	C	342/386 (89%)	275 (80%)	54 (16%)	13 (4%)	3	19
3	F	342/386 (89%)	280 (82%)	48 (14%)	14 (4%)	3	16
3	I	342/386 (89%)	278 (81%)	50 (15%)	14 (4%)	3	16
3	L	342/386 (89%)	278 (81%)	50 (15%)	14 (4%)	3	16
All	All	2760/3212 (86%)	2201 (80%)	375 (14%)	184 (7%)	1	7

5 of 184 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	106	ARG
1	A	135	SER
1	A	179	MET
1	A	186	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/185 (96%)	164 (92%)	14 (8%)	12	40
1	D	178/185 (96%)	161 (90%)	17 (10%)	8	31
1	G	178/185 (96%)	163 (92%)	15 (8%)	11	38
1	J	178/185 (96%)	161 (90%)	17 (10%)	8	31
2	B	131/176 (74%)	117 (89%)	14 (11%)	6	26
2	E	131/176 (74%)	118 (90%)	13 (10%)	8	29
2	H	131/176 (74%)	117 (89%)	14 (11%)	6	26
2	K	131/176 (74%)	119 (91%)	12 (9%)	9	33
3	C	295/334 (88%)	277 (94%)	18 (6%)	18	49
3	F	295/334 (88%)	277 (94%)	18 (6%)	18	49
3	I	295/334 (88%)	277 (94%)	18 (6%)	18	49
3	L	295/334 (88%)	277 (94%)	18 (6%)	18	49
All	All	2416/2780 (87%)	2228 (92%)	188 (8%)	12	40

5 of 188 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	173	ASN
1	G	155	TYR
3	L	43	THR
3	F	192	VAL
1	G	18	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 69 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	224	GLN
2	H	123	ASN
3	L	147	GLN
3	F	252	ASN
1	G	145	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	AF3	A	1252	5,4	0,3,3	0.00	-	-		
5	GDP	D	2250	4,6	24,30,30	1.84	6 (25%)	31,47,47	2.34	8 (25%)
6	AF3	J	4252	5,4	0,3,3	0.00	-	-		
5	GDP	G	3250	4,6	24,30,30	1.76	5 (20%)	31,47,47	2.41	10 (32%)
6	AF3	G	3252	5,4	0,3,3	0.00	-	-		
5	GDP	J	4250	4,6	24,30,30	1.80	4 (16%)	31,47,47	2.45	12 (38%)
6	AF3	D	2252	5,4	0,3,3	0.00	-	-		
5	GDP	A	1250	4,6	24,30,30	1.79	4 (16%)	31,47,47	2.37	10 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GDP	G	3250	4,6	-	3/12/32/32	0/3/3/3
5	GDP	D	2250	4,6	-	3/12/32/32	0/3/3/3
5	GDP	J	4250	4,6	-	3/12/32/32	0/3/3/3
5	GDP	A	1250	4,6	-	3/12/32/32	0/3/3/3

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1250	GDP	C6-N1	5.11	1.41	1.33
5	G	3250	GDP	C6-N1	5.02	1.41	1.33
5	J	4250	GDP	C6-N1	4.72	1.41	1.33
5	D	2250	GDP	C6-N1	4.55	1.41	1.33
5	D	2250	GDP	C6-C5	4.09	1.48	1.41

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	3250	GDP	C5-C6-N1	-8.51	111.79	123.43
5	D	2250	GDP	C5-C6-N1	-8.39	111.95	123.43
5	J	4250	GDP	C5-C6-N1	-8.30	112.08	123.43
5	A	1250	GDP	C5-C6-N1	-8.28	112.10	123.43
5	J	4250	GDP	C6-N1-C2	5.92	125.34	115.93

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	2250	GDP	PA-O3A-PB-O2B
5	G	3250	GDP	PA-O3A-PB-O2B
5	J	4250	GDP	PA-O3A-PB-O2B
5	A	1250	GDP	PA-O3A-PB-O2B
5	D	2250	GDP	PA-O3A-PB-O1B


There are no ring outliers.


4 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	2250	GDP	5	0
5	G	3250	GDP	5	0
5	J	4250	GDP	4	0
5	A	1250	GDP	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	206/216 (95%)	-0.48	5 (2%) 59 37	21, 57, 156, 186	0
1	D	206/216 (95%)	-0.50	9 (4%) 34 17	13, 46, 161, 199	0
1	G	206/216 (95%)	-0.55	1 (0%) 91 81	14, 50, 159, 198	0
1	J	206/216 (95%)	-0.62	2 (0%) 82 67	11, 47, 148, 199	0
2	B	146/201 (72%)	-0.27	3 (2%) 63 43	43, 88, 148, 200	0
2	E	146/201 (72%)	0.03	6 (4%) 37 18	43, 100, 162, 195	0
2	H	146/201 (72%)	-0.25	4 (2%) 54 31	32, 80, 158, 187	0
2	K	146/201 (72%)	-0.31	2 (1%) 75 56	44, 80, 145, 188	0
3	C	344/386 (89%)	-0.72	1 (0%) 94 88	18, 48, 107, 144	0
3	F	344/386 (89%)	-0.82	1 (0%) 94 88	11, 47, 93, 143	0
3	I	344/386 (89%)	-0.73	0 100 100	15, 50, 100, 148	0
3	L	344/386 (89%)	-0.78	0 100 100	8, 43, 97, 150	0
All	All	2784/3212 (86%)	-0.58	34 (1%) 79 61	8, 55, 139, 200	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	191	PRO	6.7
2	H	63	ASP	5.0
1	D	195	ALA	3.7
1	D	188	VAL	3.7
1	A	206	THR	3.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MG	D	2251	1/1	0.70	0.30	45,45,45,45	0
4	MG	G	3251	1/1	0.81	0.36	44,44,44,44	0
4	MG	J	4251	1/1	0.89	0.36	33,33,33,33	0
4	MG	A	1251	1/1	0.91	0.37	31,31,31,31	0
6	AF3	G	3252	4/4	0.95	0.20	57,57,57,57	0
5	GDP	A	1250	28/28	0.95	0.15	43,43,43,43	0
6	AF3	A	1252	4/4	0.95	0.26	57,57,57,57	0
5	GDP	G	3250	28/28	0.96	0.13	35,35,35,35	0
6	AF3	D	2252	4/4	0.96	0.27	57,57,57,57	0
5	GDP	J	4250	28/28	0.97	0.12	29,29,29,29	0
5	GDP	D	2250	28/28	0.97	0.12	31,31,31,31	0
6	AF3	J	4252	4/4	0.98	0.25	57,57,57,57	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GDP A 1250:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GDP G 3250:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP J 4250:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GDP D 2250:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.