



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 15, 2020 – 02:03 am BST

PDB ID : 3K5H  
Title : Crystal structure of carboxyaminoimidazole ribonucleotide synthase from asperigillus clavatus complexed with ATP  
Authors : Thoden, J.B.; Holden, H.M.; Paritala, H.; Firestine, S.M.  
Deposited on : 2009-10-07  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	<div> <div>50%</div> <div>37%</div> <div>7%</div> <div>5%</div> </div>
1	B	403	<div>2%</div> <div>46%</div> <div>43%</div> <div>5%</div> <div>6%</div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoribosyl-aminoimidazole carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	4	0
			2977	1869	525	566	17			
1	B	380	Total	C	N	O	S	0	5	0
			2969	1865	519	570	15			
1	C	382	Total	C	N	O	S	0	2	0
			2975	1868	527	564	16			
1	D	380	Total	C	N	O	S	0	1	0
			2948	1854	518	561	15			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP A1CII2
A	-18	GLY	-	EXPRESSION TAG	UNP A1CII2
A	-17	SER	-	EXPRESSION TAG	UNP A1CII2
A	-16	SER	-	EXPRESSION TAG	UNP A1CII2
A	-15	HIS	-	EXPRESSION TAG	UNP A1CII2
A	-14	HIS	-	EXPRESSION TAG	UNP A1CII2
A	-13	HIS	-	EXPRESSION TAG	UNP A1CII2
A	-12	HIS	-	EXPRESSION TAG	UNP A1CII2
A	-11	HIS	-	EXPRESSION TAG	UNP A1CII2
A	-10	HIS	-	EXPRESSION TAG	UNP A1CII2
A	-9	SER	-	EXPRESSION TAG	UNP A1CII2
A	-8	SER	-	EXPRESSION TAG	UNP A1CII2
A	-7	GLU	-	EXPRESSION TAG	UNP A1CII2
A	-6	ASN	-	EXPRESSION TAG	UNP A1CII2
A	-5	LEU	-	EXPRESSION TAG	UNP A1CII2
A	-4	TYR	-	EXPRESSION TAG	UNP A1CII2
A	-3	PHE	-	EXPRESSION TAG	UNP A1CII2
A	-2	GLN	-	EXPRESSION TAG	UNP A1CII2
A	-1	GLY	-	EXPRESSION TAG	UNP A1CII2
A	0	HIS	-	EXPRESSION TAG	UNP A1CII2
B	-19	MET	-	EXPRESSION TAG	UNP A1CII2

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	GLY	-	EXPRESSION TAG	UNP A1CII2
B	-17	SER	-	EXPRESSION TAG	UNP A1CII2
B	-16	SER	-	EXPRESSION TAG	UNP A1CII2
B	-15	HIS	-	EXPRESSION TAG	UNP A1CII2
B	-14	HIS	-	EXPRESSION TAG	UNP A1CII2
B	-13	HIS	-	EXPRESSION TAG	UNP A1CII2
B	-12	HIS	-	EXPRESSION TAG	UNP A1CII2
B	-11	HIS	-	EXPRESSION TAG	UNP A1CII2
B	-10	HIS	-	EXPRESSION TAG	UNP A1CII2
B	-9	SER	-	EXPRESSION TAG	UNP A1CII2
B	-8	SER	-	EXPRESSION TAG	UNP A1CII2
B	-7	GLU	-	EXPRESSION TAG	UNP A1CII2
B	-6	ASN	-	EXPRESSION TAG	UNP A1CII2
B	-5	LEU	-	EXPRESSION TAG	UNP A1CII2
B	-4	TYR	-	EXPRESSION TAG	UNP A1CII2
B	-3	PHE	-	EXPRESSION TAG	UNP A1CII2
B	-2	GLN	-	EXPRESSION TAG	UNP A1CII2
B	-1	GLY	-	EXPRESSION TAG	UNP A1CII2
B	0	HIS	-	EXPRESSION TAG	UNP A1CII2
C	-19	MET	-	EXPRESSION TAG	UNP A1CII2
C	-18	GLY	-	EXPRESSION TAG	UNP A1CII2
C	-17	SER	-	EXPRESSION TAG	UNP A1CII2
C	-16	SER	-	EXPRESSION TAG	UNP A1CII2
C	-15	HIS	-	EXPRESSION TAG	UNP A1CII2
C	-14	HIS	-	EXPRESSION TAG	UNP A1CII2
C	-13	HIS	-	EXPRESSION TAG	UNP A1CII2
C	-12	HIS	-	EXPRESSION TAG	UNP A1CII2
C	-11	HIS	-	EXPRESSION TAG	UNP A1CII2
C	-10	HIS	-	EXPRESSION TAG	UNP A1CII2
C	-9	SER	-	EXPRESSION TAG	UNP A1CII2
C	-8	SER	-	EXPRESSION TAG	UNP A1CII2
C	-7	GLU	-	EXPRESSION TAG	UNP A1CII2
C	-6	ASN	-	EXPRESSION TAG	UNP A1CII2
C	-5	LEU	-	EXPRESSION TAG	UNP A1CII2
C	-4	TYR	-	EXPRESSION TAG	UNP A1CII2
C	-3	PHE	-	EXPRESSION TAG	UNP A1CII2
C	-2	GLN	-	EXPRESSION TAG	UNP A1CII2
C	-1	GLY	-	EXPRESSION TAG	UNP A1CII2
C	0	HIS	-	EXPRESSION TAG	UNP A1CII2
D	-19	MET	-	EXPRESSION TAG	UNP A1CII2
D	-18	GLY	-	EXPRESSION TAG	UNP A1CII2
D	-17	SER	-	EXPRESSION TAG	UNP A1CII2

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	EXPRESSION TAG	UNP A1CII2
D	-15	HIS	-	EXPRESSION TAG	UNP A1CII2
D	-14	HIS	-	EXPRESSION TAG	UNP A1CII2
D	-13	HIS	-	EXPRESSION TAG	UNP A1CII2
D	-12	HIS	-	EXPRESSION TAG	UNP A1CII2
D	-11	HIS	-	EXPRESSION TAG	UNP A1CII2
D	-10	HIS	-	EXPRESSION TAG	UNP A1CII2
D	-9	SER	-	EXPRESSION TAG	UNP A1CII2
D	-8	SER	-	EXPRESSION TAG	UNP A1CII2
D	-7	GLU	-	EXPRESSION TAG	UNP A1CII2
D	-6	ASN	-	EXPRESSION TAG	UNP A1CII2
D	-5	LEU	-	EXPRESSION TAG	UNP A1CII2
D	-4	TYR	-	EXPRESSION TAG	UNP A1CII2
D	-3	PHE	-	EXPRESSION TAG	UNP A1CII2
D	-2	GLN	-	EXPRESSION TAG	UNP A1CII2
D	-1	GLY	-	EXPRESSION TAG	UNP A1CII2
D	0	HIS	-	EXPRESSION TAG	UNP A1CII2

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	A	2	Total	Mg	0	0
			2	2		
3	D	2	Total	Mg	0	0
			2	2		
3	C	2	Total	Mg	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	227	Total	O	0	0
			227	227		
4	B	247	Total	O	0	0
			247	247		
4	C	179	Total	O	0	0
			179	179		
4	D	196	Total	O	0	0
			196	196		


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Phosphoribosyl-aminoimidazole carboxylase




#### • Molecule 1: Phosphoribosyl-aminoimidazole carboxylase








• Molecule 1: Phosphoribosyl-aminoimidazole carboxylase



• Molecule 1: Phosphoribosyl-aminoimidazole carboxylase







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.90 Å 134.40 Å 99.60 Å 90.00° 107.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 40.62 – 2.10	Depositor EDS
% Data completeness (in resolution range)	93.6 (30.00-2.10) 93.5 (40.62-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.63 (at 2.10 Å)	Xtriage
Refinement program	TNT	Depositor
R, $R_{free}$	0.196 , 0.261 0.195 , 0.193	Depositor DCC
$R_{free}$ test set	10356 reflections (9.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.5	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 104.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12850	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.79	1/3046 (0.0%)	1.49	29/4124 (0.7%)
1	B	0.79	1/3042 (0.0%)	1.53	31/4120 (0.8%)
1	C	0.77	1/3036 (0.0%)	1.46	20/4110 (0.5%)
1	D	0.73	1/3005 (0.0%)	1.55	42/4071 (1.0%)
All	All	0.77	4/12129 (0.0%)	1.51	122/16425 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
1	D	1	0
All	All	1	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	67	CYS	CB-SG	-6.34	1.71	1.82
1	A	189	GLU	CD-OE1	-5.96	1.19	1.25
1	C	137	GLU	CB-CG	-5.77	1.41	1.52
1	D	188	MET	SD-CE	-5.67	1.46	1.77

The worst 5 of 122 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	143	LEU	CB-CG-CD1	-11.52	91.42	111.00
1	C	144	MET	CG-SD-CE	-10.85	82.85	100.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57	ARG	NE-CZ-NH2	-10.76	114.92	120.30
1	A	127	ASN	N-CA-CB	-10.72	91.30	110.60
1	A	376	ASP	CB-CG-OD1	-9.95	109.34	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	280	ILE	CB

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	TYR	Sidechain
1	C	114	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2977	0	2988	135	0
1	B	2969	0	2965	161	0
1	C	2975	0	2991	162	0
1	D	2948	0	2960	205	0
2	A	31	0	12	1	0
2	B	31	0	12	2	0
2	C	31	0	12	1	0
2	D	31	0	12	2	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	227	0	0	3	0
4	B	247	0	0	11	0
4	C	179	0	0	9	0
4	D	196	0	0	12	0
All	All	12850	0	11952	659	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 659 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:LEU:HD13	1:A:173:LEU:HD12	1.25	1.15
1:B:174:LYS:HD2	1:B:174:LYS:H	1.11	1.07
1:C:41:LYS:HE2	1:C:49:HIS:HB3	1.36	1.05
1:D:132:LEU:HD23	1:D:166:ILE:HG23	1.40	1.03
1:D:135:VAL:HA	1:D:138:GLN:HE21	1.19	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/403 (95%)	363 (94%)	20 (5%)	1 (0%)	41	41
1	B	383/403 (95%)	368 (96%)	15 (4%)	0	100	100
1	C	382/403 (95%)	361 (94%)	19 (5%)	2 (0%)	29	26
1	D	379/403 (94%)	351 (93%)	27 (7%)	1 (0%)	41	41
All	All	1528/1612 (95%)	1443 (94%)	81 (5%)	4 (0%)	41	41

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	259	GLU
1	C	227	SER
1	D	324	THR
1	A	57	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/336 (96%)	282 (88%)	40 (12%)	4	2
1	B	321/336 (96%)	293 (91%)	28 (9%)	10	7
1	C	320/336 (95%)	280 (88%)	40 (12%)	4	2
1	D	317/336 (94%)	281 (89%)	36 (11%)	5	3
All	All	1280/1344 (95%)	1136 (89%)	144 (11%)	6	3

5 of 144 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	344	SER
1	C	139	LEU
1	D	187	LYS
1	B	381	ILE
1	C	87	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	102	GLN
1	C	106	ASN
1	D	127	ASN
1	B	232	GLN
1	D	106	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	ATP	B	400	3	26,33,33	1.04	2 (7%)	31,52,52	2.27	10 (32%)
2	ATP	D	400	3	26,33,33	0.94	1 (3%)	31,52,52	2.95	7 (22%)
2	ATP	A	400	3	26,33,33	0.89	1 (3%)	31,52,52	2.75	11 (35%)
2	ATP	C	400	3	26,33,33	0.78	0	31,52,52	1.79	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	B	400	3	-	3/18/38/38	0/3/3/3
2	ATP	D	400	3	-	5/18/38/38	0/3/3/3
2	ATP	A	400	3	-	6/18/38/38	0/3/3/3
2	ATP	C	400	3	-	3/18/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	400	ATP	C2-N1	2.62	1.38	1.33
2	B	400	ATP	C6-N6	-2.50	1.24	1.34
2	B	400	ATP	C2-N1	2.18	1.38	1.33
2	A	400	ATP	C2-N1	2.11	1.37	1.33

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	400	ATP	C2-N1-C6	9.33	134.72	118.75
2	A	400	ATP	C5-C6-N6	8.77	133.69	120.35
2	B	400	ATP	C5-C6-N6	8.51	133.28	120.35
2	D	400	ATP	C5-C6-N1	-8.12	101.94	120.35
2	D	400	ATP	C5-C6-N6	6.58	130.35	120.35

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	400	ATP	PB-O3B-PG-O3G
2	B	400	ATP	PB-O3B-PG-O3G
2	D	400	ATP	PB-O3B-PG-O2G
2	D	400	ATP	PB-O3B-PG-O3G
2	A	400	ATP	PB-O3B-PG-O2G


There are no ring outliers.


4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	400	ATP	2	0
2	D	400	ATP	2	0
2	A	400	ATP	1	0
2	C	400	ATP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	382/403 (94%)	-0.14	2 (0%) 91 92	7, 25, 62, 81	0
1	B	380/403 (94%)	-0.10	10 (2%) 56 61	6, 24, 64, 86	0
1	C	382/403 (94%)	-0.04	5 (1%) 77 80	12, 31, 69, 93	0
1	D	380/403 (94%)	0.03	8 (2%) 63 68	11, 29, 70, 93	0
All	All	1524/1612 (94%)	-0.06	25 (1%) 72 75	6, 28, 67, 93	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	381	ILE	6.1
1	D	381	ILE	5.7
1	B	323	ASP	3.9
1	B	321	ALA	3.1
1	D	326	LEU	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	B	401	1/1	0.84	0.09	22,22,22,22	0
3	MG	C	401	1/1	0.85	0.07	40,40,40,40	0
3	MG	D	401	1/1	0.87	0.08	33,33,33,33	0
3	MG	B	402	1/1	0.92	0.08	25,25,25,25	0
2	ATP	C	400	31/31	0.96	0.10	12,30,82,100	0
3	MG	D	402	1/1	0.96	0.05	24,24,24,24	0
2	ATP	B	400	31/31	0.96	0.10	7,28,44,63	0
3	MG	C	402	1/1	0.96	0.09	22,22,22,22	0
2	ATP	D	400	31/31	0.97	0.09	12,26,56,100	0
2	ATP	A	400	31/31	0.97	0.08	9,23,33,75	0
3	MG	A	401	1/1	0.97	0.04	14,14,14,14	0
3	MG	A	402	1/1	0.97	0.06	24,24,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around ATP B 400:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ATP D 400:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.