



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 8, 2020 – 02:17 AM EST

PDB ID : 7K5E  
Title : 1.75 Å resolution structure of WT BfrB from *Pseudomonas aeruginosa* in complex with a protein-protein interaction inhibitor JAG-5-7  
Authors : Lovell, S.; Battaile, K.P.; Soldano, A.; Punchi-Hewage, A.; Meraz, K.; Annor-Gyamfi, J.K.; Yao, H.; Bunce, R.A.; Rivera, M.  
Deposited on : 2020-09-16  
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.15.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.15.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*






The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	 97% ..
1	B	158	 94% 5% .
1	C	158	 94% ..
1	D	158	 94% ..
1	E	158	 96% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	158	 95% . .
1	G	158	 92% 7% .
1	H	158	 95% . .
1	I	158	 98% ..
1	J	158	 95% . .
1	K	158	 95% . .
1	L	158	 91% 8% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	VXM	C	202	-	X	-	-
4	VXM	D	203	-	X	-	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 17271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	2	0
			1278	808	216	247	7			
1	B	156	Total	C	N	O	S	0	2	0
			1277	807	216	247	7			
1	C	156	Total	C	N	O	S	0	1	0
			1271	803	217	244	7			
1	D	156	Total	C	N	O	S	0	1	0
			1277	807	217	246	7			
1	E	156	Total	C	N	O	S	0	2	0
			1271	804	215	245	7			
1	F	156	Total	C	N	O	S	0	2	0
			1277	807	216	247	7			
1	G	156	Total	C	N	O	S	0	3	0
			1289	814	221	247	7			
1	H	156	Total	C	N	O	S	0	2	0
			1279	808	218	246	7			
1	I	156	Total	C	N	O	S	0	2	0
			1282	810	219	246	7			
1	J	156	Total	C	N	O	S	0	2	0
			1286	813	220	246	7			
1	K	156	Total	C	N	O	S	0	2	0
			1289	815	221	246	7			
1	L	156	Total	C	N	O	S	0	2	0
			1283	810	218	248	7			

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	K	0	0
			1	1		
2	A	1	Total	K	0	0
			1	1		


*Continued on next page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total K 1 1	0	0

- # HEM

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	H	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	L	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- 
- WORLD WIDE  
PDB  
PROTEIN DATA BANK



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	F	N	O	0	1
			42	30	2	4	6		
4	C	1	Total	C	N	O		0	0
			14	10	2	2			
4	D	1	Total	C	N	O		0	0
			14	10	2	2			
4	E	1	Total	C	N	O		0	0
			13	9	2	2			
4	G	1	Total	C	F	N	O	0	0
			21	15	1	2	3		
4	H	1	Total	C	F	N	O	0	1
			42	30	2	4	6		
4	I	1	Total	C	N	O		0	0
			14	10	2	2			
4	J	1	Total	C	F	N	O	0	0
			21	15	1	2	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	121	Total	O	0	0
			121	121		
5	B	100	Total	O	0	0
			100	100		
5	C	102	Total	O	0	0
			102	102		
5	D	124	Total	O	0	0
			124	124		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	123	Total 123	O 123	0	0
5	F	110	Total 110	O 110	0	0
5	G	132	Total 132	O 132	0	0
5	H	134	Total 134	O 134	0	0
5	I	132	Total 132	O 132	0	0
5	J	156	Total 156	O 156	0	0
5	K	108	Total 108	O 108	0	0
5	L	85	Total 85	O 85	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ferroxidase

Chain A:  97% ..



- Molecule 1: Ferroxidase

Chain B:  94% 5% .



- Molecule 1: Ferroxidase

Chain C:  94% . .



- Molecule 1: Ferroxidase

Chain D:  94% . .



- Molecule 1: Ferroxidase

Chain E:  96% ..



- Molecule 1: Ferroxidase

Chain F:  95% . .





- Molecule 1: Ferroxidase

Chain G: 92% 7% .



- Molecule 1: Ferroxidase

Chain H: 95% . .



- Molecule 1: Ferroxidase

Chain I: 98% . .



- Molecule 1: Ferroxidase

Chain J: 95% . .




- Molecule 1: Ferroxidase

Chain K: 95% . .



- Molecule 1: Ferroxidase

Chain L: 91% 8% .



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	129.85Å 194.28Å 203.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.36 – 1.75 48.57 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.36-1.75) 100.0 (48.57-1.75)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 1.75Å)	Xtriage
Refinement program	PHENIX 1.17rc2_3615	Depositor
R, $R_{free}$	0.182 , 0.213 0.191 , 0.218	Depositor DCC
$R_{free}$ test set	12616 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.9	Xtriage
Anisotropy	0.756	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	17271	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VXM, HEM, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/1302	0.53	0/1756
1	B	0.42	0/1301	0.53	0/1755
1	C	0.46	0/1292	0.57	0/1743
1	D	0.41	0/1298	0.51	0/1750
1	E	0.46	0/1298	0.56	0/1752
1	F	0.45	0/1301	0.53	0/1755
1	G	0.43	0/1314	0.53	0/1772
1	H	0.42	0/1303	0.53	0/1757
1	I	0.43	0/1306	0.57	0/1760
1	J	0.46	0/1310	0.54	0/1764
1	K	0.44	0/1314	0.58	0/1770
1	L	0.38	0/1307	0.53	0/1762
All	All	0.43	0/15646	0.54	0/21096

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1278	0	1236	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1277	0	1234	6	0
1	C	1271	0	1230	4	0
1	D	1277	0	1238	5	0
1	E	1271	0	1228	3	0
1	F	1277	0	1234	4	0
1	G	1289	0	1249	6	0
1	H	1279	0	1243	5	0
1	I	1282	0	1252	1	0
1	J	1286	0	1263	4	0
1	K	1289	0	1261	4	0
1	L	1283	0	1247	9	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
3	A	43	0	30	4	0
3	B	43	0	30	5	0
3	C	43	0	30	3	0
3	D	43	0	30	6	0
3	F	43	0	30	4	0
3	H	43	0	30	5	0
3	L	43	0	30	5	0
4	B	42	0	0	2	0
4	C	14	0	0	1	0
4	D	14	0	0	1	0
4	E	13	0	0	1	0
4	G	21	0	0	1	0
4	H	42	0	0	2	0
4	I	14	0	0	1	0
4	J	21	0	0	1	0
5	A	121	0	0	0	0
5	B	100	0	0	1	0
5	C	102	0	0	0	0
5	D	124	0	0	1	0
5	E	123	0	0	1	0
5	F	110	0	0	1	0
5	G	132	0	0	1	0
5	H	134	0	0	1	0
5	I	132	0	0	0	0
5	J	156	0	0	0	0
5	K	108	0	0	1	0
5	L	85	0	0	0	0
All	All	17271	0	15125	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:201:HEM:HBC2	3:H:201:HEM:HMC2	1.60	0.83
3:B:202:HEM:HMC2	3:B:202:HEM:HBC2	1.71	0.72
3:H:201:HEM:CMC	3:H:201:HEM:HBC2	2.22	0.69
1:K:20:ILE:HD11	1:K:75:GLY:HA3	1.79	0.65
1:G:69:PRO:O	4:G:201:VXM:N03	2.30	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/158 (99%)	155 (99%)	1 (1%)	0	100	100
1	B	156/158 (99%)	155 (99%)	1 (1%)	0	100	100
1	C	155/158 (98%)	155 (100%)	0	0	100	100
1	D	155/158 (98%)	154 (99%)	1 (1%)	0	100	100
1	E	156/158 (99%)	155 (99%)	1 (1%)	0	100	100
1	F	156/158 (99%)	156 (100%)	0	0	100	100
1	G	157/158 (99%)	157 (100%)	0	0	100	100
1	H	156/158 (99%)	155 (99%)	1 (1%)	0	100	100
1	I	156/158 (99%)	156 (100%)	0	0	100	100
1	J	156/158 (99%)	156 (100%)	0	0	100	100
1	K	156/158 (99%)	155 (99%)	1 (1%)	0	100	100
1	L	156/158 (99%)	154 (99%)	2 (1%)	0	100	100
All	All	1871/1896 (99%)	1863 (100%)	8 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/144 (95%)	137 (100%)	0	100	100
1	B	137/144 (95%)	136 (99%)	1 (1%)	84	75
1	C	136/144 (94%)	135 (99%)	1 (1%)	84	75
1	D	137/144 (95%)	136 (99%)	1 (1%)	84	75
1	E	136/144 (94%)	135 (99%)	1 (1%)	84	75
1	F	137/144 (95%)	137 (100%)	0	100	100
1	G	139/144 (96%)	138 (99%)	1 (1%)	84	75
1	H	138/144 (96%)	137 (99%)	1 (1%)	84	75
1	I	139/144 (96%)	139 (100%)	0	100	100
1	J	140/144 (97%)	140 (100%)	0	100	100
1	K	140/144 (97%)	140 (100%)	0	100	100
1	L	139/144 (96%)	138 (99%)	1 (1%)	84	75
All	All	1655/1728 (96%)	1648 (100%)	7 (0%)	91	87

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	121	LYS
1	L	84	GLN
1	G	121	LYS
1	C	121	LYS
1	H	121	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	142	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	L	130	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 20 ligands modelled in this entry, 3 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	HEM	F	201	1	27,50,50	1.91	5 (18%)	17,82,82	1.93	5 (29%)
4	VXM	B	203[A]	-	23,23,23	2.44	11 (47%)	33,33,33	1.68	5 (15%)
3	HEM	C	201	1	27,50,50	1.84	4 (14%)	17,82,82	1.92	4 (23%)
4	VXM	I	201	-	15,15,23	3.18	9 (60%)	21,21,33	2.05	7 (33%)
4	VXM	H	202[B]	-	23,23,23	2.41	13 (56%)	33,33,33	1.67	7 (21%)
4	VXM	C	202	-	15,15,23	3.42	13 (86%)	21,21,33	2.05	7 (33%)
4	VXM	J	201	-	23,23,23	2.29	11 (47%)	33,33,33	1.70	6 (18%)
4	VXM	E	201	-	14,14,23	3.90	12 (85%)	19,20,33	2.65	5 (26%)
4	VXM	D	203	-	15,15,23	3.21	12 (80%)	21,21,33	2.66	8 (38%)
4	VXM	G	201	-	23,23,23	2.21	12 (52%)	33,33,33	1.75	7 (21%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	VXM	H	202[A]	-	23,23,23	2.50	14 (60%)	33,33,33	1.65	6 (18%)
3	HEM	L	201	1	27,50,50	1.85	4 (14%)	17,82,82	1.58	4 (23%)
3	HEM	H	201	1	27,50,50	1.73	5 (18%)	17,82,82	2.04	6 (35%)
3	HEM	D	202	1	27,50,50	1.72	4 (14%)	17,82,82	1.71	4 (23%)
3	HEM	A	202	1	27,50,50	1.86	5 (18%)	17,82,82	1.88	5 (29%)
3	HEM	B	202	1	27,50,50	1.86	6 (22%)	17,82,82	1.96	6 (35%)
4	VXM	B	203[B]	-	23,23,23	2.42	11 (47%)	33,33,33	1.69	5 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	F	201	1	-	0/6/54/54	-
4	VXM	B	203[A]	-	-	0/5/17/17	0/3/3/3
3	HEM	C	201	1	-	0/6/54/54	-
4	VXM	I	201	-	-	3/3/15/17	0/2/2/3
4	VXM	H	202[B]	-	-	0/5/17/17	0/3/3/3
4	VXM	C	202	-	-	3/3/15/17	0/2/2/3
4	VXM	J	201	-	-	0/5/17/17	0/3/3/3
4	VXM	E	201	-	-	2/2/14/17	0/2/2/3
4	VXM	D	203	-	-	3/3/15/17	0/2/2/3
4	VXM	G	201	-	-	2/5/17/17	0/3/3/3
4	VXM	H	202[A]	-	-	0/5/17/17	0/3/3/3
3	HEM	L	201	1	-	0/6/54/54	-
3	HEM	H	201	1	-	0/6/54/54	-
3	HEM	D	202	1	-	0/6/54/54	-
3	HEM	A	202	1	-	0/6/54/54	-
3	HEM	B	202	1	-	0/6/54/54	-
4	VXM	B	203[B]	-	-	0/5/17/17	0/3/3/3

The worst 5 of 151 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	202	VXM	C20-C02	7.91	1.60	1.48
4	E	201	VXM	C06-N07	7.90	1.52	1.37
4	I	201	VXM	C20-C02	7.62	1.60	1.48
4	E	201	VXM	C20-C02	5.54	1.57	1.48
4	E	201	VXM	C08-N07	5.52	1.54	1.45



The worst 5 of 97 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	201	VXM	C04-N03-C02	-6.63	106.87	112.52
4	D	203	VXM	C04-N03-C02	-6.52	106.96	112.52
4	B	203[B]	VXM	C04-N03-C02	-6.36	107.10	112.52
4	B	203[A]	VXM	C04-N03-C02	-6.21	107.22	112.52
4	H	202[A]	VXM	C04-N03-C02	-5.19	108.09	112.52

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	201	VXM	C17-C06-N07-C08
4	E	201	VXM	C05-C06-N07-C08
4	D	203	VXM	C17-C06-N07-C08
4	D	203	VXM	C05-C06-N07-C08
4	C	202	VXM	C17-C06-N07-C08


There are no ring outliers.


17 monomers are involved in 42 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	201	HEM	4	0
4	B	203[A]	VXM	1	0
3	C	201	HEM	3	0
4	I	201	VXM	1	0
4	H	202[B]	VXM	1	0
4	C	202	VXM	1	0
4	J	201	VXM	1	0
4	E	201	VXM	1	0
4	D	203	VXM	1	0
4	G	201	VXM	1	0
4	H	202[A]	VXM	1	0
3	L	201	HEM	5	0
3	H	201	HEM	5	0
3	D	202	HEM	6	0
3	A	202	HEM	4	0
3	B	202	HEM	5	0
4	B	203[B]	VXM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## Ligand VXM I 201





## Ligand VXM H 202 (B)
































## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	156/158 (98%)	-0.19	0 100 100	19, 27, 37, 48	0
1	B	156/158 (98%)	0.09	0 100 100	22, 29, 39, 51	0
1	C	156/158 (98%)	-0.18	0 100 100	20, 26, 37, 55	0
1	D	156/158 (98%)	-0.02	0 100 100	21, 26, 37, 50	0
1	E	156/158 (98%)	-0.11	0 100 100	19, 25, 34, 50	0
1	F	156/158 (98%)	-0.19	0 100 100	21, 25, 35, 52	0
1	G	156/158 (98%)	-0.24	0 100 100	17, 25, 34, 48	0
1	H	156/158 (98%)	-0.29	0 100 100	18, 23, 33, 48	0
1	I	156/158 (98%)	-0.27	0 100 100	18, 23, 33, 54	0
1	J	156/158 (98%)	-0.29	0 100 100	16, 20, 28, 51	0
1	K	156/158 (98%)	-0.22	0 100 100	18, 24, 37, 47	0
1	L	156/158 (98%)	-0.00	0 100 100	26, 32, 42, 53	0
All	All	1872/1896 (98%)	-0.16	0 100 100	16, 26, 37, 55	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	VXM	C	202	14/21	0.65	0.29	32,40,49,50	0
4	VXM	I	201	14/21	0.70	0.25	33,44,56,56	0
4	VXM	G	201	21/21	0.75	0.23	33,46,58,61	0
4	VXM	J	201	21/21	0.78	0.28	28,44,63,65	0
4	VXM	H	202[B]	21/21	0.78	0.29	30,39,49,58	21
4	VXM	H	202[A]	21/21	0.78	0.29	30,38,49,50	21
4	VXM	B	203[A]	21/21	0.83	0.21	29,36,49,51	21
4	VXM	B	203[B]	21/21	0.83	0.21	29,36,49,50	21
4	VXM	E	201	13/21	0.89	0.19	35,42,46,53	0
4	VXM	D	203	14/21	0.90	0.15	32,40,48,60	0
3	HEM	B	202	43/43	0.94	0.12	21,27,42,46	0
3	HEM	D	202	43/43	0.94	0.12	23,26,35,38	43
3	HEM	L	201	43/43	0.95	0.11	23,29,36,39	43
3	HEM	A	202	43/43	0.95	0.11	20,25,39,47	0
3	HEM	C	201	43/43	0.96	0.10	19,27,40,44	0
2	K	B	201	1/1	0.96	0.11	24,24,24,24	0
3	HEM	F	201	43/43	0.96	0.10	21,25,39,45	0
3	HEM	H	201	43/43	0.96	0.10	17,22,37,39	0
2	K	A	201	1/1	0.99	0.11	17,17,17,17	0
2	K	D	201	1/1	0.99	0.09	18,18,18,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around VXM C 202:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around VXM I 201:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around VXM G 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around VXM J 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around VXM H 202 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around VXM H 202 (A):**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around VXM B 203 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around VXM B 203 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around VXM E 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around VXM D 203:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM B 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM D 202:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around HEM L 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM A 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM C 201:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM F 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.