



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 01:21 PM BST

PDB ID : 5K8D  
Title : Crystal structure of rFVIII<sup>Fc</sup>  
Authors : Leksa, N.; Quan, C.  
Deposited on : 2016-05-29  
Resolution : 4.19 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 4.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1005 (4.62-3.78)
Clashscore	141614	1044 (4.60-3.80)
Ramachandran outliers	138981	1000 (4.60-3.80)
Sidechain outliers	138945	1007 (4.62-3.78)
RSRZ outliers	127900	1063 (4.70-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	740	<div> <div>21%</div> <div>45%</div> <div>16%</div> <div>•</div> <div>17%</div> </div>
2	B	865	<div> <div>20%</div> <div>36%</div> <div>15%</div> <div>•</div> <div>28%</div> </div>
3	C	4	<div> <div>25%</div> <div>75%</div> </div>
4	D	2	<div> <div>100%</div> </div>
5	E	7	<div> <div>14%</div> <div>43%</div> <div>43%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BMA	C	3	-	-	X	-
4	NAG	D	1	-	-	-	X
4	NAG	D	2	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor VIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	611	Total	C	N	O	S	0	0	0
			4937	3190	824	898	25			

- Molecule 2 is a protein called Coagulation factor VIII,Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	619	Total	C	N	O	S	0	0	0
			5035	3234	862	908	31			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	2332	TYR	-	linker	UNP P00451

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	4	Total	C	N	O		0	0	0
			50	28	2	20				

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-beta-D-mannopyranos e-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	Ca	0	0
			3	3		


- Molecule 7 is COPPER (II) ION (three-letter code: CU) (formula: Cu).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Cu	0	0
			1	1		
7	A	1	Total	Cu	0	0
			1	1		

### 3 Residue-property plots



These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Coagulation factor VIII





• Molecule 2: Coagulation factor VIII,Ig gamma-1 chain C region



- Molecule 3: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  25% 75%

MAN1  
MAN2  
MAN3  
MAN4

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAN1  
MAN2

- Molecule 5: alpha-D-mannopyranose-(1-2)-beta-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  14% 43% 43%

MAN1  
MAN2  
MAN3  
MAN4  
MAN5  
MAN6  
MAN7



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	136.29 Å   136.29 Å   365.10 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	37.60 – 4.19 37.60 – 4.19	Depositor EDS
% Data completeness (in resolution range)	98.8 (37.60-4.19) 99.0 (37.60-4.19)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 4.13 Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.298   ,   0.362 0.298   ,   0.361	Depositor DCC
$R_{free}$ test set	1309 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	126.3	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.19   ,   131.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	10138	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	205.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, BMA, NAG, CU, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.86	15/5077 (0.3%)	0.90	4/6890 (0.1%)
2	B	0.80	21/5178 (0.4%)	0.86	1/7011 (0.0%)
All	All	0.83	36/10255 (0.4%)	0.88	5/13901 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	2
All	All	0	4

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	434	GLU	CD-OE1	14.90	1.42	1.25
1	A	389	GLU	CD-OE2	14.10	1.41	1.25
1	A	697	PHE	CG-CD2	13.56	1.59	1.38
1	A	697	PHE	CG-CD1	11.83	1.56	1.38
1	A	697	PHE	CE1-CZ	10.48	1.57	1.37

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	491	LEU	C-N-CD	-6.36	106.62	120.60
1	A	179	CYS	CA-CB-SG	-5.99	103.21	114.00
2	B	2330	ASP	CB-CG-OD2	-5.94	112.96	118.30
1	A	186	LYS	CD-CE-NZ	5.28	123.84	111.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	250	ARG	NE-CZ-NH1	5.12	122.86	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	430	ALA	Peptide
1	A	693	HIS	Peptide
2	B	1790	ILE	Peptide
2	B	1828	ASP	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4937	0	4820	803	1
2	B	5035	0	4902	791	0
3	C	50	0	43	8	0
4	D	28	0	25	0	0
5	E	83	0	70	4	0
6	A	3	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
All	All	10138	0	9860	1508	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 75.

The worst 5 of 1508 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:2046:TRP:CZ3	2:B:2059:ILE:HA	1.37	1.59
1:A:287:GLU:HB3	1:A:673:PHE:CE2	1.51	1.42
2:B:1934:MET:CE	2:B:1940:ILE:HD13	1.46	1.41
2:B:1738:PHE:CD2	2:B:1747:LEU:HD12	1.57	1.36
1:A:65:ARG:CG	1:A:66:PRO:HD3	1.57	1.35

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:250:ARG:NH1	1:A:496:LYS:NZ[7_555]	1.94	0.26

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	599/740 (81%)	448 (75%)	111 (18%)	40 (7%)	1	18
2	B	613/865 (71%)	498 (81%)	87 (14%)	28 (5%)	2	24
All	All	1212/1605 (76%)	946 (78%)	198 (16%)	68 (6%)	2	21

5 of 68 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	TRP
1	A	127	LYS
1	A	229	PRO
1	A	243	PRO
1	A	312	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	542/660 (82%)	415 (77%)	127 (23%)	1	5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	550/776 (71%)	413 (75%)	137 (25%)	0	4
All	All	1092/1436 (76%)	828 (76%)	264 (24%)	0	5

5 of 264 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	688	TRP
2	B	1781	ARG
2	B	2243	VAL
1	A	699	ASN
2	B	1708	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	HIS
1	A	232	HIS
1	A	615	HIS
1	A	684	ASN
2	B	2287	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

13 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	C	1	1,3	14,14,15	0.45	0	17,19,21	0.74	0
3	NAG	C	2	3	14,14,15	0.38	0	17,19,21	1.27	1 (5%)
3	BMA	C	3	3	11,11,12	0.34	0	15,15,17	1.84	3 (20%)
3	MAN	C	4	3	11,11,12	0.29	0	15,15,17	0.94	1 (6%)
4	NAG	D	1	2,4	14,14,15	0.63	0	17,19,21	1.48	2 (11%)
4	NAG	D	2	4	14,14,15	0.43	0	17,19,21	1.70	5 (29%)
5	NAG	E	1	2,5	14,14,15	0.51	0	17,19,21	2.48	5 (29%)
5	NAG	E	2	5	14,14,15	0.55	0	17,19,21	1.41	3 (17%)
5	BMA	E	3	5	11,11,12	0.58	0	15,15,17	1.08	1 (6%)
5	BMA	E	4	5	11,11,12	0.28	0	15,15,17	0.76	0
5	MAN	E	5	5	11,11,12	0.27	0	15,15,17	0.57	0
5	MAN	E	6	5	11,11,12	0.62	0	15,15,17	1.38	2 (13%)
5	MAN	E	7	5	11,11,12	0.48	0	15,15,17	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	2/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
4	NAG	D	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	4/6/23/26	0/1/1/1
5	NAG	E	1	2,5	-	2/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	2/2/19/22	0/1/1/1
5	BMA	E	4	5	-	0/2/19/22	0/1/1/1
5	MAN	E	5	5	-	0/2/19/22	0/1/1/1
5	MAN	E	6	5	-	0/2/19/22	0/1/1/1
5	MAN	E	7	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
5	E	1	NAG	O5-C5-C6	5.68	116.11	107.20
5	E	1	NAG	C2-N2-C7	5.50	130.73	122.90
3	C	3	BMA	C1-C2-C3	5.19	116.05	109.67
3	C	2	NAG	O5-C5-C6	3.88	113.28	107.20
4	D	2	NAG	C4-C3-C2	3.78	116.56	111.02

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	3	BMA	O5-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6
4	D	2	NAG	C8-C7-N2-C2


There are no ring outliers.


9 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0
5	E	6	MAN	2	0
5	E	3	BMA	3	0
5	E	7	MAN	1	0
5	E	2	NAG	1	0
3	C	3	BMA	7	0
3	C	4	MAN	4	0
3	C	2	NAG	4	0
5	E	5	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	611/740 (82%)	-0.42	1 (0%) 95 93	99, 186, 319, 397	0
2	B	619/865 (71%)	-0.36	3 (0%) 91 86	106, 195, 319, 424	0
All	All	1230/1605 (76%)	-0.39	4 (0%) 94 90	99, 191, 319, 424	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	2058	SER	2.4
1	A	190	GLN	2.3
2	B	1744	THR	2.3
2	B	1743	PHE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	D	2	14/15	0.64	0.42	307,363,404,410	0
5	BMA	E	3	11/12	0.69	0.23	278,295,314,335	0
4	NAG	D	1	14/15	0.70	0.45	275,361,420,429	0
5	MAN	E	7	11/12	0.74	0.32	303,330,373,383	0
3	MAN	C	4	11/12	0.78	0.17	193,218,224,226	11

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BMA	C	3	11/12	0.81	0.14	181,207,223,224	11
5	MAN	E	5	11/12	0.82	0.27	192,215,228,233	11
3	NAG	C	2	14/15	0.85	0.12	169,231,241,258	14
3	NAG	C	1	14/15	0.85	0.25	199,244,264,269	0
5	MAN	E	6	11/12	0.87	0.22	348,364,380,392	0
5	BMA	E	4	11/12	0.90	0.17	222,241,252,258	11
5	NAG	E	1	14/15	0.92	0.26	151,202,230,232	0
5	NAG	E	2	14/15	0.94	0.39	228,247,271,273	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	CA	A	806	1/1	0.93	0.21	236,236,236,236	0
7	CU	A	808	1/1	0.98	0.11	167,167,167,167	0
6	CA	A	805	1/1	0.99	0.14	215,215,215,215	0
6	CA	A	807	1/1	0.99	0.14	165,165,165,165	0
7	CU	B	2610	1/1	0.99	0.12	150,150,150,150	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.