



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 05:57 AM EDT

PDB ID : 1K9F
Title : Crystal structure of a mutated family-67 alpha-D-glucuronidase (E285N) from *Bacillus stearothermophilus* T-6, complexed with aldotetrauronic acid
Authors : Golan, G.; Shallom, D.; Teplitsky, A.; Zaide, G.; Shulami, S.; Baasov, T.; Stojanoff, V.; Thompson, A.; Shoham, Y.; Shoham, G.
Deposited on : 2001-10-29
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	679	
2	B	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GCV	B	3	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-D-glucuronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	671	Total	C	N	O	S	0	12	0
			5550	3542	952	1035	21			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	285	ASN	GLU	engineered mutation	GB 16876433

- Molecule 2 is an oligosaccharide called 4-O-methyl-alpha-D-glucopyranuronic acid-(1-2)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	3	Total	C	O	0	0	0
			32	17	15			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	1
			12	6	6		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	445	Total 445	O 445	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: alpha-D-glucuronidase



- Molecule 2: 4-O-methyl-alpha-D-glucopyranuronic acid-(1-2)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.56 Å 73.56 Å 330.01 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.55 – 1.75	Depositor
% Data completeness (in resolution range)	91.6 (36.55-1.75)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.189 , 0.211	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6105	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GCV, XYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/5701	0.59	0/7735

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5550	0	5290	51	0
2	B	32	0	8	0	0
3	A	78	0	104	8	0
4	A	445	0	0	2	0
All	All	6105	0	5402	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:HIS:HE1	1:A:321:VAL:HG21	1.44	0.80
1:A:87:HIS:HD2	1:A:89:GLU:H	1.28	0.79
1:A:526:HIS:HD2	1:A:528:TYR:H	1.44	0.64
1:A:77:PHE:O	1:A:78:GLU:HB3	1.99	0.63
1:A:526:HIS:CD2	1:A:528:TYR:H	2.16	0.63
1:A:297:HIS:CE1	1:A:321:VAL:HG21	2.33	0.62
1:A:174:VAL:HG12	1:A:176:GLN:H	1.66	0.61
1:A:31:THR:HG22	1:A:31:THR:O	2.02	0.59
1:A:674:GLY:HA2	4:A:1286:HOH:O	2.03	0.57
1:A:253:LEU:HD12	1:A:295:ARG:CZ	2.34	0.57
1:A:150:TRP:HA	1:A:199:ASN:HA	1.88	0.56
1:A:478:GLU:O	1:A:482:GLU:HG3	2.06	0.56
1:A:344:LYS:HB3	1:A:345:PRO:HD3	1.87	0.56
1:A:34:ILE:O	1:A:38:VAL:HG23	2.06	0.55
1:A:555:ARG:HB3	1:A:563:TYR:HB3	1.90	0.54
1:A:38:VAL:CG1	1:A:118:VAL:HG21	2.38	0.54
1:A:541:GLY:HA2	1:A:543:TYR:CE2	2.44	0.53
1:A:637:ARG:HD3	3:A:809:GOL:H11	1.92	0.52
1:A:309:ALA:HB3	1:A:310:PRO:HD3	1.92	0.51
1:A:594:TYR:HB3	1:A:605:ILE:HB	1.92	0.51
1:A:623:ARG:O	1:A:627:GLU:HG3	2.10	0.51
1:A:50[A]:MET:HB2	1:A:52:ILE:HG12	1.93	0.51
1:A:319:CYS:SG	1:A:356:LEU:HB3	2.51	0.50
1:A:409:GLU:HB3	3:A:802:GOL:H11	1.95	0.49
1:A:388:GLN:HG2	1:A:390:THR:O	2.14	0.48
1:A:382:ASN:HB3	1:A:437:TYR:O	2.14	0.48
1:A:97:VAL:HG12	1:A:97:VAL:O	2.14	0.47
1:A:320:PHE:HA	3:A:811:GOL:H11	1.97	0.47
1:A:355:ILE:HD11	1:A:437:TYR:HB3	1.96	0.47
1:A:75:GLU:OE2	1:A:107:ILE:HD13	2.15	0.46
1:A:597:ARG:HD2	1:A:601:GLY:O	2.16	0.46
1:A:359:LYS:HG3	3:A:811:GOL:H12	1.96	0.46
1:A:119:PHE:CG	1:A:189:SER:HA	2.51	0.45
1:A:418:LYS:HE3	1:A:433:PHE:CE1	2.52	0.45
1:A:55:GLN:HA	3:A:808:GOL:H11	1.99	0.44
1:A:484:VAL:HG11	1:A:495:VAL:HA	1.99	0.44
1:A:339:ALA:HB2	3:A:811:GOL:O2	2.17	0.44
1:A:302:ASN:O	1:A:306:GLU:HG3	2.18	0.44
1:A:624:LYS:O	1:A:628:GLN:HG3	2.18	0.44
1:A:236:ILE:HD12	1:A:236:ILE:C	2.38	0.43
1:A:6:GLU:HB3	3:A:809:GOL:H12	2.00	0.43
1:A:33:PRO:HD2	4:A:1227:HOH:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:GLN:HB2	1:A:369:GLU:HG3	2.02	0.42
1:A:397:GLN:O	1:A:455:HIS:HB2	2.18	0.42
1:A:636:LYS:HE3	1:A:640:ASP:OD2	2.20	0.41
1:A:635:GLU:OE1	1:A:639:HIS:HE1	2.03	0.41
1:A:408:LYS:HZ1	3:A:805:GOL:H2	1.86	0.41
1:A:523:ASN:O	1:A:527:HIS:HA	2.21	0.41
1:A:96:ASP:HB2	1:A:103:ARG:HB2	2.02	0.40
1:A:123:ARG:O	1:A:127[A]:MET:HG2	2.21	0.40
1:A:352:GLU:H	1:A:352:GLU:CD	2.25	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	679/679 (100%)	651 (96%)	26 (4%)	2 (0%)	41 22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	77	PHE
1	A	78	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	584/587 (100%)	581 (100%)	3 (0%)	88 83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	PHE
1	A	164	ARG
1	A	679	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	HIS
1	A	326	GLN
1	A	526	HIS
1	A	639	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	XYP	B	1	2	10,10,10	0.43	0	14,14,14	0.51	0
2	XYP	B	2	2	9,9,10	1.18	1 (11%)	10,12,14	1.56	1 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GCV	B	3	2	10,13,14	2.24	3 (30%)	11,18,20	5.86	10 (90%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	B	1	2	-	-	0/1/1/1
2	XYP	B	2	2	-	-	0/1/1/1
2	GCV	B	3	2	1/1/5/6	0/2/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	GCV	C2-C3	5.44	1.60	1.52
2	B	3	GCV	O5-C5	2.96	1.46	1.43
2	B	3	GCV	C3-C4	-2.49	1.45	1.52
2	B	2	XYP	C4-C3	2.05	1.55	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	GCV	O2-C2-C3	10.05	130.28	110.14
2	B	3	GCV	O4-C4-C3	8.24	130.00	110.29
2	B	3	GCV	O5-C1-C2	7.11	121.74	110.77
2	B	3	GCV	O3-C3-C2	-6.96	96.66	109.99
2	B	3	GCV	C3-C4-C5	-5.90	102.72	110.28
2	B	3	GCV	C2-C3-C4	5.44	121.34	110.41
2	B	3	GCV	C1-C2-C3	-4.68	103.92	109.67
2	B	2	XYP	C1-C2-C3	-4.14	104.58	109.67
2	B	3	GCV	C7-O4-C4	3.42	123.49	114.52
2	B	3	GCV	O3-C3-C4	2.44	116.40	109.94
2	B	3	GCV	C1-O5-C5	-2.31	108.10	112.17

All (1) chirality outliers are listed below:


Mol	Chain	Res	Type	Atom
2	B	3	GCV	C5

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	812	-	5,5,5	0.75	0	5,5,5	0.45	0
3	GOL	A	809	-	5,5,5	0.79	0	5,5,5	0.41	0
3	GOL	A	807	-	5,5,5	0.73	0	5,5,5	0.39	0
3	GOL	A	811	-	5,5,5	0.80	0	5,5,5	0.44	0
3	GOL	A	801	-	5,5,5	0.73	0	5,5,5	0.46	0
3	GOL	A	810	-	5,5,5	0.80	0	5,5,5	0.43	0
3	GOL	A	804	-	5,5,5	0.77	0	5,5,5	0.43	0
3	GOL	A	806	-	5,5,5	0.78	0	5,5,5	0.46	0
3	GOL	A	808	-	5,5,5	0.76	0	5,5,5	0.44	0
3	GOL	A	802	-	5,5,5	0.76	0	5,5,5	0.45	0
3	GOL	A	805	-	5,5,5	0.76	0	5,5,5	0.43	0
3	GOL	A	803[B]	-	5,5,5	0.76	0	5,5,5	0.39	0
3	GOL	A	803[A]	-	5,5,5	0.75	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	812	-	-	1/4/4/4	-
3	GOL	A	809	-	-	2/4/4/4	-
3	GOL	A	807	-	-	2/4/4/4	-
3	GOL	A	811	-	-	0/4/4/4	-
3	GOL	A	801	-	-	0/4/4/4	-
3	GOL	A	810	-	-	0/4/4/4	-
3	GOL	A	804	-	-	0/4/4/4	-
3	GOL	A	806	-	-	4/4/4/4	-
3	GOL	A	808	-	-	1/4/4/4	-
3	GOL	A	802	-	-	0/4/4/4	-
3	GOL	A	805	-	-	2/4/4/4	-
3	GOL	A	803[B]	-	-	1/4/4/4	-
3	GOL	A	803[A]	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	805	GOL	O1-C1-C2-C3
3	A	807	GOL	O1-C1-C2-C3
3	A	803[A]	GOL	O1-C1-C2-C3
3	A	806	GOL	C1-C2-C3-O3
3	A	808	GOL	C1-C2-C3-O3
3	A	809	GOL	C1-C2-C3-O3
3	A	812	GOL	C1-C2-C3-O3
3	A	803[A]	GOL	O1-C1-C2-O2
3	A	805	GOL	O1-C1-C2-O2
3	A	807	GOL	O1-C1-C2-O2
3	A	806	GOL	O2-C2-C3-O3
3	A	809	GOL	O2-C2-C3-O3
3	A	806	GOL	O1-C1-C2-C3
3	A	803[B]	GOL	O1-C1-C2-C3
3	A	806	GOL	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	809	GOL	2	0
3	A	811	GOL	3	0
3	A	808	GOL	1	0
3	A	802	GOL	1	0
3	A	805	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.