



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 02:58 PM BST

PDB ID : 5K9K
Title : Crystal structure of multidonor HV6-1-class broadly neutralizing Influenza A antibody 56.a.09 in complex with Hemagglutinin Hong Kong 1968.
Authors : Joyce, M.G.; Thomas, P.V.; Wheatley, A.K.; McDermott, A.B.; Mascola, J.R.; Kwong, P.D.
Deposited on : 2016-05-31
Resolution : 2.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2754 (3.00-2.96)
Clashscore	141614	3103 (3.00-2.96)
Ramachandran outliers	138981	2993 (3.00-2.96)
Sidechain outliers	138945	2996 (3.00-2.96)
RSRZ outliers	127900	2644 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	<div> <div>3%</div> <div>79%</div> <div>18%</div> <div>..</div> </div>
1	H	229	<div> <div>%</div> <div>79%</div> <div>17%</div> <div>..</div> </div>
2	B	215	<div> <div>2%</div> <div>76%</div> <div>21%</div> <div>.</div> </div>
2	L	215	<div> <div>3%</div> <div>79%</div> <div>20%</div> <div>.</div> </div>
3	F	503	<div> <div>9%</div> <div>73%</div> <div>23%</div> <div>..</div> </div>
3	I	503	<div> <div>6%</div> <div>77%</div> <div>18%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	C	5	
4	J	5	
4	K	5	
4	P	5	
5	D	3	
5	N	3	
6	E	2	
7	G	5	
8	M	6	
9	O	5	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	NAG	F	601	-	-	-	X
10	NAG	I	601	-	-	-	X
10	NAG	I	602	-	-	-	X
4	NAG	K	1	-	-	-	X
4	NAG	K	2	-	-	-	X
4	BMA	K	3	-	-	-	X
4	MAN	K	4	-	-	-	X
4	MAN	K	5	-	-	-	X
4	MAN	P	4	-	-	-	X
4	MAN	P	5	-	-	-	X
5	NAG	D	2	-	-	-	X
5	BMA	D	3	-	-	-	X
5	NAG	N	2	-	-	-	X
5	BMA	N	3	-	-	-	X
6	NAG	E	1	X	-	-	X
7	MAN	G	5	-	-	-	X
9	MAN	O	5	-	-	-	X

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 15108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 56.a.09 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1710	1082	288	333	7			
1	H	224	Total	C	N	O	S	0	0	0
			1695	1073	285	330	7			

- Molecule 2 is a protein called 56.a.09 Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	215	Total	C	N	O	S	0	0	0
			1652	1032	279	336	5			
2	L	215	Total	C	N	O	S	0	0	0
			1652	1032	279	336	5			


- Molecule 3 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	490	Total	C	N	O	S	0	1	0
			3876	2416	681	760	19			
3	F	497	Total	C	N	O	S	0	1	0
			3923	2447	691	766	19			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	218	GLU	GLY	conflict	UNP Q91MA7
F	218	GLU	GLY	conflict	UNP Q91MA7

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	5	Total	C	N	O	0	0	0
			61	34	2	25			
4	J	5	Total	C	N	O	0	0	0
			61	34	2	25			
4	K	5	Total	C	N	O	0	0	0
			61	34	2	25			
4	P	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	D	3	Total	C	N	O	0	0	0
			39	22	2	15			
5	N	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	G	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	M	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 9 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	O	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	I	1	Total	C	N	O	0	0
			14	8	1	5		
10	I	1	Total	C	N	O	0	0
			14	8	1	5		
10	F	1	Total	C	N	O	0	0
			14	8	1	5		
10	F	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 56.a.09 Heavy chain




- Molecule 1: 56.a.09 Heavy chain




- Molecule 2: 56.a.09 Light chain





- Molecule 2: 56.a.09 Light chain








• Molecule 3: Hemagglutinin



• Molecule 3: Hemagglutinin



• Molecule 4: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  60% 40%



- Molecule 4: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain J:  20% 60% 20%



- Molecule 4: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain K:  60% 40%



- Molecule 4: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain P:  40% 60%



- Molecule 5: β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain D:  100%



- Molecule 5: β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain N:  100%




- Molecule 6: 2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain E:  50% 50%




- Molecule 7: alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 8: alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 9: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	123.90Å 136.56Å 311.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.01 – 2.97 46.01 – 2.97	Depositor EDS
% Data completeness (in resolution range)	75.0 (46.01-2.97) 71.6 (46.01-2.97)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.96Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, R_{free}	0.225 , 0.265 0.229 , 0.259	Depositor DCC
R_{free} test set	2058 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	60.9	Xtriage
Anisotropy	0.419	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 20.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	15108	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1750	0.55	1/2386 (0.0%)
1	H	0.36	0/1735	0.52	0/2367
2	B	0.41	0/1687	0.49	0/2288
2	L	0.42	0/1687	0.49	0/2288
3	F	0.34	0/4009	0.52	0/5436
3	I	0.38	0/3960	0.53	3/5372 (0.1%)
All	All	0.38	0/14828	0.52	4/20137 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	I	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	148	GLU	C-N-CD	-9.57	99.55	120.60
3	I	350[A]	TRP	CB-CA-C	-5.78	98.84	110.40
3	I	350[B]	TRP	CB-CA-C	-5.78	98.84	110.40
3	I	404	GLY	N-CA-C	-5.36	99.70	113.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	I	350[A]	TRP	Mainchain
3	I	350[B]	TRP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1710	0	1709	36	0
1	H	1695	0	1691	36	0
2	B	1652	0	1599	47	0
2	L	1652	0	1599	27	0
3	F	3923	0	3776	133	0
3	I	3876	0	3729	84	0
4	C	61	0	52	0	0
4	J	61	0	52	3	0
4	K	61	0	52	0	0
4	P	61	0	52	0	0
5	D	39	0	34	0	0
5	N	39	0	34	0	0
6	E	28	0	25	0	0
7	G	61	0	52	0	0
8	M	72	0	61	1	0
9	O	61	0	52	0	0
10	F	28	0	26	0	0
10	I	28	0	26	0	0
All	All	15108	0	14621	354	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 354 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:402:VAL:CG1	3:F:409:LEU:HD11	1.52	1.39
3:F:402:VAL:CG1	3:F:409:LEU:CD1	2.06	1.31
3:F:402:VAL:HG13	3:F:409:LEU:CD1	1.64	1.24
3:F:156:LYS:HE2	3:F:193:SER:O	1.40	1.21

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:27:GLN:OE1	4:J:1:NAG:H81	1.43	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/229 (97%)	210 (95%)	11 (5%)	1 (0%)	29	66
1	H	220/229 (96%)	207 (94%)	13 (6%)	0	100	100
2	B	213/215 (99%)	207 (97%)	6 (3%)	0	100	100
2	L	213/215 (99%)	208 (98%)	5 (2%)	0	100	100
3	F	494/503 (98%)	470 (95%)	24 (5%)	0	100	100
3	I	487/503 (97%)	467 (96%)	20 (4%)	0	100	100
All	All	1849/1894 (98%)	1769 (96%)	79 (4%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	149	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/200 (99%)	190 (96%)	8 (4%)	31	66
1	H	196/200 (98%)	187 (95%)	9 (5%)	27	61
2	B	185/185 (100%)	171 (92%)	14 (8%)	13	41
2	L	185/185 (100%)	177 (96%)	8 (4%)	29	64
3	F	435/440 (99%)	411 (94%)	24 (6%)	21	55
3	I	432/440 (98%)	412 (95%)	20 (5%)	27	61
All	All	1631/1650 (99%)	1548 (95%)	83 (5%)	24	57

5 of 83 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	I	407	GLN
1	H	124	LEU
3	F	329	ARG
3	I	409	LEU
1	H	1	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
3	I	371	GLN
3	F	75	HIS
2	L	147	GLN
2	B	158	ASN
1	H	1	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

44 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	1	3,4	14,14,15	0.39	0	17,19,21	0.55	0
4	NAG	C	2	4	14,14,15	0.24	0	17,19,21	0.78	0
4	BMA	C	3	4	11,11,12	0.54	0	15,15,17	0.78	0
4	MAN	C	4	4	11,11,12	0.59	0	15,15,17	0.98	2 (13%)
4	MAN	C	5	4	11,11,12	0.64	0	15,15,17	1.03	2 (13%)
5	NAG	D	1	3,5	14,14,15	0.37	0	17,19,21	0.56	0
5	NAG	D	2	5	14,14,15	0.27	0	17,19,21	0.46	0
5	BMA	D	3	5	11,11,12	0.56	0	15,15,17	0.78	0
6	NAG	E	1	3,6	14,14,15	0.47	0	17,19,21	1.84	3 (17%)
6	NAG	E	2	6	14,14,15	0.28	0	17,19,21	0.48	0
7	NAG	G	1	3,7	14,14,15	0.31	0	17,19,21	0.50	0
7	NAG	G	2	7	14,14,15	0.37	0	17,19,21	0.79	1 (5%)
7	BMA	G	3	7	11,11,12	0.68	0	15,15,17	1.13	1 (6%)
7	MAN	G	4	7	11,11,12	0.77	0	15,15,17	0.95	2 (13%)
7	MAN	G	5	7	11,11,12	0.71	0	15,15,17	1.09	2 (13%)
4	NAG	J	1	3,4	14,14,15	0.39	0	17,19,21	0.76	1 (5%)
4	NAG	J	2	4	14,14,15	0.22	0	17,19,21	0.56	0
4	BMA	J	3	4	11,11,12	1.73	2 (18%)	15,15,17	1.54	2 (13%)
4	MAN	J	4	4	11,11,12	0.60	0	15,15,17	1.02	2 (13%)
4	MAN	J	5	4	11,11,12	0.75	0	15,15,17	1.15	2 (13%)
4	NAG	K	1	3,4	14,14,15	0.34	0	17,19,21	0.46	0
4	NAG	K	2	4	14,14,15	0.25	0	17,19,21	0.51	0
4	BMA	K	3	4	11,11,12	0.73	0	15,15,17	0.91	0
4	MAN	K	4	4	11,11,12	0.63	0	15,15,17	1.01	2 (13%)
4	MAN	K	5	4	11,11,12	0.84	1 (9%)	15,15,17	1.34	2 (13%)
8	NAG	M	1	8,3	14,14,15	0.36	0	17,19,21	0.64	0
8	NAG	M	2	8	14,14,15	0.22	0	17,19,21	0.73	0
8	BMA	M	3	8	11,11,12	0.62	0	15,15,17	0.86	0
8	MAN	M	4	8	11,11,12	0.78	0	15,15,17	1.14	2 (13%)
8	MAN	M	5	8	11,11,12	0.74	0	15,15,17	0.96	1 (6%)
8	MAN	M	6	8	11,11,12	0.63	0	15,15,17	1.01	2 (13%)
5	NAG	N	1	3,5	14,14,15	0.53	0	17,19,21	0.65	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	N	2	5	14,14,15	0.29	0	17,19,21	0.38	0
5	BMA	N	3	5	11,11,12	0.65	0	15,15,17	0.71	0
9	NAG	O	1	9,3	14,14,15	1.03	1 (7%)	17,19,21	0.83	0
9	NAG	O	2	9	14,14,15	0.21	0	17,19,21	0.54	0
9	BMA	O	3	9	11,11,12	0.58	0	15,15,17	0.75	0
9	MAN	O	4	9	11,11,12	0.70	0	15,15,17	0.94	1 (6%)
9	MAN	O	5	9	11,11,12	0.62	0	15,15,17	1.00	2 (13%)
4	NAG	P	1	3,4	14,14,15	0.37	0	17,19,21	0.49	0
4	NAG	P	2	4	14,14,15	0.34	0	17,19,21	0.68	0
4	BMA	P	3	4	11,11,12	0.88	0	15,15,17	1.49	1 (6%)
4	MAN	P	4	4	11,11,12	0.67	0	15,15,17	1.00	2 (13%)
4	MAN	P	5	4	11,11,12	0.93	0	15,15,17	0.91	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	3,4	-	2/6/23/26	0/1/1/1
4	NAG	C	2	4	-	2/6/23/26	0/1/1/1
4	BMA	C	3	4	-	0/2/19/22	0/1/1/1
4	MAN	C	4	4	-	1/2/19/22	0/1/1/1
4	MAN	C	5	4	-	0/2/19/22	0/1/1/1
5	NAG	D	1	3,5	-	2/6/23/26	0/1/1/1
5	NAG	D	2	5	-	0/6/23/26	0/1/1/1
5	BMA	D	3	5	-	0/2/19/22	0/1/1/1
6	NAG	E	1	3,6	1/1/5/7	2/6/23/26	0/1/1/1
6	NAG	E	2	6	-	1/6/23/26	0/1/1/1
7	NAG	G	1	3,7	-	0/6/23/26	0/1/1/1
7	NAG	G	2	7	-	0/6/23/26	0/1/1/1
7	BMA	G	3	7	-	2/2/19/22	0/1/1/1
7	MAN	G	4	7	-	2/2/19/22	0/1/1/1
7	MAN	G	5	7	-	0/2/19/22	1/1/1/1
4	NAG	J	1	3,4	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	0/6/23/26	0/1/1/1
4	BMA	J	3	4	-	0/2/19/22	0/1/1/1
4	MAN	J	4	4	-	0/2/19/22	0/1/1/1
4	MAN	J	5	4	-	0/2/19/22	1/1/1/1
4	NAG	K	1	3,4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	K	2	4	-	3/6/23/26	0/1/1/1
4	BMA	K	3	4	-	2/2/19/22	0/1/1/1
4	MAN	K	4	4	-	0/2/19/22	0/1/1/1
4	MAN	K	5	4	-	0/2/19/22	0/1/1/1
8	NAG	M	1	8,3	-	3/6/23/26	0/1/1/1
8	NAG	M	2	8	-	2/6/23/26	0/1/1/1
8	BMA	M	3	8	-	0/2/19/22	0/1/1/1
8	MAN	M	4	8	-	0/2/19/22	0/1/1/1
8	MAN	M	5	8	-	0/2/19/22	0/1/1/1
8	MAN	M	6	8	-	0/2/19/22	0/1/1/1
5	NAG	N	1	3,5	-	2/6/23/26	0/1/1/1
5	NAG	N	2	5	-	0/6/23/26	0/1/1/1
5	BMA	N	3	5	-	0/2/19/22	0/1/1/1
9	NAG	O	1	9,3	-	0/6/23/26	0/1/1/1
9	NAG	O	2	9	-	3/6/23/26	0/1/1/1
9	BMA	O	3	9	-	1/2/19/22	0/1/1/1
9	MAN	O	4	9	-	1/2/19/22	0/1/1/1
9	MAN	O	5	9	-	0/2/19/22	0/1/1/1
4	NAG	P	1	3,4	-	0/6/23/26	0/1/1/1
4	NAG	P	2	4	-	0/6/23/26	0/1/1/1
4	BMA	P	3	4	-	2/2/19/22	1/1/1/1
4	MAN	P	4	4	-	0/2/19/22	0/1/1/1
4	MAN	P	5	4	-	0/2/19/22	1/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	J	3	BMA	C1-C2	4.24	1.61	1.52
9	O	1	NAG	O5-C1	-3.64	1.37	1.43
4	J	3	BMA	C2-C3	3.39	1.57	1.52
4	K	5	MAN	C1-C2	2.44	1.57	1.52

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	1	NAG	C1-O5-C5	6.08	120.42	112.19
4	P	3	BMA	C1-O5-C5	4.46	118.23	112.19
4	J	3	BMA	C1-C2-C3	4.01	114.60	109.67
4	K	5	MAN	C1-O5-C5	3.71	117.22	112.19
4	J	5	MAN	C1-O5-C5	3.23	116.57	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	E	1	NAG	C1

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	P	3	BMA	O5-C5-C6-O6
5	D	1	NAG	O5-C5-C6-O6
7	G	3	BMA	C4-C5-C6-O6
5	N	1	NAG	O5-C5-C6-O6
5	D	1	NAG	C4-C5-C6-O6









All (4) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	P	3	BMA	C1-C2-C3-C4-C5-O5
4	P	5	MAN	C1-C2-C3-C4-C5-O5
7	G	5	MAN	C1-C2-C3-C4-C5-O5
4	J	5	MAN	C1-C2-C3-C4-C5-O5





2 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	J	1	NAG	3	0
8	M	6	MAN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain G**Bond lengths****Bond angles****Torsions****Rings****Oligosaccharide Chain K****Bond lengths****Bond angles****Torsions****Rings**



Oligosaccharide Chain J**Bond lengths****Bond angles****Torsions****Rings**





5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	NAG	I	602	3	14,14,15	0.24	0	17,19,21	0.45	0
10	NAG	I	601	-	14,14,15	0.21	0	17,19,21	0.40	0
10	NAG	F	601	3	14,14,15	0.18	0	17,19,21	0.40	0
10	NAG	F	621	3	14,14,15	0.26	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	I	602	3	-	1/6/23/26	0/1/1/1
10	NAG	I	601	-	-	0/6/23/26	0/1/1/1
10	NAG	F	601	3	-	1/6/23/26	0/1/1/1
10	NAG	F	621	3	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	F	621	NAG	O5-C5-C6-O6
10	F	601	NAG	O5-C5-C6-O6
10	I	602	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/229 (98%)	0.37	6 (2%) 54 35	37, 63, 98, 162	1 (0%)
1	H	224/229 (97%)	0.33	3 (1%) 77 59	24, 65, 94, 155	0
2	B	215/215 (100%)	0.31	5 (2%) 60 40	45, 68, 97, 149	1 (0%)
2	L	215/215 (100%)	0.41	6 (2%) 53 34	37, 61, 100, 162	1 (0%)
3	F	497/503 (98%)	0.68	46 (9%) 8 5	39, 85, 137, 182	2 (0%)
3	I	490/503 (97%)	0.45	29 (5%) 22 12	49, 70, 117, 168	4 (0%)
All	All	1867/1894 (98%)	0.47	95 (5%) 28 16	24, 70, 122, 182	9 (0%)

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	I	6	ASN	10.2
2	B	214	CYS	10.2
3	F	4	PRO	9.5
3	F	5	GLY	7.0
3	F	128	THR	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	BMA	D	3	11/12	0.24	0.48	134,151,164,164	0
4	MAN	K	5	11/12	0.32	1.04	187,191,196,197	0
4	MAN	K	4	11/12	0.39	0.75	171,189,191,193	0
5	BMA	N	3	11/12	0.43	0.56	168,171,173,173	0
7	MAN	G	5	11/12	0.43	0.83	159,168,173,177	0
5	NAG	N	2	14/15	0.44	0.51	131,148,158,165	0
7	MAN	G	4	11/12	0.44	0.39	140,153,157,163	0
4	NAG	K	2	14/15	0.45	0.79	185,188,196,197	0
4	BMA	K	3	11/12	0.50	0.73	194,195,199,199	0
5	NAG	N	1	14/15	0.58	0.32	109,135,142,144	0
4	NAG	K	1	14/15	0.58	0.68	124,150,158,173	0
9	MAN	O	4	11/12	0.59	0.24	150,158,162,163	0
5	NAG	D	2	14/15	0.60	0.46	121,147,156,161	0
7	BMA	G	3	11/12	0.62	0.25	127,131,145,151	0
4	MAN	C	5	11/12	0.62	0.24	82,102,117,121	0
4	MAN	P	5	11/12	0.62	0.41	105,126,133,133	0
4	MAN	J	4	11/12	0.63	0.37	149,153,156,164	0
4	MAN	P	4	11/12	0.67	0.43	140,143,146,148	0
8	MAN	M	4	11/12	0.69	0.26	114,121,126,127	0
9	BMA	O	3	11/12	0.70	0.32	128,131,151,154	0
9	MAN	O	5	11/12	0.71	0.42	150,154,156,161	0
4	BMA	J	3	11/12	0.73	0.24	129,146,152,155	0
4	BMA	P	3	11/12	0.74	0.24	128,134,137,139	0
4	MAN	J	5	11/12	0.75	0.28	117,124,127,130	0
6	NAG	E	2	14/15	0.76	0.37	127,145,172,173	0
6	NAG	E	1	14/15	0.77	0.46	103,136,165,170	0
8	MAN	M	5	11/12	0.78	0.31	127,130,136,137	0
4	NAG	J	2	14/15	0.80	0.28	108,115,138,139	0
4	MAN	C	4	11/12	0.80	0.23	77,80,86,92	0
9	NAG	O	2	14/15	0.81	0.27	73,107,115,121	0
5	NAG	D	1	14/15	0.84	0.22	87,99,121,129	0
7	NAG	G	1	14/15	0.85	0.20	59,83,90,100	0
7	NAG	G	2	14/15	0.86	0.22	81,100,113,121	0
8	MAN	M	6	11/12	0.86	0.21	96,101,115,118	0
4	NAG	P	2	14/15	0.87	0.18	86,96,108,118	0
4	NAG	J	1	14/15	0.88	0.24	68,85,104,107	0
4	NAG	P	1	14/15	0.89	0.20	60,71,93,97	0
8	NAG	M	2	14/15	0.89	0.17	32,51,58,72	0
8	BMA	M	3	11/12	0.89	0.11	59,75,97,106	0
9	NAG	O	1	14/15	0.90	0.19	63,81,93,96	0
4	BMA	C	3	11/12	0.90	0.12	63,70,79,89	0
8	NAG	M	1	14/15	0.93	0.16	26,38,57,61	0
4	NAG	C	2	14/15	0.93	0.14	32,41,56,56	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	C	1	14/15	0.94	0.13	29,35,60,70	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain P:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain N:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain M:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	NAG	I	602	14/15	0.37	0.59	118,133,151,151	0
10	NAG	F	601	14/15	0.52	0.79	127,140,147,149	0
10	NAG	I	601	14/15	0.62	0.58	128,134,148,150	0
10	NAG	F	621	14/15	0.74	0.33	96,110,124,127	0

6.5 Other polymers [i](#)

There are no such residues in this entry.