



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 08:38 AM BST

PDB ID : 5KAK
Title : Crystal structure of an uncharacterized NIPSNAP-like domain protein from *Burkholderia xenovorans*
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2016-06-01
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	126	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>8%</div> </div> </div>
1	B	126	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>7%</div> </div> </div>
1	C	126	<div> <div>11%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>6%</div> </div> </div>
1	D	126	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>11%</div> <div>6%</div> </div> </div>
1	E	126	<div> <div>4%</div> <div> <div></div> <div>88%</div> <div>7%</div> <div>5%</div> </div> </div>
1	F	126	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>6%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	126	<div><div></div><div>4%</div><div>82%</div><div>11%</div><div>7%</div></div>
1	H	126	<div><div></div><div>5%</div><div>83%</div><div>12%</div><div>5%</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9050 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	116	Total	C	N	O	S	0	5	0
			984	639	173	167	5			
1	B	117	Total	C	N	O	S	0	2	0
			965	626	167	167	5			
1	C	118	Total	C	N	O	S	0	4	0
			974	633	170	166	5			
1	D	119	Total	C	N	O	S	0	3	0
			999	645	179	171	4			
1	E	120	Total	C	N	O	S	0	3	0
			1001	647	182	168	4			
1	F	118	Total	C	N	O	S	0	4	0
			996	643	178	171	4			
1	G	117	Total	C	N	O	S	0	1	0
			963	623	166	170	4			
1	H	120	Total	C	N	O	S	0	4	0
			1023	658	186	175	4			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	expression tag	UNP Q13PY7
A	-7	ALA	-	expression tag	UNP Q13PY7
A	-6	HIS	-	expression tag	UNP Q13PY7
A	-5	HIS	-	expression tag	UNP Q13PY7
A	-4	HIS	-	expression tag	UNP Q13PY7
A	-3	HIS	-	expression tag	UNP Q13PY7
A	-2	HIS	-	expression tag	UNP Q13PY7
A	-1	HIS	-	expression tag	UNP Q13PY7
B	-8	MET	-	expression tag	UNP Q13PY7
B	-7	ALA	-	expression tag	UNP Q13PY7
B	-6	HIS	-	expression tag	UNP Q13PY7
B	-5	HIS	-	expression tag	UNP Q13PY7
B	-4	HIS	-	expression tag	UNP Q13PY7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	HIS	-	expression tag	UNP Q13PY7
B	-2	HIS	-	expression tag	UNP Q13PY7
B	-1	HIS	-	expression tag	UNP Q13PY7
C	-8	MET	-	expression tag	UNP Q13PY7
C	-7	ALA	-	expression tag	UNP Q13PY7
C	-6	HIS	-	expression tag	UNP Q13PY7
C	-5	HIS	-	expression tag	UNP Q13PY7
C	-4	HIS	-	expression tag	UNP Q13PY7
C	-3	HIS	-	expression tag	UNP Q13PY7
C	-2	HIS	-	expression tag	UNP Q13PY7
C	-1	HIS	-	expression tag	UNP Q13PY7
D	-8	MET	-	expression tag	UNP Q13PY7
D	-7	ALA	-	expression tag	UNP Q13PY7
D	-6	HIS	-	expression tag	UNP Q13PY7
D	-5	HIS	-	expression tag	UNP Q13PY7
D	-4	HIS	-	expression tag	UNP Q13PY7
D	-3	HIS	-	expression tag	UNP Q13PY7
D	-2	HIS	-	expression tag	UNP Q13PY7
D	-1	HIS	-	expression tag	UNP Q13PY7
E	-8	MET	-	expression tag	UNP Q13PY7
E	-7	ALA	-	expression tag	UNP Q13PY7
E	-6	HIS	-	expression tag	UNP Q13PY7
E	-5	HIS	-	expression tag	UNP Q13PY7
E	-4	HIS	-	expression tag	UNP Q13PY7
E	-3	HIS	-	expression tag	UNP Q13PY7
E	-2	HIS	-	expression tag	UNP Q13PY7
E	-1	HIS	-	expression tag	UNP Q13PY7
F	-8	MET	-	expression tag	UNP Q13PY7
F	-7	ALA	-	expression tag	UNP Q13PY7
F	-6	HIS	-	expression tag	UNP Q13PY7
F	-5	HIS	-	expression tag	UNP Q13PY7
F	-4	HIS	-	expression tag	UNP Q13PY7
F	-3	HIS	-	expression tag	UNP Q13PY7
F	-2	HIS	-	expression tag	UNP Q13PY7
F	-1	HIS	-	expression tag	UNP Q13PY7
G	-8	MET	-	expression tag	UNP Q13PY7
G	-7	ALA	-	expression tag	UNP Q13PY7
G	-6	HIS	-	expression tag	UNP Q13PY7
G	-5	HIS	-	expression tag	UNP Q13PY7
G	-4	HIS	-	expression tag	UNP Q13PY7
G	-3	HIS	-	expression tag	UNP Q13PY7
G	-2	HIS	-	expression tag	UNP Q13PY7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-1	HIS	-	expression tag	UNP Q13PY7
H	-8	MET	-	expression tag	UNP Q13PY7
H	-7	ALA	-	expression tag	UNP Q13PY7
H	-6	HIS	-	expression tag	UNP Q13PY7
H	-5	HIS	-	expression tag	UNP Q13PY7
H	-4	HIS	-	expression tag	UNP Q13PY7
H	-3	HIS	-	expression tag	UNP Q13PY7
H	-2	HIS	-	expression tag	UNP Q13PY7
H	-1	HIS	-	expression tag	UNP Q13PY7

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	F	1	Total C O 4 2 2	0	0
2	H	1	Total C O 4 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	141	Total 145	O 145	0	4
3	B	149	Total 159	O 159	0	9
3	C	103	Total 105	O 105	0	2
3	D	163	Total 167	O 167	0	4
3	E	136	Total 142	O 142	0	6
3	F	123	Total 124	O 124	0	1
3	G	125	Total 128	O 128	0	3
3	H	149	Total 151	O 151	0	2


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Uncharacterized protein





- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein




- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.63 Å 70.83 Å 109.02 Å 90.00° 98.62° 90.00°	Depositor
Resolution (Å)	43.43 – 1.65 43.43 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.43-1.65) 99.7 (43.43-1.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.65 Å)	Xtriage
Refinement program	PHENIX (dev_2386: ???)	Depositor
R, R_{free}	0.206 , 0.251 0.206 , 0.251	Depositor DCC
R_{free} test set	1992 reflections (1.56%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 66.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9050	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/1027	0.56	0/1397
1	B	0.37	0/999	0.57	0/1361
1	C	0.35	0/1014	0.52	0/1385
1	D	0.38	0/1037	0.57	0/1412
1	E	0.35	0/1039	0.53	0/1413
1	F	0.35	0/1035	0.54	0/1409
1	G	0.37	0/993	0.59	0/1353
1	H	0.35	0/1060	0.56	0/1440
All	All	0.36	0/8204	0.56	0/11170

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	984	0	983	9	0
1	B	965	0	943	8	0
1	C	974	0	953	16	0
1	D	999	0	984	9	0
1	E	1001	0	991	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	996	0	991	10	0
1	G	963	0	942	9	0
1	H	1023	0	1017	13	0
2	A	4	0	6	1	0
2	C	8	0	12	4	0
2	D	4	0	6	1	0
2	F	4	0	6	3	0
2	H	4	0	6	3	0
3	A	145	0	0	3	1
3	B	159	0	0	3	0
3	C	105	0	0	2	0
3	D	167	0	0	1	1
3	E	142	0	0	0	0
3	F	124	0	0	0	0
3	G	128	0	0	2	0
3	H	151	0	0	1	0
All	All	9050	0	7840	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:22[B]:GLU:OE2	3:C:301:HOH:O	2.08	0.71
1:H:52:HIS:CE1	2:H:201:EDO:H12	2.27	0.69
1:G:91:GLN:NE2	3:G:201:HOH:O	2.19	0.66
1:H:29:GLN:OE1	2:H:201:EDO:H11	2.01	0.61
1:B:57:GLU:OE1	3:B:201:HOH:O	2.16	0.60
1:D:89:ARG:NH2	3:D:303:HOH:O	2.34	0.59
1:B:109:GLN:NE2	3:B:205:HOH:O	2.35	0.56
1:A:66:ARG:HH12	2:A:201:EDO:H11	1.71	0.56
1:D:100:PHE:HB3	2:D:201:EDO:H12	1.88	0.56
1:C:0:MET:SD	1:C:104:LEU:HG	2.47	0.53
1:H:52:HIS:HE1	2:H:201:EDO:H12	1.73	0.53
1:B:76:GLN:HG2	3:B:300:HOH:O	2.09	0.53
1:C:95:LEU:HB3	1:D:40:PHE:HB3	1.91	0.53
1:F:47:LEU:HD13	2:F:201:EDO:H12	1.90	0.52
1:G:7:TYR:CD1	1:G:90:MET:HG2	2.44	0.51
1:G:0:MET:HB2	1:G:104:LEU:HD13	1.91	0.51
1:E:1:ILE:HD13	1:E:59:LEU:HD23	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:28:ILE:HD11	2:C:202:EDO:C2	2.41	0.51
1:D:0:MET:HB3	1:D:104:LEU:HD22	1.93	0.51
1:F:84[A]:GLN:NE2	1:H:12:ASN:HB2	2.25	0.51
1:C:28:ILE:HD11	2:C:202:EDO:H21	1.93	0.49
1:H:84:GLN:HB3	1:H:85:PRO:HD3	1.95	0.49
1:H:7:TYR:HB2	1:H:50:VAL:CG2	2.43	0.48
1:A:84[B]:GLN:HG3	3:A:397:HOH:O	2.12	0.48
1:D:19:GLN:HA	1:D:22[B]:GLU:HG2	1.96	0.48
1:C:22[A]:GLU:HG2	1:D:99:SER:OG	2.13	0.48
1:C:108:TRP:CH2	1:C:112:ASP:HB3	2.48	0.48
1:F:84[B]:GLN:HA	1:F:87:LEU:HD12	1.96	0.47
1:F:84[A]:GLN:HA	1:F:87:LEU:HD12	1.96	0.47
1:C:84[B]:GLN:HG3	3:C:367:HOH:O	2.14	0.47
1:H:0:MET:SD	1:H:104:LEU:HG	2.55	0.46
1:A:68:ILE:O	1:A:72:MET:HG3	2.16	0.45
1:F:7:TYR:HB2	1:F:50:VAL:HG22	1.97	0.45
1:G:0:MET:HB3	1:G:104:LEU:HD22	1.98	0.45
1:B:104:LEU:HD12	1:B:105:PRO:HD2	1.98	0.45
1:G:18:ALA:O	1:G:22[B]:GLU:HG3	2.16	0.45
1:C:55:GLY:HA3	1:C:104:LEU:CD2	2.47	0.45
1:D:16:GLU:HG2	1:D:20:HIS:CD2	2.52	0.45
1:C:7:TYR:HB2	1:C:50:VAL:CG2	2.46	0.45
1:E:16:GLU:HG2	1:E:20:HIS:CD2	2.52	0.45
1:F:47:LEU:HD22	2:F:201:EDO:H12	1.99	0.45
1:G:100:PHE:CD1	1:H:26:ILE:HG12	2.52	0.45
1:G:76:GLN:HG2	3:G:268:HOH:O	2.17	0.45
1:A:95:LEU:HB3	1:B:40:PHE:HB3	2.00	0.44
1:A:59:LEU:HB3	1:H:59:LEU:HD13	2.00	0.44
1:A:0[A]:MET:HG2	1:A:97:PRO:HG3	2.00	0.44
1:B:84:GLN:HB3	1:B:85:PRO:HD3	2.00	0.44
1:C:55:GLY:HA3	1:C:104:LEU:HD21	2.00	0.43
1:H:7:TYR:HB2	1:H:50:VAL:HG22	2.00	0.43
1:C:24:GLU:OE2	2:C:201:EDO:H12	2.17	0.43
1:E:99:SER:HG	1:F:22[A]:GLU:HG2	1.84	0.43
1:F:16:GLU:HG2	1:F:20:HIS:CD2	2.54	0.42
1:E:7:TYR:HB2	1:E:50:VAL:CG2	2.50	0.42
1:C:104:LEU:HD23	1:C:104:LEU:HA	1.79	0.41
1:F:47:LEU:HD22	2:F:201:EDO:C1	2.50	0.41
1:C:100:PHE:CD1	1:D:26:ILE:HG12	2.55	0.41
1:G:20:HIS:CG	1:G:82:ASN:HB3	2.56	0.41
1:H:89:ARG:NH2	3:H:313:HOH:O	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:LYS:HD2	3:A:318:HOH:O	2.19	0.41
1:F:91[A]:GLN:HG2	1:F:92:ASN:N	2.36	0.41
1:C:74:GLU:HG2	2:C:202:EDO:H22	2.03	0.41
1:A:40:PHE:HB3	1:B:95:LEU:HB3	2.03	0.41
1:C:7:TYR:HB2	1:C:50:VAL:HG22	2.02	0.41
1:G:40:PHE:HB3	1:H:95:LEU:HB3	2.03	0.41
1:B:68:ILE:O	1:B:72:MET:HG3	2.21	0.41
1:D:0:MET:HG2	1:D:57:GLU:HA	2.03	0.40
1:A:81:LYS:NZ	3:A:304:HOH:O	2.35	0.40
1:H:70:LEU:HD12	1:H:76:GLN:NE2	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:376:HOH:O	3:D:379:HOH:O[1_545]	2.13	0.07

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/126 (94%)	115 (97%)	4 (3%)	0	100	100
1	B	117/126 (93%)	113 (97%)	4 (3%)	0	100	100
1	C	120/126 (95%)	118 (98%)	2 (2%)	0	100	100
1	D	120/126 (95%)	118 (98%)	2 (2%)	0	100	100
1	E	121/126 (96%)	117 (97%)	4 (3%)	0	100	100
1	F	120/126 (95%)	117 (98%)	3 (2%)	0	100	100
1	G	116/126 (92%)	113 (97%)	3 (3%)	0	100	100
1	H	122/126 (97%)	118 (97%)	4 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	955/1008 (95%)	929 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/112 (93%)	101 (97%)	3 (3%)	42	16
1	B	100/112 (89%)	97 (97%)	3 (3%)	41	15
1	C	100/112 (89%)	96 (96%)	4 (4%)	31	9
1	D	104/112 (93%)	101 (97%)	3 (3%)	42	16
1	E	103/112 (92%)	101 (98%)	2 (2%)	57	34
1	F	104/112 (93%)	101 (97%)	3 (3%)	42	16
1	G	100/112 (89%)	98 (98%)	2 (2%)	55	32
1	H	107/112 (96%)	105 (98%)	2 (2%)	57	34
All	All	822/896 (92%)	800 (97%)	22 (3%)	49	19

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0[A]	MET
1	A	0[B]	MET
1	A	39	TYR
1	B	0[A]	MET
1	B	0[B]	MET
1	B	39	TYR
1	C	0	MET
1	C	1	ILE
1	C	24	GLU
1	C	39	TYR
1	D	39	TYR
1	D	91[A]	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	91[B]	GLN
1	E	-3	HIS
1	E	39	TYR
1	F	39	TYR
1	F	52	HIS
1	F	70	LEU
1	G	39	TYR
1	G	70	LEU
1	H	0	MET
1	H	39	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	F	201	-	3,3,3	0.42	0	2,2,2	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	C	201	-	3,3,3	0.47	0	2,2,2	0.25	0
2	EDO	A	201	-	3,3,3	0.43	0	2,2,2	0.43	0
2	EDO	H	201	-	3,3,3	0.38	0	2,2,2	0.47	0
2	EDO	C	202	-	3,3,3	0.49	0	2,2,2	0.27	0
2	EDO	D	201	-	3,3,3	0.49	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	F	201	-	-	0/1/1/1	-
2	EDO	C	201	-	-	0/1/1/1	-
2	EDO	A	201	-	-	0/1/1/1	-
2	EDO	H	201	-	-	0/1/1/1	-
2	EDO	C	202	-	-	0/1/1/1	-
2	EDO	D	201	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	201	EDO	3	0
2	C	201	EDO	1	0
2	A	201	EDO	1	0
2	H	201	EDO	3	0
2	C	202	EDO	3	0
2	D	201	EDO	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/126 (92%)	0.42	6 (5%) 27 26	8, 16, 45, 79	0
1	B	117/126 (92%)	0.51	7 (5%) 21 20	9, 18, 43, 53	0
1	C	118/126 (93%)	0.66	14 (11%) 4 3	10, 23, 57, 68	0
1	D	119/126 (94%)	0.45	7 (5%) 22 21	9, 18, 42, 70	0
1	E	120/126 (95%)	0.50	5 (4%) 36 35	11, 20, 47, 75	0
1	F	118/126 (93%)	0.52	7 (5%) 22 21	11, 21, 39, 44	0
1	G	117/126 (92%)	0.49	5 (4%) 35 34	10, 19, 43, 59	0
1	H	120/126 (95%)	0.51	6 (5%) 28 27	10, 19, 43, 60	0
All	All	945/1008 (93%)	0.51	57 (6%) 21 20	8, 19, 44, 79	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	114	HIS	5.9
1	E	115	ALA	5.9
1	D	114	HIS	5.6
1	C	114	HIS	5.2
1	F	114	HIS	4.9
1	G	114	HIS	4.9
1	D	115	ALA	4.9
1	D	-2	HIS	4.4
1	H	114	HIS	4.2
1	C	110	PRO	4.2
1	H	-2	HIS	4.2
1	B	104	LEU	4.0
1	G	110	PRO	4.0
1	A	-2	HIS	4.0
1	A	111	GLU	3.9
1	A	110	PRO	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	-2	HIS	3.8
1	B	113	GLU	3.7
1	G	115	ALA	3.6
1	C	113	GLU	3.4
1	G	104	LEU	3.4
1	G	-1	HIS	3.3
1	H	115	ALA	3.3
1	C	112	ASP	3.3
1	E	110	PRO	3.1
1	F	110	PRO	3.1
1	A	23	ARG	3.0
1	E	-2	HIS	3.0
1	C	104	LEU	3.0
1	B	-1	HIS	2.8
1	C	-2	HIS	2.8
1	F	107	LEU	2.8
1	D	113	GLU	2.7
1	H	110	PRO	2.7
1	F	71	ALA	2.6
1	B	110	PRO	2.6
1	D	110	PRO	2.6
1	C	59	LEU	2.5
1	C	109	GLN	2.5
1	C	73	PRO	2.4
1	F	108	TRP	2.4
1	F	-1	HIS	2.4
1	B	109	GLN	2.2
1	B	112	ASP	2.2
1	F	115	ALA	2.2
1	H	-1	HIS	2.2
1	C	63	ALA	2.2
1	C	-1	HIS	2.1
1	C	72[A]	MET	2.1
1	E	108	TRP	2.1
1	C	71	ALA	2.1
1	D	104	LEU	2.1
1	D	-3	HIS	2.1
1	C	23	ARG	2.0
1	A	-3	HIS	2.0
1	A	104	LEU	2.0
1	H	-3	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	C	202	4/4	0.69	0.28	42,43,46,47	0
2	EDO	C	201	4/4	0.70	0.27	42,42,44,45	0
2	EDO	A	201	4/4	0.84	0.19	26,29,33,37	0
2	EDO	D	201	4/4	0.84	0.23	28,32,39,42	0
2	EDO	H	201	4/4	0.93	0.26	21,26,34,44	0
2	EDO	F	201	4/4	0.95	0.15	17,30,36,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.