



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 30, 2021 – 06:13 PM EST

PDB ID : 3KCN  
Title : The crystal structure of adenylate cyclase from *Rhodopirellula baltica*  
Authors : Zhang, Z.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center  
for Structural Genomics (NYSGXRC)  
Deposited on : 2009-10-21  
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.16
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	 2% 63% 23% 11%
1	B	151	 2% 68% 19% 5% 9%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylate cyclase homolog.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	135	Total	C	N	O	S	Se	0	0	0
			1050	661	173	206	3	7			
1	B	137	Total	C	N	O	S	Se	0	0	0
			1068	674	175	209	3	7			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MSE	-	expression tag	UNP Q7UJS6
A	4	SER	-	expression tag	UNP Q7UJS6
A	5	LEU	-	expression tag	UNP Q7UJS6
A	146	GLU	-	expression tag	UNP Q7UJS6
A	147	GLY	-	expression tag	UNP Q7UJS6
A	148	HIS	-	expression tag	UNP Q7UJS6
A	149	HIS	-	expression tag	UNP Q7UJS6
A	150	HIS	-	expression tag	UNP Q7UJS6
A	151	HIS	-	expression tag	UNP Q7UJS6
A	152	HIS	-	expression tag	UNP Q7UJS6
A	153	HIS	-	expression tag	UNP Q7UJS6
B	3	MSE	-	expression tag	UNP Q7UJS6
B	4	SER	-	expression tag	UNP Q7UJS6
B	5	LEU	-	expression tag	UNP Q7UJS6
B	146	GLU	-	expression tag	UNP Q7UJS6
B	147	GLY	-	expression tag	UNP Q7UJS6
B	148	HIS	-	expression tag	UNP Q7UJS6
B	149	HIS	-	expression tag	UNP Q7UJS6
B	150	HIS	-	expression tag	UNP Q7UJS6
B	151	HIS	-	expression tag	UNP Q7UJS6
B	152	HIS	-	expression tag	UNP Q7UJS6
B	153	HIS	-	expression tag	UNP Q7UJS6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	65	Total 65	O 65	0	0
2	B	61	Total 61	O 61	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Adenylate cyclase homolog



#### • Molecule 1: Adenylate cyclase homolog



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.93Å 103.10Å 33.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.55 – 2.45 103.10 – 2.45	Depositor EDS
% Data completeness (in resolution range)	98.7 (65.55-2.45) 99.7 (103.10-2.45)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.97 (at 2.45Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.217 , 0.267 0.214 , 0.259	Depositor DCC
$R_{free}$ test set	548 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.1	Xtriage
Anisotropy	0.761	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 36.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2244	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.22 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.7865e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/1055	0.79	0/1410
1	B	0.77	0/1074	0.80	0/1436
All	All	0.77	0/2129	0.79	0/2846

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1050	0	1069	44	0
1	B	1068	0	1088	37	0
2	A	65	0	0	4	0
2	B	61	0	0	4	0
All	All	2244	0	2157	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:42:LEU:O	1:B:46:LYS:HD2	1.68	0.93

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:LEU:HD11	1:A:63:MSE:HE1	1.48	0.92
1:A:42:LEU:CD1	1:A:63:MSE:HE1	2.06	0.86
1:A:63:MSE:HE3	1:A:68:VAL:HA	1.60	0.83
1:B:65:GLY:HA3	1:B:83:MSE:HE1	1.60	0.82
1:B:58:MSE:HE1	1:B:66:THR:HG22	1.68	0.76
1:A:63:MSE:CE	1:A:68:VAL:HG22	2.16	0.75
1:B:107:ASN:O	1:B:110:CYS:HB2	1.90	0.72
1:A:58:MSE:HA	1:A:83:MSE:HE1	1.74	0.69
1:B:99:GLU:HA	2:B:210:HOH:O	1.92	0.68
1:A:66:THR:HG23	2:A:191:HOH:O	1.95	0.66
1:A:104:ARG:HH12	1:B:99:GLU:CG	2.08	0.66
1:A:19:LEU:HD13	1:A:34:THR:HB	1.79	0.65
1:B:65:GLY:CA	1:B:83:MSE:HE1	2.27	0.65
1:B:58:MSE:HE1	1:B:66:THR:CG2	2.28	0.63
1:B:25:ASN:HB3	2:B:199:HOH:O	1.99	0.62
1:A:104:ARG:HH12	1:B:99:GLU:HG3	1.65	0.61
1:B:103:PHE:O	1:B:104:ARG:HD3	2.01	0.61
1:A:38:GLY:HA3	1:A:63:MSE:HE2	1.81	0.61
1:A:63:MSE:HE3	1:A:68:VAL:HG22	1.84	0.60
1:A:134:GLU:O	1:A:138:LYS:HG2	2.03	0.57
1:B:22:LEU:HD23	1:B:112:MSE:HE1	1.87	0.57
1:A:38:GLY:CA	1:A:63:MSE:HE2	2.35	0.56
1:B:66:THR:HG23	2:B:194:HOH:O	2.06	0.54
1:A:26:LEU:HD21	1:A:112:MSE:HE1	1.91	0.53
1:A:105:PHE:CE2	1:B:94:MSE:HE1	2.44	0.53
1:A:97:VAL:HG12	1:B:104:ARG:HD2	1.90	0.53
1:A:112:MSE:HE3	1:A:112:MSE:O	2.09	0.53
1:B:84:LEU:HG	1:B:108:LYS:HG2	1.90	0.53
1:A:104:ARG:HH12	1:B:99:GLU:HG2	1.75	0.52
1:A:58:MSE:HA	1:A:83:MSE:CE	2.38	0.52
1:B:65:GLY:HA3	1:B:83:MSE:CE	2.34	0.52
1:A:55:MSE:HE1	1:A:84:LEU:HD11	1.92	0.52
1:A:104:ARG:NH1	1:B:99:GLU:CG	2.73	0.51
1:B:131:SER:O	1:B:135:LEU:HD13	2.11	0.51
1:A:96:ALA:HB1	1:A:102:VAL:HG23	1.93	0.51
1:A:96:ALA:HB1	1:A:102:VAL:CG2	2.41	0.51
1:A:63:MSE:HE2	1:A:68:VAL:HG22	1.93	0.50
1:A:59:ARG:HB3	2:A:168:HOH:O	2.10	0.50
1:B:55:MSE:HE1	1:B:119:ILE:HD11	1.93	0.49
1:A:135:LEU:CD2	1:B:135:LEU:HD11	2.43	0.48
1:A:63:MSE:HG2	1:A:68:VAL:HG22	1.96	0.48

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:LEU:HD11	1:A:94:MSE:HE1	1.96	0.48
1:B:112:MSE:HG3	1:B:116:LYS:HE3	1.94	0.48
1:A:63:MSE:HE3	1:A:68:VAL:CA	2.38	0.48
1:A:104:ARG:NH1	1:B:99:GLU:HG2	2.28	0.47
1:B:15:ASP:C	1:B:15:ASP:OD1	2.53	0.47
1:A:94:MSE:HB2	1:A:94:MSE:HE2	1.60	0.47
1:B:88:GLN:HG2	1:B:89:ASP:N	2.30	0.47
1:A:59:ARG:NH2	1:A:108:LYS:HE3	2.30	0.47
1:A:58:MSE:HE3	1:A:89:ASP:HB3	1.97	0.46
1:B:65:GLY:C	1:B:83:MSE:HE1	2.35	0.46
1:A:63:MSE:HG2	1:A:68:VAL:CG2	2.45	0.46
1:A:75:ILE:HD11	2:A:205:HOH:O	2.16	0.46
1:A:104:ARG:HG2	1:A:104:ARG:HH11	1.81	0.45
1:A:108:LYS:HB3	1:A:109:PRO:HA	1.97	0.45
1:B:58:MSE:HE3	1:B:93:ALA:CB	2.47	0.45
1:B:66:THR:O	1:B:70:GLN:HG2	2.16	0.45
1:A:104:ARG:NH1	1:B:99:GLU:HG3	2.30	0.45
1:B:58:MSE:CE	1:B:66:THR:HG22	2.44	0.43
1:B:85:THR:OG1	1:B:89:ASP:HB2	2.18	0.43
1:A:135:LEU:HD22	1:B:135:LEU:HD11	2.00	0.43
1:A:55:MSE:CE	1:A:84:LEU:HD11	2.49	0.43
1:B:63:MSE:SE	1:B:68:VAL:HG22	2.68	0.42
1:A:55:MSE:HB3	1:A:82:LEU:HB2	2.01	0.42
1:A:64:GLU:HB2	2:A:191:HOH:O	2.18	0.42
1:A:127:ASP:O	1:A:131:SER:HB3	2.19	0.42
1:B:77:PRO:HD2	2:B:168:HOH:O	2.19	0.42
1:B:19:LEU:HA	1:B:19:LEU:HD23	1.84	0.42
1:A:59:ARG:HH21	1:A:108:LYS:HE3	1.85	0.42
1:B:42:LEU:O	1:B:46:LYS:CD	2.55	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/151 (88%)	131 (98%)	1 (1%)	1 (1%)	19	22
1	B	135/151 (89%)	135 (100%)	0	0	100	100
All	All	268/302 (89%)	266 (99%)	1 (0%)	1 (0%)	34	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	87	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/125 (96%)	108 (90%)	12 (10%)	7	7
1	B	122/125 (98%)	112 (92%)	10 (8%)	11	13
All	All	242/250 (97%)	220 (91%)	22 (9%)	9	10

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	SER
1	A	18	LEU
1	A	25	ASN
1	A	47	LYS
1	A	82	LEU
1	A	87	ASN
1	A	104	ARG
1	A	113	SER
1	A	128	LEU
1	A	135	LEU
1	A	136	LEU
1	A	138	LYS
1	B	15	ASP
1	B	46	LYS
1	B	58	MSE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	66	THR
1	B	83	MSE
1	B	91	THR
1	B	99	GLU
1	B	104	ARG
1	B	128	LEU
1	B	137	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	107	ASN
1	B	20	ASN
1	B	78	ASN
1	B	111	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	128/151 (84%)	-0.17	3 (2%) 60 56	8, 16, 34, 48	0
1	B	130/151 (86%)	-0.17	3 (2%) 60 56	8, 15, 38, 46	0
All	All	258/302 (85%)	-0.17	6 (2%) 60 56	8, 16, 37, 48	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	LYS	4.5
1	B	110	CYS	3.1
1	A	62	GLY	2.6
1	B	137	LYS	2.5
1	B	97	VAL	2.4
1	A	87	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.