



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 01:06 am BST

PDB ID : 5KC7
Title : Crystal structure of Cbln1 (Val55-Gly58 deletion mutant)
Authors : Elegheert, J.; Aricescu, A.R.
Deposited on : 2016-06-05
Resolution : 7.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 7.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1004 (10.00-3.90)
Clashscore	141614	1069 (10.00-3.90)
Ramachandran outliers	138981	1002 (10.00-3.90)
Sidechain outliers	138945	1002 (10.00-3.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	178	
1	B	178	
1	C	178	
1	D	178	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4276 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cerebellin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	135	Total	C	N	O	S	0	0	0
			1073	686	186	196	5			
1	B	135	Total	C	N	O	S	0	0	0
			1063	681	183	194	5			
1	C	135	Total	C	N	O	S	0	0	0
			1073	686	186	196	5			
1	D	135	Total	C	N	O	S	0	0	0
			1067	683	183	196	5			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	GLU	-	expression tag	UNP P23435
A	26	THR	-	expression tag	UNP P23435
A	27	GLY	-	expression tag	UNP P23435
A	?	-	VAL	deletion	UNP P23435
A	?	-	ARG	deletion	UNP P23435
A	?	-	SER	deletion	UNP P23435
A	?	-	GLY	deletion	UNP P23435
A	194	GLY	-	expression tag	UNP P23435
A	195	THR	-	expression tag	UNP P23435
A	196	LYS	-	expression tag	UNP P23435
A	197	HIS	-	expression tag	UNP P23435
A	198	HIS	-	expression tag	UNP P23435
A	199	HIS	-	expression tag	UNP P23435
A	200	HIS	-	expression tag	UNP P23435
A	201	HIS	-	expression tag	UNP P23435
A	202	HIS	-	expression tag	UNP P23435
B	25	GLU	-	expression tag	UNP P23435
B	26	THR	-	expression tag	UNP P23435
B	27	GLY	-	expression tag	UNP P23435
B	?	-	VAL	deletion	UNP P23435
B	?	-	ARG	deletion	UNP P23435

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	SER	deletion	UNP P23435
B	?	-	GLY	deletion	UNP P23435
B	194	GLY	-	expression tag	UNP P23435
B	195	THR	-	expression tag	UNP P23435
B	196	LYS	-	expression tag	UNP P23435
B	197	HIS	-	expression tag	UNP P23435
B	198	HIS	-	expression tag	UNP P23435
B	199	HIS	-	expression tag	UNP P23435
B	200	HIS	-	expression tag	UNP P23435
B	201	HIS	-	expression tag	UNP P23435
B	202	HIS	-	expression tag	UNP P23435
C	25	GLU	-	expression tag	UNP P23435
C	26	THR	-	expression tag	UNP P23435
C	27	GLY	-	expression tag	UNP P23435
C	?	-	VAL	deletion	UNP P23435
C	?	-	ARG	deletion	UNP P23435
C	?	-	SER	deletion	UNP P23435
C	?	-	GLY	deletion	UNP P23435
C	194	GLY	-	expression tag	UNP P23435
C	195	THR	-	expression tag	UNP P23435
C	196	LYS	-	expression tag	UNP P23435
C	197	HIS	-	expression tag	UNP P23435
C	198	HIS	-	expression tag	UNP P23435
C	199	HIS	-	expression tag	UNP P23435
C	200	HIS	-	expression tag	UNP P23435
C	201	HIS	-	expression tag	UNP P23435
C	202	HIS	-	expression tag	UNP P23435
D	25	GLU	-	expression tag	UNP P23435
D	26	THR	-	expression tag	UNP P23435
D	27	GLY	-	expression tag	UNP P23435
D	?	-	VAL	deletion	UNP P23435
D	?	-	ARG	deletion	UNP P23435
D	?	-	SER	deletion	UNP P23435
D	?	-	GLY	deletion	UNP P23435
D	194	GLY	-	expression tag	UNP P23435
D	195	THR	-	expression tag	UNP P23435
D	196	LYS	-	expression tag	UNP P23435
D	197	HIS	-	expression tag	UNP P23435
D	198	HIS	-	expression tag	UNP P23435
D	199	HIS	-	expression tag	UNP P23435
D	200	HIS	-	expression tag	UNP P23435
D	201	HIS	-	expression tag	UNP P23435

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	202	HIS	-	expression tag	UNP P23435

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cerebellin-1

Chain A:  71% 24%




• Molecule 1: Cerebellin-1

Chain B:  72% 24%




• Molecule 1: Cerebellin-1

Chain C:  72% 24%



• Molecule 1: Cerebellin-1

Chain D:  72% 24%



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	187.45Å 187.45Å 187.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.10 – 7.04 76.53 – 7.03	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.10-7.04) 100.0 (76.53-7.03)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.96 (at 6.72Å)	Xtriage
Refinement program	PHENIX (dev_2283: ???)	Depositor
R, R_{free}	0.270 , 0.351 0.270 , 0.647	Depositor DCC
R_{free} test set	173 reflections (9.61%)	wwPDB-VP
Wilson B-factor (Å ²)	402.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 612.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.047 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.81	EDS
Total number of atoms	4276	wwPDB-VP
Average B, all atoms (Å ²)	469.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/1097	0.42	0/1481
1	B	0.24	0/1086	0.42	0/1466
1	C	0.24	0/1097	0.42	0/1481
1	D	0.24	0/1091	0.42	0/1474
All	All	0.24	0/4371	0.42	0/5902

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1073	0	1058	4	4
1	B	1063	0	1044	4	1
1	C	1073	0	1058	3	1
1	D	1067	0	1047	3	4
All	All	4276	0	4207	11	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:THR:CB	1:D:172:GLU:O	1.95	1.14
1:C:120:LYS:NZ	1:C:125:GLN:O	2.11	0.84
1:D:120:LYS:NZ	1:D:125:GLN:O	2.11	0.84
1:A:120:LYS:NZ	1:A:125:GLN:O	2.11	0.84
1:B:120:LYS:NZ	1:B:125:GLN:O	2.11	0.83
1:B:146:GLN:NE2	1:B:151:GLU:OE2	2.43	0.49
1:A:146:GLN:NE2	1:A:151:GLU:OE2	2.43	0.48
1:D:146:GLN:NE2	1:D:151:GLU:OE2	2.43	0.45
1:A:117:HIS:CE1	1:C:155:ASN:HA	2.56	0.41
1:A:155:ASN:HA	1:B:117:HIS:CE1	2.56	0.41
1:C:146:GLN:NE2	1:C:151:GLU:OE2	2.43	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:GLU:O	1:C:70:THR:OG1[14_545]	1.45	0.75
1:A:173:ARG:CA	1:D:72:HIS:CE1[12_455]	1.63	0.57
1:A:172:GLU:O	1:D:70:THR:OG1[12_455]	1.88	0.32
1:A:173:ARG:N	1:D:72:HIS:CE1[12_455]	2.02	0.18
1:A:173:ARG:CA	1:D:72:HIS:ND1[12_455]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/178 (75%)	130 (98%)	3 (2%)	0	100	100
1	B	133/178 (75%)	130 (98%)	3 (2%)	0	100	100
1	C	133/178 (75%)	130 (98%)	3 (2%)	0	100	100
1	D	133/178 (75%)	130 (98%)	3 (2%)	0	100	100
All	All	532/712 (75%)	520 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/153 (76%)	115 (99%)	1 (1%)	78	87
1	B	113/153 (74%)	112 (99%)	1 (1%)	78	87
1	C	116/153 (76%)	115 (99%)	1 (1%)	78	87
1	D	115/153 (75%)	114 (99%)	1 (1%)	78	87
All	All	460/612 (75%)	456 (99%)	4 (1%)	78	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	GLN
1	B	146	GLN
1	C	146	GLN
1	D	146	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.