



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2020 – 10:07 AM EDT

PDB ID : 7KCP
Title : Crystal structure of Acetyl-CoA Synthetase in Complex with Adenosine-5'-pyrophosphate from *Coccidioides posadasii* C735
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2020-10-07
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.14.6
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	706	<div> <div>7%</div> <div>77%</div> <div>7%</div> <div>16%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	710	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5046 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetyl-coenzyme A synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	594	Total	C	N	O	S	0	7	0
			4604	2960	790	836	18			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	MET	-	initiating methionine	UNP C5PGB4
A	-14	HIS	-	expression tag	UNP C5PGB4
A	-13	HIS	-	expression tag	UNP C5PGB4
A	-12	HIS	-	expression tag	UNP C5PGB4
A	-11	HIS	-	expression tag	UNP C5PGB4
A	-10	HIS	-	expression tag	UNP C5PGB4
A	-9	HIS	-	expression tag	UNP C5PGB4
A	-8	HIS	-	expression tag	UNP C5PGB4
A	-7	HIS	-	expression tag	UNP C5PGB4
A	-6	GLU	-	expression tag	UNP C5PGB4
A	-5	ASN	-	expression tag	UNP C5PGB4
A	-4	LEU	-	expression tag	UNP C5PGB4
A	-3	TYR	-	expression tag	UNP C5PGB4
A	-2	PHE	-	expression tag	UNP C5PGB4
A	-1	GLN	-	expression tag	UNP C5PGB4
A	0	GLY	-	expression tag	UNP C5PGB4

- Molecule 2 is ADENOSINE-5'-MONOPHOSPHATE-PROPYL ESTER (three-letter code: PRX) (formula: C₁₃H₂₀N₅O₇P) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			26	13	5	7	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	366	Total O 371 371	0	5

- Molecule 1: Acetyl-coenzyme A synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	107.17Å 107.17Å 116.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.26 – 2.15 49.24 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (36.26-2.15) 100.0 (49.24-2.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.84 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.154 , 0.192 0.154 , 0.192	Depositor DCC
R_{free} test set	1989 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 59.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.045 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5046	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO, PRX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/4756	0.53	0/6483

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4604	0	4415	32	0
2	A	26	0	19	1	0
3	A	1	0	0	0	0
4	A	44	0	66	7	0
5	A	371	0	0	3	0
All	All	5046	0	4500	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:363:GLU:OE2	5:A:801:HOH:O	2.08	0.72
1:A:265:PRO:O	4:A:710:EDO:H11	1.91	0.71
1:A:12:VAL:HG21	1:A:522:ASP:HA	1.81	0.63
1:A:267:GLU:HG2	4:A:710:EDO:C2	2.33	0.58
1:A:471:GLU:OE2	1:A:484:ARG:NH1	2.39	0.56
1:A:280:THR:HG21	1:A:431:THR:HG21	1.88	0.55
1:A:523:HIS:CE1	1:A:524:GLU:HG3	2.42	0.55
1:A:429:TRP:CE2	2:A:701:PRX:H3P1	2.44	0.52
1:A:12:VAL:HG22	1:A:476:ASP:OD1	2.09	0.52
1:A:580:PHE:HA	1:A:615:PHE:HB2	1.92	0.51
1:A:372:VAL:HG22	1:A:373:ALA:H	1.77	0.49
1:A:395:LEU:HD21	1:A:398:LEU:HD21	1.95	0.49
1:A:471:GLU:HG3	4:A:712:EDO:H22	1.95	0.48
1:A:143[B]:LYS:HE2	1:A:264:LEU:HD11	1.96	0.48
1:A:350:PRO:HB3	1:A:358:TYR:CE2	2.48	0.48
1:A:267:GLU:HG2	4:A:710:EDO:H22	1.96	0.47
1:A:485:LYS:HE2	5:A:1008:HOH:O	2.14	0.47
1:A:485:LYS:NZ	5:A:811:HOH:O	2.38	0.46
1:A:288:LYS:HG2	1:A:500:ARG:CZ	2.45	0.46
1:A:115:TYR:CD1	1:A:161:PRO:HD3	2.52	0.44
1:A:280:THR:CG2	1:A:431:THR:HG21	2.49	0.43
1:A:61:MET:HE2	4:A:709:EDO:H11	2.00	0.43
1:A:545:SER:HB3	1:A:548:GLU:HB2	2.01	0.42
1:A:264:LEU:HD21	4:A:710:EDO:H12	2.00	0.42
1:A:425[B]:VAL:HG13	1:A:440:PRO:HG2	2.03	0.41
1:A:452:ALA:O	1:A:453:SER:HB2	2.20	0.41
1:A:61:MET:HB2	1:A:61:MET:HE2	1.96	0.41
1:A:26:VAL:HG22	1:A:30:ASN:ND2	2.35	0.41
1:A:65:LEU:HA	4:A:704:EDO:H11	2.03	0.41
1:A:461:PRO:HG2	1:A:527:TYR:OH	2.21	0.41
1:A:115:TYR:CD2	1:A:123:GLY:HA2	2.56	0.41
1:A:374:PRO:HG2	1:A:403:GLU:HG3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	595/706 (84%)	573 (96%)	21 (4%)	1 (0%)	47	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	453	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	468/584 (80%)	463 (99%)	5 (1%)	73	78

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	ASP
1	A	316	ARG
1	A	413	TYR
1	A	430	GLN
1	A	592	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	523	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	708	-	3,3,3	0.53	0	2,2,2	0.19	0
4	EDO	A	703	-	3,3,3	0.53	0	2,2,2	0.45	0
4	EDO	A	709	-	3,3,3	0.55	0	2,2,2	0.09	0
4	EDO	A	707	-	3,3,3	0.60	0	2,2,2	0.11	0
4	EDO	A	710	-	3,3,3	0.46	0	2,2,2	0.09	0
4	EDO	A	713	-	3,3,3	0.46	0	2,2,2	0.43	0
4	EDO	A	712	-	3,3,3	0.57	0	2,2,2	0.21	0
4	EDO	A	704	-	3,3,3	0.49	0	2,2,2	0.27	0
4	EDO	A	705	-	3,3,3	0.53	0	2,2,2	0.61	0
4	EDO	A	711	-	3,3,3	0.52	0	2,2,2	0.27	0
4	EDO	A	706	-	3,3,3	0.37	0	2,2,2	0.73	0
2	PRX	A	701	-	25,28,28	0.63	0	27,41,41	0.71	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	708	-	-	1/1/1/1	-
4	EDO	A	703	-	-	0/1/1/1	-
4	EDO	A	709	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	707	-	-	0/1/1/1	-
4	EDO	A	710	-	-	0/1/1/1	-
4	EDO	A	713	-	-	0/1/1/1	-
4	EDO	A	712	-	-	1/1/1/1	-
4	EDO	A	704	-	-	0/1/1/1	-
4	EDO	A	705	-	-	1/1/1/1	-
4	EDO	A	711	-	-	1/1/1/1	-
4	EDO	A	706	-	-	0/1/1/1	-
2	PRX	A	701	-	-	0/11/31/31	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	PRX	C5-C6-N6	2.20	123.69	120.35

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	712	EDO	O1-C1-C2-O2
4	A	708	EDO	O1-C1-C2-O2
4	A	705	EDO	O1-C1-C2-O2
4	A	711	EDO	O1-C1-C2-O2


There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	709	EDO	1	0
4	A	710	EDO	4	0
4	A	712	EDO	1	0
4	A	704	EDO	1	0
2	A	701	PRX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	594/706 (84%)	0.15	50 (8%) 11 15	17, 37, 101, 128	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	580	PHE	9.3
1	A	581	VAL	7.6
1	A	594	ARG	6.6
1	A	614	VAL	6.6
1	A	560	ALA	6.5
1	A	613	VAL	6.5
1	A	615	PHE	6.2
1	A	554	ILE	5.4
1	A	579	ALA	5.2
1	A	562	ALA	5.1
1	A	565	VAL	5.1
1	A	616	VAL	4.9
1	A	283	SER	4.8
1	A	611	PRO	4.5
1	A	610	ALA	4.5
1	A	467	VAL	4.4
1	A	559	VAL	4.4
1	A	558	CYS	4.3
1	A	593	LEU	4.0
1	A	578	HIS	3.9
1	A	583	LEU	3.7
1	A	576	ALA	3.6
1	A	582	ALA	3.5
1	A	590	ARG	3.4
1	A	563	ALA	3.3
1	A	589	ASN	3.3
1	A	609	ALA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	536	VAL	3.2
1	A	546	THR	3.1
1	A	564	VAL	3.1
1	A	284	THR	3.0
1	A	534	ASP	2.9
1	A	533	VAL	2.9
1	A	608	PHE	2.9
1	A	591	GLU	2.8
1	A	547	ALA	2.6
1	A	535	ASP	2.6
1	A	537	VAL	2.5
1	A	382	ARG	2.5
1	A	541	GLY	2.4
1	A	561	GLU	2.4
1	A	592	GLN	2.4
1	A	551	ALA	2.3
1	A	595	LYS	2.3
1	A	550	GLU	2.3
1	A	538	ASN	2.1
1	A	468	THR	2.1
1	A	557	HIS	2.1
1	A	577	VAL	2.1
1	A	612	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	A	709	4/4	0.67	0.18	45,47,54,56	0
4	EDO	A	712	4/4	0.72	0.38	60,61,64,65	0
4	EDO	A	703	4/4	0.79	0.12	40,44,50,55	0
4	EDO	A	710	4/4	0.85	0.19	32,36,41,41	0
4	EDO	A	707	4/4	0.87	0.14	39,42,43,46	0
4	EDO	A	704	4/4	0.91	0.18	38,41,42,43	0
4	EDO	A	705	4/4	0.91	0.16	32,33,36,38	0
4	EDO	A	713	4/4	0.94	0.24	49,53,53,55	0
3	MG	A	702	1/1	0.95	0.09	39,39,39,39	0
4	EDO	A	711	4/4	0.95	0.15	26,35,35,45	0
4	EDO	A	708	4/4	0.96	0.13	44,45,49,58	0
4	EDO	A	706	4/4	0.96	0.10	34,35,35,36	0
2	PRX	A	701	26/26	0.97	0.10	28,34,38,39	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.