



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 18, 2021 – 12:37 PM EST

PDB ID : 7KEV  
Title : PCSK9 in complex with a cyclic peptide LDLR disruptor  
Authors : Spraggon, G.; Chopra, R.  
Deposited on : 2020-10-12  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	124	
2	B	546	
3	C	14	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proprotein convertase subtilisin/kexin type 9 Propeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	92	Total	C	N	O	S	0	1	0
			748	479	136	131	2			

- Molecule 2 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	471	Total	C	N	O	S	0	0	0
			3497	2154	645	667	31			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	474	ILE	VAL	variant	UNP Q8NBP7
B	670	GLU	GLY	variant	UNP Q8NBP7
B	693	HIS	-	expression tag	UNP Q8NBP7
B	694	HIS	-	expression tag	UNP Q8NBP7
B	695	HIS	-	expression tag	UNP Q8NBP7
B	696	HIS	-	expression tag	UNP Q8NBP7
B	697	HIS	-	expression tag	UNP Q8NBP7
B	698	HIS	-	expression tag	UNP Q8NBP7

- Molecule 3 is a protein called cyclic peptide LDLR disruptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	14	Total	C	N	O	S	0	0	1
			124	85	18	20	1			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Ca 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total 5	O 5	0	0
5	B	12	Total 12	O 12	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proprotein convertase subtilisin/kexin type 9 Propeptide



- Molecule 2: Proprotein convertase subtilisin/kexin type 9



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.23Å 70.44Å 149.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.64 – 2.80 40.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	93.9 (40.64-2.80) 85.6 (40.64-2.80)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.33 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, $R_{free}$	0.226 , 0.280 0.225 , 0.279	Depositor DCC
$R_{free}$ test set	785 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.7	Xtriage
Anisotropy	0.615	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 38.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4387	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: WCM, CA, ALO, ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/768	0.60	0/1037
2	B	0.40	1/3554 (0.0%)	0.56	2/4820 (0.0%)
3	C	1.48	0/68	1.12	0/90
All	All	0.43	1/4390 (0.0%)	0.58	2/5947 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	612	GLU	CG-CD	6.23	1.61	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	167	ARG	NE-CZ-NH2	-10.29	115.15	120.30
2	B	167	ARG	NE-CZ-NH1	6.39	123.49	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	748	0	763	4	0
2	B	3497	0	3418	66	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	124	0	69	3	0
4	B	1	0	0	0	0
5	A	5	0	0	1	0
5	B	12	0	0	0	0
All	All	4387	0	4250	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:494:LYS:HG3	2:B:517:GLY:HA2	1.64	0.80
3:C:10:ARG:HG3	3:C:11:PRO:HD2	1.65	0.76
2:B:641:THR:HG1	2:B:671:ALA:N	1.87	0.72
2:B:558:VAL:HG21	2:B:606:LEU:HD23	1.76	0.67
2:B:375:CYS:HG	2:B:378:CYS:HG	0.69	0.65
2:B:302:GLN:HB2	2:B:331:PRO:HG2	1.79	0.65
2:B:310:VAL:HG22	2:B:408:LEU:HD13	1.80	0.63
2:B:641:THR:OG1	2:B:671:ALA:N	2.31	0.63
2:B:199:ARG:NH2	2:B:235:SER:O	2.33	0.62
2:B:589:VAL:HG13	2:B:638:LEU:HD21	1.81	0.62
2:B:283:LEU:HB3	2:B:309:VAL:HG13	1.82	0.62
2:B:275:GLN:HE21	2:B:283:LEU:HD22	1.64	0.60
2:B:468:THR:OG1	2:B:471:ALA:HB2	2.04	0.58
2:B:619:GLN:N	2:B:619:GLN:HE21	2.02	0.57
2:B:469:ARG:O	2:B:496:ARG:NH1	2.37	0.57
2:B:490:SER:HB2	2:B:520:VAL:HG12	1.86	0.57
2:B:372:SER:HB2	2:B:380:VAL:HB	1.85	0.56
2:B:306:ARG:HH12	2:B:479:PRO:HG2	1.70	0.56
2:B:375:CYS:CB	2:B:378:CYS:SG	2.95	0.55
2:B:391:HIS:O	2:B:395:ILE:HG13	2.08	0.54
2:B:186:ASP:N	2:B:186:ASP:OD1	2.40	0.54
2:B:645:LEU:HD23	2:B:657:ARG:O	2.09	0.53
2:B:275:GLN:NE2	2:B:283:LEU:HD22	2.24	0.53
2:B:638:LEU:HB2	2:B:673:THR:HB	1.92	0.51
2:B:196:ILE:HB	2:B:200:VAL:HG21	1.91	0.51
2:B:375:CYS:HB2	2:B:378:CYS:SG	2.50	0.51
2:B:465:SER:HB2	2:B:473:ALA:HB2	1.92	0.51
2:B:619:GLN:N	2:B:619:GLN:NE2	2.59	0.51
2:B:614:GLY:HA2	2:B:673:THR:HA	1.93	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:199:ARG:HH22	2:B:237:ARG:HG3	1.77	0.50
2:B:503:GLN:OE1	2:B:510:ARG:NH2	2.44	0.50
2:B:490:SER:O	2:B:645:LEU:HD12	2.12	0.50
2:B:186:ASP:HB2	2:B:253:LEU:HD12	1.94	0.49
2:B:478:ALA:HB3	2:B:481:GLU:HB2	1.94	0.49
2:B:334:ILE:HD11	2:B:408:LEU:HD11	1.94	0.49
2:B:323:CYS:HA	2:B:329:SER:HB3	1.95	0.48
2:B:319:ARG:HG3	2:B:352:GLY:HA3	1.95	0.48
3:C:10:ARG:HG3	3:C:11:PRO:CD	2.38	0.48
2:B:410:GLU:HA	2:B:528:LEU:HD11	1.96	0.48
2:B:554:GLN:OE1	2:B:554:GLN:N	2.44	0.48
1:A:82:LEU:HD23	1:A:140:VAL:HA	1.97	0.47
2:B:331:PRO:O	2:B:332:GLU:HB2	2.13	0.47
2:B:208:VAL:HG21	2:B:251:ARG:NH1	2.30	0.47
3:C:3:VAL:HG23	3:C:5:THR:H	1.79	0.47
2:B:272:ARG:NH1	2:B:275:GLN:OE1	2.49	0.46
2:B:613:HIS:NE2	2:B:615:ILE:HD11	2.30	0.46
2:B:628:GLU:OE1	2:B:628:GLU:N	2.46	0.46
2:B:644:VAL:HG23	2:B:657:ARG:O	2.16	0.46
2:B:499:ARG:HD2	2:B:501:GLU:OE2	2.15	0.46
2:B:153:SER:O	2:B:153:SER:OG	2.29	0.44
2:B:645:LEU:HD23	2:B:645:LEU:H	1.81	0.44
1:A:151:ALA:HB2	2:B:253:LEU:HD13	2.00	0.44
2:B:658:SER:OG	2:B:672:VAL:HG21	2.17	0.44
2:B:229:HIS:O	2:B:233:VAL:HG23	2.18	0.44
2:B:565:HIS:CE1	2:B:597:HIS:HE2	2.36	0.43
2:B:532:ALA:HA	2:B:602:HIS:O	2.18	0.43
2:B:638:LEU:HA	2:B:638:LEU:HD23	1.79	0.43
2:B:620:GLU:OE1	2:B:621:GLN:HB2	2.18	0.43
2:B:415:LEU:HD23	2:B:415:LEU:HA	1.90	0.43
2:B:465:SER:OG	2:B:471:ALA:HB1	2.19	0.42
2:B:157:ASN:HB2	2:B:391:HIS:NE2	2.34	0.42
2:B:194:ARG:HA	2:B:194:ARG:HD3	1.86	0.42
2:B:343:ASP:O	2:B:424:ILE:HA	2.20	0.42
2:B:589:VAL:HG11	2:B:638:LEU:HD11	2.02	0.42
2:B:607:GLU:O	2:B:679:CYS:HA	2.20	0.41
2:B:455:LEU:HD13	2:B:529:LEU:HD13	2.01	0.41
1:A:108:LEU:HD12	1:A:108:LEU:HA	1.80	0.41
2:B:306:ARG:HG2	2:B:332:GLU:OE2	2.21	0.41
1:A:132:GLU:HG3	5:A:204:HOH:O	2.21	0.40
2:B:185:LEU:HD23	2:B:252:VAL:HG11	2.04	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:648:TYR:OH	2:B:657:ARG:NH1	2.48	0.40
2:B:230:LEU:HD22	2:B:390:ALA:HB2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	91/124 (73%)	86 (94%)	4 (4%)	1 (1%)	14	41
2	B	451/546 (83%)	428 (95%)	19 (4%)	4 (1%)	17	46
3	C	8/14 (57%)	6 (75%)	2 (25%)	0	100	100
All	All	550/684 (80%)	520 (94%)	25 (4%)	5 (1%)	17	46

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	84	GLU
2	B	186	ASP
2	B	585	PRO
2	B	280	VAL
2	B	604	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/104 (77%)	80 (100%)	0	100	100
2	B	379/437 (87%)	369 (97%)	10 (3%)	46	79
3	C	8/8 (100%)	6 (75%)	2 (25%)	0	2
All	All	467/549 (85%)	455 (97%)	12 (3%)	46	79

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	167	ARG
2	B	249	SER
2	B	344	GLN
2	B	485	SER
2	B	499	ARG
2	B	533	ASN
2	B	619	GLN
2	B	658	SER
2	B	659	ARG
2	B	682	ARG
3	C	4	SER
3	C	11	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	344	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ALO	C	6	3	5,6,7	1.50	0	6,7,9	3.19	3 (50%)
3	WCM	C	7	3	23,24,25	1.71	7 (30%)	30,32,34	1.50	5 (16%)
3	WCM	C	8	3	23,24,25	1.57	4 (17%)	30,32,34	2.28	14 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ALO	C	6	3	-	2/5/6/8	-
3	WCM	C	7	3	-	0/20/22/24	0/2/2/2
3	WCM	C	8	3	-	2/20/22/24	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	8	WCM	C11-N6	4.92	1.46	1.35
3	C	7	WCM	C11-N6	4.54	1.45	1.35
3	C	7	WCM	C72-C69	2.75	1.55	1.49
3	C	7	WCM	C68-C67	2.51	1.43	1.38
3	C	7	WCM	C71-C70	2.44	1.43	1.38
3	C	7	WCM	O12-C11	-2.34	1.18	1.22
3	C	7	WCM	C73-C72	2.33	1.44	1.39
3	C	8	WCM	O12-C11	-2.10	1.18	1.22
3	C	8	WCM	O11-C14	2.06	1.28	1.19
3	C	7	WCM	C71-C66	2.04	1.43	1.38
3	C	8	WCM	C71-C70	2.01	1.42	1.38

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	6	ALO	CG2-CB-CA	6.20	126.68	112.14
3	C	8	WCM	C67-C68-C69	-4.13	115.19	121.13
3	C	7	WCM	C13-C12-N6	4.09	116.29	110.64
3	C	8	WCM	C76-C77-C72	-3.92	115.63	120.56
3	C	8	WCM	C13-C12-N6	-3.86	105.30	110.64
3	C	8	WCM	C70-C69-C68	3.70	124.96	117.59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	8	WCM	C68-C69-C72	-3.65	115.03	121.36
3	C	6	ALO	CB-CA-C	-3.35	106.42	111.77
3	C	8	WCM	C77-C72-C73	3.28	124.12	117.59
3	C	6	ALO	OG1-CB-CA	-3.14	102.40	109.06
3	C	7	WCM	C65-C10-C11	3.03	116.53	109.27
3	C	8	WCM	C73-C72-C69	-2.95	116.25	121.36
3	C	8	WCM	C10-C11-N6	-2.92	113.27	119.39
3	C	8	WCM	C65-C66-C67	-2.82	115.31	120.91
3	C	7	WCM	C64-N6-C12	2.64	121.34	116.17
3	C	7	WCM	C71-C66-C67	-2.52	114.21	118.17
3	C	7	WCM	O12-C11-N6	-2.41	117.67	121.71
3	C	8	WCM	C12-N6-C11	2.33	125.92	117.64
3	C	8	WCM	C71-C70-C69	-2.27	117.87	121.13
3	C	8	WCM	C75-C76-C77	2.24	123.60	120.19
3	C	8	WCM	C71-C66-C67	2.16	121.56	118.17
3	C	8	WCM	C70-C71-C66	-2.11	118.12	121.03

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	8	WCM	C10-C65-C66-C67
3	C	8	WCM	C10-C65-C66-C71
3	C	6	ALO	N-CA-CB-CG2
3	C	6	ALO	C-CA-CB-CG2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	92/124 (74%)	-0.06	2 (2%) 62 52	40, 44, 54, 64	0
2	B	471/546 (86%)	0.27	32 (6%) 17 10	39, 54, 82, 103	0
3	C	9/14 (64%)	0.50	1 (11%) 5 3	48, 54, 68, 73	0
All	All	572/684 (83%)	0.22	35 (6%) 21 13	39, 50, 80, 103	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	640	GLY	4.4
2	B	602	HIS	4.1
2	B	531	GLN	4.0
2	B	641	THR	3.9
2	B	615	ILE	3.8
2	B	533	ASN	3.7
2	B	212	ASP	3.5
2	B	590	GLY	3.4
2	B	555	GLN	3.3
2	B	568	VAL	3.2
2	B	504	GLY	3.1
2	B	470	MET	3.1
2	B	204	ASP	3.0
2	B	467	PRO	2.9
2	B	192	ASP	2.8
2	B	659	ARG	2.8
2	B	194	ARG	2.8
2	B	530	PRO	2.7
2	B	496	ARG	2.7
2	B	620	GLU	2.6
2	B	505	GLY	2.5
2	B	682	ARG	2.5
2	B	220	ALA	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	671	ALA	2.4
2	B	604	PRO	2.4
2	B	556	GLY	2.4
2	B	553	HIS	2.4
1	A	109	THR	2.3
3	C	13	GLY	2.2
2	B	211	GLU	2.1
2	B	593	GLU	2.1
2	B	672	VAL	2.1
2	B	469	ARG	2.1
1	A	108	LEU	2.1
2	B	503	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	WCM	C	8	23/24	0.92	0.24	42,51,56,60	0
3	WCM	C	7	23/24	0.93	0.29	39,45,51,54	0
3	ALO	C	6	7/8	0.94	0.19	45,48,50,51	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CA	B	801	1/1	0.60	0.22	79,79,79,79	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers



as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers ⓘ

There are no such residues in this entry.