



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 04:11 pm BST

PDB ID : 5KIW
Title : p97 ND1-L198W in complex with VIMP
Authors : Tang, W.K.; Xia, D.
Deposited on : 2016-06-17
Resolution : 3.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1486 (3.50-3.34)
Clashscore	141614	1572 (3.50-3.34)
Ramachandran outliers	138981	1534 (3.50-3.34)
Sidechain outliers	138945	1535 (3.50-3.34)
RSRZ outliers	127900	1395 (3.50-3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>6%</div> </div> </div>
1	B	468	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>.</div> </div> </div>
2	C	81	<div> <div></div> <div> <div>42%</div> <div>5%</div> <div>53%</div> </div> </div>
2	D	81	<div> <div>2%</div> <div> <div></div> <div>65%</div> <div>11%</div> <div>23%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transitional endoplasmic reticulum ATPase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	440	Total	C	N	O	S	0	0	0
			3453	2167	613	655	18			
1	B	447	Total	C	N	O	S	0	0	0
			3506	2202	623	663	18			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	198	TRP	LEU	engineered mutation	UNP P55072
A	461	ARG	-	expression tag	UNP P55072
A	462	SER	-	expression tag	UNP P55072
A	463	HIS	-	expression tag	UNP P55072
A	464	HIS	-	expression tag	UNP P55072
A	465	HIS	-	expression tag	UNP P55072
A	466	HIS	-	expression tag	UNP P55072
A	467	HIS	-	expression tag	UNP P55072
A	468	HIS	-	expression tag	UNP P55072
B	198	TRP	LEU	engineered mutation	UNP P55072
B	461	ARG	-	expression tag	UNP P55072
B	462	SER	-	expression tag	UNP P55072
B	463	HIS	-	expression tag	UNP P55072
B	464	HIS	-	expression tag	UNP P55072
B	465	HIS	-	expression tag	UNP P55072
B	466	HIS	-	expression tag	UNP P55072
B	467	HIS	-	expression tag	UNP P55072
B	468	HIS	-	expression tag	UNP P55072

- Molecule 2 is a protein called Selenoprotein S.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	38	Total	C	N	O	S	0	0	0
			304	189	57	57	1			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	62	Total	C	N	O	S	0	0	0
			517	317	112	87	1			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	42	MET	-	initiating methionine	UNP Q9BQE4
C	43	HIS	-	expression tag	UNP Q9BQE4
C	44	HIS	-	expression tag	UNP Q9BQE4
C	45	HIS	-	expression tag	UNP Q9BQE4
C	46	HIS	-	expression tag	UNP Q9BQE4
C	47	HIS	-	expression tag	UNP Q9BQE4
C	48	HIS	-	expression tag	UNP Q9BQE4
D	42	MET	-	initiating methionine	UNP Q9BQE4
D	43	HIS	-	expression tag	UNP Q9BQE4
D	44	HIS	-	expression tag	UNP Q9BQE4
D	45	HIS	-	expression tag	UNP Q9BQE4
D	46	HIS	-	expression tag	UNP Q9BQE4
D	47	HIS	-	expression tag	UNP Q9BQE4
D	48	HIS	-	expression tag	UNP Q9BQE4

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		



4 Data and refinement statistics

Property	Value	Source
Space group	P 6 2 2	Depositor
Cell constants a, b, c, α , β , γ	153.77Å 153.77Å 240.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.27 – 3.41 49.27 – 3.41	Depositor EDS
% Data completeness (in resolution range)	97.0 (49.27-3.41) 97.0 (49.27-3.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 3.40Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.233 , 0.294 0.231 , 0.287	Depositor DCC
R_{free} test set	1135 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	86.2	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 65.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.19$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7845	wwPDB-VP
Average B, all atoms (Å ²)	135.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.6606e-03.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/3508	0.98	5/4741 (0.1%)
1	B	0.77	0/3561	0.99	7/4811 (0.1%)
2	C	0.78	0/304	0.86	0/403
2	D	0.86	0/523	0.93	0/695
All	All	0.76	0/7896	0.97	12/10650 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	5
2	D	0	1
All	All	0	10

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	140	LEU	CD1-CG-CD2	-8.92	83.74	110.50
1	A	140	LEU	CB-CG-CD1	8.88	126.10	111.00
1	A	140	LEU	CD1-CG-CD2	-8.31	85.56	110.50
1	B	140	LEU	CB-CG-CD2	8.24	125.01	111.00
1	A	140	LEU	CB-CG-CD2	7.65	124.01	111.00

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	ILE	Peptide
1	A	169	ASP	Peptide
1	A	171	SER	Peptide
1	A	297	ALA	Peptide
1	B	169	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3453	0	3496	30	0
1	B	3506	0	3559	35	0
2	C	304	0	319	2	0
2	D	517	0	530	6	0
3	A	31	0	13	1	0
3	B	31	0	13	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
All	All	7845	0	7930	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 69 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:VAL:HB	1:B:126:ILE:HG22	1.70	0.73
1:A:126:ILE:HD11	1:A:130:LEU:HD11	1.71	0.72
1:B:131:PHE:O	1:B:136:LYS:HB2	1.90	0.71
1:A:123:VAL:HB	1:A:126:ILE:HD13	1.71	0.71
1:A:31:ALA:HA	1:A:83:ARG:HB3	1.74	0.69

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/468 (94%)	397 (91%)	38 (9%)	3 (1%)	22	58
1	B	445/468 (95%)	404 (91%)	39 (9%)	2 (0%)	34	69
2	C	36/81 (44%)	36 (100%)	0	0	100	100
2	D	58/81 (72%)	53 (91%)	5 (9%)	0	100	100
All	All	977/1098 (89%)	890 (91%)	82 (8%)	5 (0%)	29	65

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	172	PRO
1	A	127	THR
1	B	170	PRO
1	A	298	PRO
1	A	125	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	377/400 (94%)	377 (100%)	0	100	100
1	B	382/400 (96%)	381 (100%)	1 (0%)	92	97
2	C	31/70 (44%)	31 (100%)	0	100	100
2	D	53/70 (76%)	51 (96%)	2 (4%)	33	64
All	All	843/940 (90%)	840 (100%)	3 (0%)	91	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	40	SER
2	D	47	HIS
2	D	54	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ANP	B	800	4	29,33,33	1.91	8 (27%)	31,52,52	1.77	8 (25%)
3	ANP	A	800	4	29,33,33	1.78	7 (24%)	31,52,52	2.18	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	B	800	4	-	6/14/38/38	0/3/3/3
3	ANP	A	800	4	-	5/14/38/38	0/3/3/3

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	800	ANP	PG-O1G	4.74	1.53	1.46
3	B	800	ANP	PG-N3B	4.42	1.74	1.63
3	A	800	ANP	PG-N3B	4.20	1.74	1.63
3	B	800	ANP	PB-N3B	4.13	1.74	1.63
3	A	800	ANP	PB-N3B	3.89	1.73	1.63

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	800	ANP	O1G-PG-N3B	-6.35	102.42	111.77
3	A	800	ANP	O1B-PB-N3B	-4.79	104.72	111.77
3	B	800	ANP	O1G-PG-N3B	-4.60	105.00	111.77
3	A	800	ANP	O2B-PB-O1B	4.38	119.10	109.92
3	A	800	ANP	N3-C2-N1	-3.81	122.73	128.68

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	800	ANP	PB-N3B-PG-O1G
3	B	800	ANP	PG-N3B-PB-O1B
3	B	800	ANP	C5'-O5'-PA-O3A
3	A	800	ANP	PG-N3B-PB-O1B
3	A	800	ANP	C5'-O5'-PA-O2A


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	800	ANP	2	0
3	A	800	ANP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	440/468 (94%)	-0.19	10 (2%) 60 59	61, 126, 203, 257	0
1	B	447/468 (95%)	-0.27	5 (1%) 80 79	49, 120, 199, 276	0
2	C	38/81 (46%)	-0.10	0 100 100	145, 175, 235, 263	0
2	D	62/81 (76%)	0.05	2 (3%) 47 47	158, 193, 227, 258	0
All	All	987/1098 (89%)	-0.21	17 (1%) 70 69	49, 130, 215, 276	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437	ILE	5.1
1	B	158	MET	4.1
1	A	435	GLU	4.0
1	A	436	THR	3.6
1	A	429	LEU	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ANP	A	800	31/31	0.95	0.21	68,82,99,117	0
3	ANP	B	800	31/31	0.96	0.20	58,63,73,77	0
4	MG	A	801	1/1	0.96	0.23	75,75,75,75	0
4	MG	B	801	1/1	0.99	0.25	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.