



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 2, 2021 – 10:10 AM EST

PDB ID : 7KIE  
Title : Crystal structure of FGFR2 kinase domain gatekeeper mutant V564F in complex with covalent compound 3  
Authors : Ke, J.; Wibowo, A.S.; Carter, J.J.; Larsen, N.A.  
Deposited on : 2020-10-23  
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	
1	B	308	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	A	802	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4951 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fibroblast growth factor receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	S	0	1	0
			2333	1484	397	431	21			
1	B	292	Total	C	N	O	S	0	1	0
			2349	1499	395	434	21			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	564	PHE	VAL	engineered mutation	UNP P21802
B	564	PHE	VAL	engineered mutation	UNP P21802

- Molecule 2 is N-{4-[(E)-2-{4-(4-methylpiperazin-1-yl)-6-[(5-methyl-1H-pyrazol-3-yl)amino]pyrimidin-2-yl}ethenyl]phenyl}prop-2-enamide (three-letter code: WF7) (formula: C<sub>24</sub>H<sub>28</sub>N<sub>8</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			33	24	8	1		
2	B	1	Total	C	N	O	0	0
			33	24	8	1		

- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula:  $C_6H_5O_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	81	Total	O	0	0
			81	81		
5	B	84	Total	O	0	0
			84	84		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fibroblast growth factor receptor 2



- Molecule 1: Fibroblast growth factor receptor 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.30Å 116.91Å 64.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.01 – 2.47 56.59 – 2.47	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.01-2.47) 100.0 (56.59-2.47)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.201 , 0.229 0.202 , 0.230	Depositor DCC
$R_{free}$ test set	1471 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.4	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 27.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4951	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FLC, WF7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/2383	0.69	0/3218
1	B	0.31	0/2404	0.68	0/3246
All	All	0.31	0/4787	0.68	0/6464

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2333	0	2331	23	0
1	B	2349	0	2347	16	0
2	A	33	0	0	0	0
2	B	33	0	0	1	0
3	A	13	0	5	7	0
3	B	13	0	5	3	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	81	0	0	3	0
5	B	84	0	0	1	0
All	All	4951	0	4704	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:649[B]:ARG:NH1	3:A:802:FLC:CG	2.44	0.81
1:A:649[B]:ARG:NH1	3:A:802:FLC:HG2	2.01	0.76
1:A:649[B]:ARG:NH1	3:A:802:FLC:HG1	2.05	0.72
1:A:491:CYS:HB3	1:B:491:CYS:HB3	1.71	0.72
1:A:625:ARG:HH22	3:A:802:FLC:HA2	1.56	0.71
1:A:649[B]:ARG:HH12	3:A:802:FLC:HG1	1.57	0.69
1:A:741:HIS:HB3	5:A:951:HOH:O	1.92	0.69
1:B:541:ILE:HD11	1:B:617:LEU:CD2	2.26	0.66
1:B:508:PRO:O	1:B:509:LYS:HB2	1.95	0.64
1:B:625:ARG:HH22	3:B:802:FLC:HA1	1.65	0.60
1:A:487:LEU:HD12	1:A:495:VAL:HG12	1.82	0.60
1:A:664:ARG:NH2	3:A:802:FLC:OB2	2.30	0.59
1:B:541:ILE:HD11	1:B:617:LEU:HD22	1.86	0.56
1:B:664:ARG:HE	3:B:802:FLC:HG1	1.71	0.55
1:B:625:ARG:HH22	3:B:802:FLC:CA	2.18	0.55
1:A:667:VAL:HG13	1:A:713:PHE:CE2	2.42	0.54
1:B:577:ARG:NH1	5:B:901:HOH:O	2.41	0.53
1:B:573:ARG:O	1:B:577:ARG:HG3	2.11	0.50
1:A:503:ILE:HG22	1:A:512:VAL:HG13	1.94	0.50
1:A:625:ARG:NH2	3:A:802:FLC:HA2	2.26	0.49
1:A:660:THR:HG23	5:A:941:HOH:O	2.13	0.48
2:B:801:WF7:C10	2:B:801:WF7:N6	2.74	0.48
1:B:726:ALA:O	1:B:727:ASN:HB3	2.14	0.47
1:B:544:HIS:HB3	1:B:547:ILE:HD12	1.97	0.47
1:A:538:MET:HB2	1:A:550:LEU:HD22	1.98	0.46
1:A:667:VAL:HG13	1:A:713:PHE:HE2	1.80	0.46
1:B:731:GLU:OE2	1:B:759:ARG:NH2	2.45	0.46
1:A:668:LYS:HG2	1:A:712:LEU:HD22	1.98	0.45
1:A:621:LYS:HD2	5:A:962:HOH:O	2.15	0.45
1:A:495:VAL:HG22	1:A:517:LYS:HG3	1.98	0.45
1:A:724:LYS:NZ	1:A:728:CYS:O	2.44	0.44
1:B:658:LYS:HA	1:B:658:LYS:HD3	1.81	0.43
1:A:538:MET:HE1	1:A:645:PHE:CD1	2.54	0.43
1:A:610:LEU:HD13	1:A:692:LEU:HD21	2.01	0.42
1:A:617:LEU:HD23	1:A:617:LEU:HA	1.82	0.42
1:B:541:ILE:HD11	1:B:617:LEU:HD21	2.02	0.42
1:A:600:PHE:CZ	1:A:729:THR:HG23	2.54	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:610:LEU:HD13	1:B:692:LEU:HD21	2.03	0.41
1:B:676[A]:PHE:HZ	1:B:713:PHE:HB3	1.86	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/308 (94%)	279 (97%)	9 (3%)	0	100	100
1	B	289/308 (94%)	277 (96%)	11 (4%)	1 (0%)	41	59
All	All	577/616 (94%)	556 (96%)	20 (4%)	1 (0%)	47	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	597	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/270 (94%)	240 (94%)	14 (6%)	21	39
1	B	257/270 (95%)	245 (95%)	12 (5%)	26	46
All	All	511/540 (95%)	485 (95%)	26 (5%)	24	42

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	489	GLU
1	A	491	CYS
1	A	512	VAL
1	A	520	LYS
1	A	521	ASP
1	A	543	LYS
1	A	594	PRO
1	A	595	GLU
1	A	601	LYS
1	A	606	CYS
1	A	621	LYS
1	A	659	LYS
1	A	660	THR
1	A	763	LEU
1	B	470	GLU
1	B	485	LYS
1	B	492	PHE
1	B	506	ASP
1	B	507	LYS
1	B	521	ASP
1	B	540	MET
1	B	595	GLU
1	B	630	ARG
1	B	644	ASP
1	B	727	ASN
1	B	767	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	B	803	-	5,5,5	0.14	0	5,5,5	0.40	0
2	WF7	A	801	1	35,36,36	1.89	8 (22%)	43,49,49	2.30	10 (23%)
3	FLC	A	802	-	3,12,12	1.25	0	3,17,17	0.30	0
2	WF7	B	801	1	35,36,36	1.97	10 (28%)	43,49,49	2.28	13 (30%)
3	FLC	B	802	-	3,12,12	1.26	0	3,17,17	0.82	0
4	GOL	A	803	-	5,5,5	0.10	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	803	-	-	4/4/4/4	-
2	WF7	A	801	1	-	6/17/29/29	0/4/4/4
3	FLC	A	802	-	-	1/6/16/16	-
2	WF7	B	801	1	-	0/17/29/29	0/4/4/4
3	FLC	B	802	-	-	4/6/16/16	-
4	GOL	A	803	-	-	3/4/4/4	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	WF7	C22-C21	5.06	1.55	1.30
2	B	801	WF7	C22-C21	4.96	1.54	1.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	WF7	C21-C20	4.87	1.56	1.48
2	B	801	WF7	C14-C15	4.78	1.55	1.31
2	A	801	WF7	C14-C15	4.61	1.54	1.31
2	B	801	WF7	C21-C20	3.95	1.54	1.48
2	B	801	WF7	C8-N3	-3.30	1.32	1.38
2	B	801	WF7	N5-N4	-2.87	1.32	1.37
2	A	801	WF7	C4-N2	2.86	1.51	1.46
2	A	801	WF7	C19-N7	-2.68	1.36	1.41
2	B	801	WF7	C19-N7	-2.51	1.36	1.41
2	B	801	WF7	C12-C11	2.47	1.56	1.50
2	B	801	WF7	C3-N2	2.47	1.50	1.46
2	B	801	WF7	C10-C11	-2.38	1.34	1.39
2	A	801	WF7	C16-C15	-2.18	1.41	1.47
2	A	801	WF7	C9-N3	-2.13	1.34	1.38
2	B	801	WF7	C9-N3	-2.09	1.34	1.38
2	A	801	WF7	C3-N2	2.08	1.50	1.46

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	WF7	C22-C21-C20	-8.59	112.45	122.27
2	A	801	WF7	C6-N8-C13	7.79	123.29	116.02
2	A	801	WF7	C22-C21-C20	-7.38	113.83	122.27
2	B	801	WF7	C6-N8-C13	4.71	120.41	116.02
2	A	801	WF7	N6-C13-N8	-4.51	120.06	125.61
2	B	801	WF7	N6-C13-N8	-4.35	120.26	125.61
2	A	801	WF7	C3-N2-C4	4.31	121.02	111.52
2	B	801	WF7	C3-N2-C4	4.07	120.49	111.52
2	B	801	WF7	N8-C6-N2	3.62	120.62	116.55
2	A	801	WF7	N3-C8-N6	3.22	126.05	116.95
2	A	801	WF7	C19-N7-C20	-3.02	123.70	128.26
2	A	801	WF7	C16-C15-C14	-3.02	113.97	125.87
2	B	801	WF7	C19-N7-C20	-2.92	123.84	128.26
2	A	801	WF7	C7-C6-N8	-2.91	117.48	123.15
2	B	801	WF7	C9-C10-C11	-2.66	104.48	107.23
2	B	801	WF7	O1-C20-N7	2.62	126.06	123.05
2	A	801	WF7	C14-C13-N6	2.45	125.34	117.12
2	B	801	WF7	N3-C8-N6	2.32	123.51	116.95
2	B	801	WF7	O1-C20-C21	-2.27	119.10	122.72
2	B	801	WF7	C7-C6-N2	-2.24	119.64	122.29
2	B	801	WF7	C16-C15-C14	-2.20	117.21	125.87
2	A	801	WF7	N8-C6-N2	2.15	118.96	116.55

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	WF7	C2-N1-C5	2.08	112.43	109.52

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	803	GOL	C1-C2-C3-O3
2	A	801	WF7	O1-C20-C21-C22
2	A	801	WF7	N7-C20-C21-C22
2	A	801	WF7	N8-C13-C14-C15
2	A	801	WF7	N6-C13-C14-C15
3	B	802	FLC	CA-CB-CG-CGC
4	A	803	GOL	O1-C1-C2-C3
3	B	802	FLC	OHB-CB-CG-CGC
2	A	801	WF7	C14-C15-C16-C24
2	A	801	WF7	C14-C15-C16-C17
4	B	803	GOL	O1-C1-C2-C3
4	B	803	GOL	O1-C1-C2-O2
4	A	803	GOL	O1-C1-C2-O2
4	B	803	GOL	O2-C2-C3-O3
4	A	803	GOL	O2-C2-C3-O3
3	B	802	FLC	CBC-CB-CG-CGC
3	A	802	FLC	CAC-CA-CB-CG
3	B	802	FLC	CAC-CA-CB-OHB



There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	FLC	7	0
2	B	801	WF7	1	0
3	B	802	FLC	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	291/308 (94%)	-0.34	3 (1%) 82 84	29, 47, 100, 142	0
1	B	292/308 (94%)	-0.14	6 (2%) 63 65	28, 45, 102, 166	0
All	All	583/616 (94%)	-0.24	9 (1%) 73 75	28, 46, 102, 166	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	466	TYR	7.7
1	B	491	CYS	7.5
1	B	492	PHE	4.1
1	A	491	CYS	3.8
1	A	492	PHE	3.2
1	B	509	LYS	2.9
1	B	470	GLU	2.7
1	B	490	GLY	2.6
1	A	466	TYR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	B	803	6/6	0.81	0.15	66,74,80,85	0
3	FLC	B	802	13/13	0.83	0.36	54,70,89,94	0
3	FLC	A	802	13/13	0.86	0.33	50,60,85,91	0
4	GOL	A	803	6/6	0.87	0.15	63,77,80,80	0
2	WF7	A	801	33/33	0.95	0.16	31,60,113,118	0
2	WF7	B	801	33/33	0.95	0.17	38,65,118,136	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.