



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 07:10 pm BST

PDB ID : 3KJR
Title : Crystal structure of dihydrofolate reductase/thymidylate synthase from *Babesia bovis* determined using SlipChip based microfluidics
Authors : Li, L.; Du, W.; Edwards, T.E.; Staker, B.L.; Phan, I.; Stacy, R.; Ismagilov, R.F.; Accelerated Technologies Center for Gene to 3D Structure (ATCG3D); Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2009-11-03
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>.</div> </div> </div>
1	B	511	<div> <div>14%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>..</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8950 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrofolate reductase/thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	498	Total	C	N	O	S	0	3	0
			3995	2569	683	724	19			
1	B	495	Total	C	N	O	S	0	6	0
			3925	2525	666	715	19			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			13	8	1	3	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	533	Total	O	0	0
			533	533		
5	B	358	Total	O	0	0
			358	358		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydrofolate reductase/thymidylate synthase



- Molecule 1: Dihydrofolate reductase/thymidylate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.11Å 97.56Å 152.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 37.09 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.1 (50.00-1.95) 98.1 (37.09-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.25 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0104	Depositor
R, R_{free}	0.169 , 0.201 0.169 , 0.201	Depositor DCC
R_{free} test set	4477 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8950	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NHE, GOL, NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/4108	0.62	0/5586
1	B	0.56	2/4045 (0.0%)	0.62	0/5513
All	All	0.58	2/8153 (0.0%)	0.62	0/11099

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	393[A]	CYS	CB-SG	-5.77	1.72	1.81
1	B	393[B]	CYS	CB-SG	-5.77	1.72	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3995	0	3940	32	0
1	B	3925	0	3826	33	0
2	A	48	0	25	6	0
2	B	48	0	25	6	0
3	A	13	0	17	1	0
4	A	24	0	32	4	0
4	B	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	533	0	0	7	0
5	B	358	0	0	2	0
All	All	8950	0	7873	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:368:THR:HG22	1:B:369[B]:ASN:ND2	1.53	1.23
1:B:393[A]:CYS:SG	1:B:414:SER:O	2.19	1.01
1:A:302:LYS:NZ	4:A:515:GOL:H32	1.89	0.87
1:B:368:THR:HG22	1:B:369[B]:ASN:HD22	1.43	0.79
1:A:302:LYS:HZ1	4:A:515:GOL:H32	1.49	0.76
1:B:368:THR:CG2	1:B:369[B]:ASN:ND2	2.43	0.75
1:B:124:GLY:HA3	2:B:512:NAP:H5N	1.72	0.72
1:A:385:LEU:HA	1:A:388:MET:HE2	1.70	0.71
4:A:517:GOL:H12	5:A:950:HOH:O	1.89	0.71
1:A:175:ASP:OD1	3:A:513:NHE:H4'1	1.91	0.70
1:B:368:THR:HG22	1:B:369[B]:ASN:HD21	1.55	0.67
1:B:125:GLY:HA3	2:B:512:NAP:O1A	1.95	0.66
1:A:385:LEU:HD23	1:A:388:MET:CE	2.24	0.66
1:A:302:LYS:HZ2	4:A:515:GOL:H32	1.62	0.63
1:A:124:GLY:HA3	2:A:512:NAP:H5N	1.80	0.63
1:B:433:ALA:HA	1:B:443:LEU:HD21	1.80	0.61
1:A:46[A]:THR:HG21	5:A:558:HOH:O	2.01	0.60
1:A:26:LYS:NZ	5:A:841:HOH:O	2.36	0.58
1:A:385:LEU:HD23	1:A:388:MET:HE1	1.84	0.58
1:B:139:ASP:HA	1:B:186:ARG:HG3	1.87	0.56
1:A:126:SER:OG	2:A:512:NAP:O1N	2.20	0.56
1:B:85:ASN:N	1:B:85:ASN:HD22	2.05	0.55
1:B:186:ARG:O	1:B:187:LYS:HB2	2.08	0.54
1:B:369[A]:ASN:ND2	5:B:677:HOH:O	2.34	0.54
4:B:513:GOL:H12	5:B:720:HOH:O	2.07	0.53
2:B:512:NAP:O2X	2:B:512:NAP:O3B	2.26	0.53
1:B:23:ILE:O	2:B:512:NAP:H2N	2.09	0.52
1:A:23:ILE:O	2:A:512:NAP:H2N	2.11	0.51
1:A:385:LEU:HA	1:A:388:MET:CE	2.39	0.50
1:A:68:LYS:HG3	2:A:512:NAP:H51A	1.94	0.49
1:B:8:CYS:HA	1:B:108:ILE:HG21	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:409:MET:HE1	1:B:258:GLY:HA2	1.95	0.48
1:A:391:PRO:HG2	1:B:374:ARG:HD2	1.96	0.48
1:B:324:ARG:NH1	1:B:387:LYS:O	2.42	0.47
1:B:125:GLY:CA	2:B:512:NAP:O1A	2.63	0.47
1:B:64:ILE:HG22	1:B:128:LEU:HD21	1.97	0.47
1:A:51:GLU:CD	1:A:51:GLU:H	2.19	0.47
1:B:368:THR:CG2	1:B:369[B]:ASN:HD22	2.20	0.46
1:A:147:ASN:HB3	1:A:498:ASP:HB3	1.96	0.46
1:B:197:PRO:HG2	1:B:200:GLN:NE2	2.31	0.46
1:A:386:LYS:HE3	5:A:689:HOH:O	2.16	0.45
1:A:102:GLU:O	2:A:512:NAP:H2A	2.16	0.45
1:B:21:LYS:NZ	1:B:158:ASP:OD1	2.45	0.44
1:A:186:ARG:O	1:A:187:LYS:HB2	2.17	0.44
1:A:186:ARG:HD2	5:A:1023:HOH:O	2.18	0.43
1:A:391:PRO:HG2	1:B:374:ARG:CD	2.48	0.43
1:B:124:GLY:HA2	1:B:129:TYR:CZ	2.53	0.43
1:B:374:ARG:HE	1:B:374:ARG:HB2	1.70	0.43
1:A:70:TYR:HB2	1:A:87:ILE:HD12	1.99	0.43
1:A:102:GLU:O	2:A:512:NAP:C2A	2.66	0.43
1:A:433:ALA:HA	1:A:443:LEU:HD21	2.00	0.43
1:A:388:MET:SD	1:A:392:PRO:HD3	2.58	0.42
1:A:371:ASN:HB3	5:A:877:HOH:O	2.19	0.42
1:A:257:PHE:HB2	1:B:400:TYR:CE2	2.55	0.42
1:B:67:ARG:HB2	1:B:89:SER:HB2	2.01	0.42
1:B:95:VAL:HA	1:B:96:PRO:HD2	1.89	0.42
1:B:125:GLY:HA3	2:B:512:NAP:PA	2.59	0.42
1:A:245:LYS:HD2	5:A:748:HOH:O	2.20	0.41
1:B:29:ILE:HA	1:B:30:PRO:HD3	1.85	0.41
1:B:78:LEU:HB3	1:B:79:PRO:HD3	2.03	0.41
1:B:89:SER:HB3	1:B:92:VAL:HG12	2.03	0.41
1:A:324:ARG:HD3	1:A:328:ASP:O	2.21	0.40
1:A:5:TYR:HB3	1:A:137:LEU:HD21	2.03	0.40
1:B:115:VAL:HA	1:B:116:PRO:HD2	1.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/511 (97%)	487 (98%)	9 (2%)	1 (0%)	47	38
1	B	497/511 (97%)	470 (95%)	24 (5%)	3 (1%)	25	14
All	All	994/1022 (97%)	957 (96%)	33 (3%)	4 (0%)	34	22

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	76	ALA
1	B	96	PRO
1	A	332	ILE
1	B	332	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/458 (96%)	439 (100%)	1 (0%)	93	93
1	B	428/458 (93%)	420 (98%)	8 (2%)	57	50
All	All	868/916 (95%)	859 (99%)	9 (1%)	76	74

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	402	SER
1	B	26	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	85	ASN
1	B	87	ILE
1	B	106	THR
1	B	175	ASP
1	B	247	ASN
1	B	365	ARG
1	B	386	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	200	GLN
1	A	303	ASN
1	B	33	HIS
1	B	200	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	513	-	5,5,5	0.44	0	5,5,5	0.72	0
2	NAP	B	512	-	45,52,52	0.91	3 (6%)	56,80,80	1.31	5 (8%)
4	GOL	A	515	-	5,5,5	0.54	0	5,5,5	0.73	0
3	NHE	A	513	-	13,13,13	2.04	1 (7%)	16,17,17	1.42	3 (18%)
4	GOL	A	516	-	5,5,5	0.35	0	5,5,5	0.33	0
2	NAP	A	512	-	45,52,52	0.80	2 (4%)	56,80,80	1.39	7 (12%)
4	GOL	A	517	-	5,5,5	0.37	0	5,5,5	0.53	0
4	GOL	A	514	-	5,5,5	0.63	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	513	-	-	2/4/4/4	-
2	NAP	B	512	-	-	7/31/67/67	0/5/5/5
4	GOL	A	515	-	-	4/4/4/4	-
3	NHE	A	513	-	-	4/7/15/15	0/1/1/1
4	GOL	A	516	-	-	4/4/4/4	-
2	NAP	A	512	-	-	5/31/67/67	0/5/5/5
4	GOL	A	517	-	-	2/4/4/4	-
4	GOL	A	514	-	-	2/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	513	NHE	C2-S	-6.95	1.67	1.77
2	B	512	NAP	O4B-C1B	2.55	1.44	1.41
2	B	512	NAP	O4D-C1D	2.43	1.44	1.41
2	A	512	NAP	C7N-N7N	2.33	1.37	1.33
2	B	512	NAP	C2A-N3A	2.12	1.35	1.32
2	A	512	NAP	O4D-C1D	2.04	1.43	1.41

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	512	NAP	N3A-C2A-N1A	-5.40	120.25	128.68
2	B	512	NAP	N3A-C2A-N1A	-4.84	121.12	128.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	513	NHE	O3-S-C2	3.77	111.87	105.77
2	A	512	NAP	O7N-C7N-C3N	-3.09	115.94	119.63
2	B	512	NAP	C3N-C7N-N7N	2.81	121.12	117.75
2	B	512	NAP	O2B-C2B-C3B	2.47	120.64	111.68
3	A	513	NHE	O2-S-C2	2.46	109.88	106.92
2	A	512	NAP	O2B-C2B-C1B	-2.34	101.69	110.10
2	A	512	NAP	O2B-C2B-C3B	2.29	119.97	111.68
3	A	513	NHE	O1-S-C2	-2.26	104.20	106.92
2	B	512	NAP	O7N-C7N-C3N	-2.24	116.95	119.63
2	A	512	NAP	C2A-N1A-C6A	2.21	122.53	118.75
2	A	512	NAP	O2B-P2B-O1X	-2.19	100.93	109.39
2	A	512	NAP	O7N-C7N-N7N	2.19	125.69	122.58
2	B	512	NAP	O2B-C2B-C1B	-2.13	102.43	110.10

There are no chirality outliers.

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	513	GOL	O1-C1-C2-C3
2	B	512	NAP	C3B-C2B-O2B-P2B
2	B	512	NAP	O4D-C1D-N1N-C6N
4	A	515	GOL	C1-C2-C3-O3
4	A	515	GOL	O2-C2-C3-O3
3	A	513	NHE	N-C1-C2-S
3	A	513	NHE	C1-C2-S-O1
4	A	516	GOL	O1-C1-C2-O2
4	A	516	GOL	O1-C1-C2-C3
4	A	516	GOL	C1-C2-C3-O3
2	A	512	NAP	PA-O3-PN-O5D
4	A	517	GOL	O1-C1-C2-C3
2	A	512	NAP	C3B-C2B-O2B-P2B
3	A	513	NHE	C1-C2-S-O3
4	A	515	GOL	O1-C1-C2-C3
4	A	514	GOL	O1-C1-C2-C3
4	B	513	GOL	O1-C1-C2-O2
4	A	516	GOL	O2-C2-C3-O3
4	A	517	GOL	O1-C1-C2-O2
4	A	515	GOL	O1-C1-C2-O2
2	B	512	NAP	PA-O3-PN-O5D
2	A	512	NAP	C2B-O2B-P2B-O1X
2	B	512	NAP	C3B-C4B-C5B-O5B
3	A	513	NHE	C1-C2-S-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	514	GOL	O1-C1-C2-O2
2	A	512	NAP	C4D-C5D-O5D-PN
2	A	512	NAP	C1B-C2B-O2B-P2B
2	B	512	NAP	C4D-C5D-O5D-PN
2	B	512	NAP	O4B-C4B-C5B-O5B
2	B	512	NAP	C5B-O5B-PA-O1A


There are no ring outliers.

6 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	513	GOL	1	0
2	B	512	NAP	6	0
4	A	515	GOL	3	0
3	A	513	NHE	1	0
2	A	512	NAP	6	0
4	A	517	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	498/511 (97%)	-0.09	12 (2%) 59 68	9, 15, 30, 38	0
1	B	495/511 (96%)	0.63	72 (14%) 2 4	8, 15, 36, 43	0
All	All	993/1022 (97%)	0.27	84 (8%) 10 17	8, 15, 33, 43	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	78	LEU	12.4
1	B	95	VAL	9.8
1	B	116	PRO	7.3
1	B	99	LEU	6.4
1	B	113	ALA	6.3
1	B	93	LYS	6.3
1	B	57	PRO	6.1
1	B	8	CYS	6.1
1	B	96	PRO	5.9
1	B	249	THR	5.7
1	B	210	VAL	5.7
1	B	76	ALA	5.6
1	B	54	SER	5.4
1	B	53	LEU	5.4
1	B	74	PRO	5.3
1	B	87	ILE	5.1
1	A	92	VAL	5.1
1	B	104	LEU	4.9
1	B	92	VAL	4.8
1	B	7	GLY	4.7
1	A	210	VAL	4.7
1	B	101	TYR	4.6
1	B	115	VAL	4.5
1	B	345	LEU	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	211	PRO	4.4
1	B	97	GLY	4.4
1	B	106	THR	4.4
1	B	55	LYS	4.3
1	B	77	SER	4.3
1	B	5	TYR	4.3
1	B	79	PRO	4.3
1	B	75	LYS	4.3
1	B	100	VAL	4.2
1	B	209	SER	4.1
1	B	98	CYS	4.0
1	B	322	THR	3.9
1	B	114	ASN	3.8
1	B	211	PRO	3.8
1	B	70	TYR	3.8
1	B	110	ASP	3.7
1	B	105	SER	3.7
1	B	323	ASP	3.6
1	B	318	LYS	3.6
1	B	58	ASP	3.6
1	A	91	THR	3.5
1	B	50	PRO	3.4
1	B	102	GLU	3.4
1	B	317	ASP	3.2
1	B	91	THR	3.1
1	B	321	PHE	3.1
1	B	9	GLY	3.0
1	B	86	VAL	3.0
1	B	111	LEU	2.8
1	A	209	SER	2.8
1	B	73	ILE	2.8
1	B	51	GLU	2.8
1	B	117	HIS	2.7
1	B	107	ALA	2.7
1	B	84	ILE	2.7
1	A	511	ALA	2.6
1	B	393[A]	CYS	2.6
1	B	94	GLU	2.6
1	B	59	ILE	2.5
1	B	186	ARG	2.5
1	B	108	ILE	2.4
1	B	320	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	103	ASP	2.4
1	A	320	GLY	2.4
1	A	54	SER	2.4
1	A	323	ASP	2.4
1	B	52	VAL	2.4
1	B	89	SER	2.3
1	A	212	LYS	2.3
1	B	315	PHE	2.3
1	B	129	TYR	2.2
1	A	50	PRO	2.2
1	B	62	VAL	2.2
1	B	396	PHE	2.1
1	B	109	ARG	2.1
1	B	127	PHE	2.1
1	B	56	ASN	2.0
1	B	88	LEU	2.0
1	B	212	LYS	2.0
1	A	51	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	515	6/6	0.76	0.37	37,40,41,42	0
2	NAP	A	512	48/48	0.76	0.27	12,29,43,44	48
4	GOL	A	514	6/6	0.81	0.31	46,49,52,54	0
4	GOL	B	513	6/6	0.85	0.28	36,42,43,48	0
2	NAP	B	512	48/48	0.89	0.20	9,16,28,29	48


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	517	6/6	0.89	0.14	25,35,37,41	0
4	GOL	A	516	6/6	0.89	0.18	45,50,51,53	0
3	NHE	A	513	13/13	0.93	0.23	34,45,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.