



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2022 – 12:03 PM EDT

PDB ID : 7KPB  
Title : mTNF-alpha hTNFR1 complex  
Authors : Fox III, D.; Conrady, D.G.; Lowe, M.; Ceska, T.  
Deposited on : 2020-11-10  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	<div> <div>4%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	B	158	<div> <div>83%</div> <div>8%</div> <div>9%</div> </div>
1	C	158	<div> <div>%</div> <div>80%</div> <div>7%</div> <div>13%</div> </div>
2	E	144	<div> <div>%</div> <div>91%</div> <div>7%</div> <div>.</div> </div>
2	F	144	<div> <div>3%</div> <div>90%</div> <div>8%</div> <div>.</div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	L	214	 2% 93% 7%
4	H	223	 4% 83% 13% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	B	201	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 8525 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tumor necrosis factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	3	0
			1187	761	207	217	2			
1	B	143	Total	C	N	O	S	0	0	0
			1096	705	188	201	2			
1	C	138	Total	C	N	O	S	0	0	0
			1046	675	176	193	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P01375
B	0	SER	-	expression tag	UNP P01375
C	0	SER	-	expression tag	UNP P01375

- Molecule 2 is a protein called Tumor necrosis factor receptor superfamily member 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	141	Total	C	N	O	S	0	0	0
			1046	629	193	203	21			
2	E	141	Total	C	N	O	S	0	0	0
			1049	630	193	205	21			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	12	GLY	-	expression tag	UNP P19438
F	25	ASP	ASN	engineered mutation	UNP P19438
F	153	SER	CYS	engineered mutation	UNP P19438
E	12	GLY	-	expression tag	UNP P19438
E	25	ASP	ASN	engineered mutation	UNP P19438
E	153	SER	CYS	engineered mutation	UNP P19438


- Molecule 3 is a protein called Fab1974 - Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	213	Total	C	N	O	S	0	0	0
			1500	940	254	300	6			

- Molecule 4 is a protein called Fab1974 - Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	212	Total	C	N	O	S	0	0	0
			1472	941	250	274	7			

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		
5	E	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	E	1	Total	C	O	0	0
			6	3	3		
5	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is 5-(1-{[2-(difluoromethoxy)phenyl]methyl}-2-{[3-(2-oxopyrrolidin-1-yl)phenoxy]methyl}-1H-benzimidazol-6-yl)pyridin-2(1H)-one (three-letter code: D84) (formula:  $C_{31}H_{26}F_2N_4O_4$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	F	N	O	
			41	31	2	4	4	

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	E	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	4	Total	O	0	0
			4	4		
8	B	5	Total	O	0	0
			5	5		
8	C	5	Total	O	0	0
			5	5		
8	F	1	Total	O	0	0
			1	1		
8	E	4	Total	O	0	0
			4	4		
8	L	1	Total	O	0	0
			1	1		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tumor necrosis factor




- Molecule 1: Tumor necrosis factor




- Molecule 1: Tumor necrosis factor




- Molecule 2: Tumor necrosis factor receptor superfamily member 1A





- Molecule 2: Tumor necrosis factor receptor superfamily member 1A



- Molecule 3: Fab1974 - Light Chain



● Molecule 4: Fab1974 - Heavy Chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.51Å 99.51Å 311.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.13 – 3.00 49.13 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.13-3.00) 100.0 (49.13-3.00)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 3.01Å)	Xtriage
Refinement program	PHENIX dev_2443	Depositor
R, $R_{free}$	0.223 , 0.256 0.223 , 0.256	Depositor DCC
$R_{free}$ test set	1618 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	71.1	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 52.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	8525	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GOL, D84

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/1223	0.47	0/1667
1	B	0.26	0/1120	0.47	0/1526
1	C	0.26	0/1069	0.46	0/1456
2	E	0.25	0/1070	0.44	0/1448
2	F	0.26	0/1067	0.44	0/1442
3	L	0.25	0/1536	0.46	0/2111
4	H	0.25	0/1516	0.46	0/2089
All	All	0.26	0/8601	0.46	0/11739

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1187	0	1167	8	0
1	B	1096	0	1079	6	0
1	C	1046	0	1008	6	0
2	E	1049	0	893	6	0
2	F	1046	0	895	6	0
3	L	1500	0	1257	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	1472	0	1204	17	0
5	A	6	0	8	1	0
5	B	12	0	16	1	0
5	E	12	0	16	1	0
5	F	18	0	24	0	0
5	L	6	0	8	0	0
6	C	41	0	0	0	0
7	E	14	0	13	0	0
8	A	4	0	0	0	0
8	B	5	0	0	0	0
8	C	5	0	0	0	0
8	E	4	0	0	0	0
8	F	1	0	0	0	0
8	L	1	0	0	0	0
All	All	8525	0	7588	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:67:ARG:NH1	4:H:90:ASP:OD2	2.14	0.81
4:H:91:THR:HG23	4:H:118:THR:HA	1.81	0.62
1:C:100:PRO:HB3	1:C:116:GLU:HG3	1.85	0.59
2:F:16:PRO:HG2	2:F:19:LYS:HD2	1.87	0.57
1:B:58:ILE:HD11	1:B:126:LEU:HD11	1.88	0.56
4:H:22:CYS:HB3	4:H:79:LEU:HB3	1.90	0.54
1:A:90:LYS:HD3	4:H:59:PHE:CZ	2.43	0.53
2:E:36:GLY:HA2	2:E:90:VAL:O	2.09	0.53
2:E:93:ASP:OD1	2:E:94:THR:N	2.41	0.53
2:F:93:ASP:OD1	2:F:94:THR:N	2.40	0.52
4:H:29:PHE:O	4:H:72:ARG:NH2	2.41	0.52
4:H:196:TRP:CG	4:H:197:PRO:HA	2.45	0.52
2:E:69:HIS:ND1	5:E:202:GOL:O3	2.43	0.52
3:L:161:ASN:HB3	3:L:177:SER:HA	1.92	0.51
1:A:62:VAL:HG12	1:A:150:VAL:HG13	1.91	0.51
3:L:108:ARG:NH1	3:L:109:THR:O	2.42	0.51
1:A:26:LEU:HD11	1:A:150:VAL:HG11	1.92	0.51
1:C:16:VAL:HG12	1:C:30:ASN:HA	1.93	0.51
4:H:83:MET:HE2	4:H:86:LEU:HD21	1.91	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:66:ARG:HG3	3:L:71:TYR:CE2	2.47	0.50
1:B:60:SER:HB3	1:B:80:ILE:HD11	1.94	0.49
4:H:52:ASN:ND2	4:H:54:ASP:OD1	2.45	0.48
3:L:170:ASP:N	3:L:170:ASP:OD1	2.46	0.48
4:H:33:TYR:HB3	4:H:99:GLU:HB3	1.96	0.47
2:F:19:LYS:HG2	2:F:31:THR:HA	1.96	0.47
3:L:166:GLN:HB3	3:L:173:TYR:CE2	2.49	0.47
1:A:20:PRO:O	2:E:75:LYS:NZ	2.33	0.46
4:H:46:GLU:OE2	4:H:61:ARG:NE	2.49	0.46
1:A:75:LEU:HD23	1:A:97:ILE:HD11	1.97	0.46
1:B:11:LYS:HD2	1:C:157:LEU:HB3	1.98	0.46
2:E:19:LYS:HG2	2:E:31:THR:HA	1.98	0.46
3:L:131:SER:HA	3:L:180:THR:HA	1.99	0.45
2:E:102:GLN:HG2	2:E:116:ASN:HA	1.98	0.45
1:B:138:ARG:HD3	5:B:202:GOL:H12	1.99	0.44
1:A:20:PRO:HB3	1:A:144:PHE:HB3	2.00	0.44
2:F:36:GLY:HA2	2:F:90:VAL:O	2.17	0.44
1:B:58:ILE:HG12	1:B:154:ILE:HG22	1.99	0.44
4:H:63:SER:O	4:H:67:ARG:NH2	2.51	0.44
1:C:57:LEU:HB3	1:C:155:ILE:HG23	1.98	0.43
3:L:35:TRP:CZ3	3:L:88:CYS:HB3	2.53	0.43
4:H:51:ILE:HD13	4:H:72:ARG:HB2	1.99	0.43
1:B:20:PRO:HB3	1:B:144:PHE:HB3	2.00	0.43
2:F:108:SER:HB3	2:F:111:LEU:HB3	2.00	0.42
4:H:34:MET:HB3	4:H:79:LEU:HD22	2.01	0.42
1:C:58:ILE:O	1:C:121:GLY:HA2	2.20	0.42
4:H:196:TRP:CD1	4:H:197:PRO:HA	2.55	0.41
2:F:80:MET:HE2	2:F:111:LEU:HD11	2.02	0.41
4:H:127:PRO:HB3	4:H:153:TYR:HB3	2.03	0.41
4:H:98:THR:O	4:H:108:PHE:HA	2.20	0.41
1:C:80:ILE:HD12	1:C:94:LEU:HD12	2.03	0.41
1:A:136:ILE:H	5:A:201:GOL:HO3	1.65	0.40
4:H:47:TRP:O	4:H:61:ARG:HD2	2.21	0.40
1:A:50:VAL:HG21	1:A:126:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles

#### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	154/158 (98%)	149 (97%)	5 (3%)	0	100	100
1	B	139/158 (88%)	136 (98%)	3 (2%)	0	100	100
1	C	132/158 (84%)	129 (98%)	3 (2%)	0	100	100
2	E	139/144 (96%)	133 (96%)	6 (4%)	0	100	100
2	F	139/144 (96%)	137 (99%)	1 (1%)	1 (1%)	22	60
3	L	211/214 (99%)	204 (97%)	7 (3%)	0	100	100
4	H	208/223 (93%)	203 (98%)	5 (2%)	0	100	100
All	All	1122/1199 (94%)	1091 (97%)	30 (3%)	1 (0%)	51	85

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	44	PRO

#### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/134 (91%)	121 (99%)	1 (1%)	81	93
1	B	114/134 (85%)	111 (97%)	3 (3%)	46	78
1	C	107/134 (80%)	107 (100%)	0	100	100
2	E	113/132 (86%)	113 (100%)	0	100	100
2	F	112/132 (85%)	111 (99%)	1 (1%)	78	92

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	L	135/190 (71%)	134 (99%)	1 (1%)	84	94
4	H	119/189 (63%)	118 (99%)	1 (1%)	81	93
All	All	822/1045 (79%)	815 (99%)	7 (1%)	78	92

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	ASP
1	B	21	GLN
1	B	86	SER
1	B	130	ASP
2	F	137	CYS
3	L	33	LEU
4	H	103	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	L	198	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	E	202	-	5,5,5	0.36	0	5,5,5	0.41	0
5	GOL	A	201	-	5,5,5	0.35	0	5,5,5	0.33	0
5	GOL	L	301	-	5,5,5	0.36	0	5,5,5	0.41	0
5	GOL	F	203	-	5,5,5	0.34	0	5,5,5	0.36	0
5	GOL	F	201	-	5,5,5	0.37	0	5,5,5	0.27	0
5	GOL	F	202	-	5,5,5	0.34	0	5,5,5	0.34	0
5	GOL	E	203	-	5,5,5	0.35	0	5,5,5	0.27	0
5	GOL	B	202	-	5,5,5	0.38	0	5,5,5	0.17	0
7	NAG	E	201	2	14,14,15	0.77	1 (7%)	17,19,21	0.65	0
5	GOL	B	201	-	5,5,5	0.40	0	5,5,5	0.27	0
6	D84	C	201	-	44,46,46	0.95	1 (2%)	55,65,65	1.57	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	E	202	-	-	2/4/4/4	-
5	GOL	A	201	-	-	2/4/4/4	-
5	GOL	L	301	-	-	0/4/4/4	-
5	GOL	F	203	-	-	2/4/4/4	-
5	GOL	F	201	-	-	2/4/4/4	-
5	GOL	F	202	-	-	4/4/4/4	-
5	GOL	E	203	-	-	0/4/4/4	-
5	GOL	B	202	-	-	4/4/4/4	-
7	NAG	E	201	2	-	4/6/23/26	0/1/1/1
5	GOL	B	201	-	-	2/4/4/4	-
6	D84	C	201	-	-	1/19/31/31	0/6/6/6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	201	D84	C19-N1	-2.29	1.33	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	E	201	NAG	C1-C2	2.06	1.55	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	201	D84	C16-N1-C19	4.79	117.42	112.95
6	C	201	D84	O2-C19-N1	4.66	128.38	125.40
6	C	201	D84	C29-C28-N3	3.92	120.47	115.38
6	C	201	D84	C27-N3-C28	-3.67	121.22	124.12
6	C	201	D84	C29-C30-C26	-3.09	118.68	122.21
6	C	201	D84	O3-C28-C29	-3.04	119.81	125.16
6	C	201	D84	C17-C16-N1	-2.08	101.19	103.42
6	C	201	D84	C27-C26-C30	2.06	119.39	117.26
6	C	201	D84	O-C1-C6	2.01	119.58	116.04

There are no chirality outliers.

All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	F	201	GOL	O1-C1-C2-C3
5	F	203	GOL	O1-C1-C2-C3
5	E	202	GOL	O1-C1-C2-C3
7	E	201	NAG	C1-C2-N2-C7
5	B	202	GOL	O2-C2-C3-O3
5	A	201	GOL	O1-C1-C2-C3
5	B	201	GOL	C1-C2-C3-O3
5	B	202	GOL	O1-C1-C2-C3
5	B	202	GOL	C1-C2-C3-O3
5	F	202	GOL	O1-C1-C2-C3
5	F	202	GOL	C1-C2-C3-O3
5	F	201	GOL	O1-C1-C2-O2
5	F	202	GOL	O1-C1-C2-O2
5	F	203	GOL	O1-C1-C2-O2
5	F	202	GOL	O2-C2-C3-O3
5	E	202	GOL	O1-C1-C2-O2
7	E	201	NAG	C4-C5-C6-O6
5	A	201	GOL	O1-C1-C2-O2
6	C	201	D84	C1-C6-C7-N
5	B	201	GOL	O2-C2-C3-O3
5	B	202	GOL	O1-C1-C2-O2
7	E	201	NAG	O5-C5-C6-O6
7	E	201	NAG	C3-C2-N2-C7

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	202	GOL	1	0
5	A	201	GOL	1	0
5	B	202	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/158 (96%)	-0.11	6 (3%) 39 15	32, 49, 112, 163	0
1	B	143/158 (90%)	-0.35	0 100 100	35, 51, 87, 120	0
1	C	138/158 (87%)	-0.34	1 (0%) 87 69	31, 50, 96, 111	0
2	E	141/144 (97%)	-0.19	2 (1%) 75 49	45, 68, 92, 103	0
2	F	141/144 (97%)	0.04	5 (3%) 44 18	40, 67, 125, 169	0
3	L	213/214 (99%)	0.05	5 (2%) 60 31	42, 86, 124, 130	0
4	H	212/223 (95%)	-0.11	9 (4%) 36 14	44, 83, 140, 157	0
All	All	1141/1199 (95%)	-0.13	28 (2%) 57 29	31, 66, 125, 169	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	148	ASN	8.4
2	F	141	ALA	6.6
1	A	71	SER	5.3
1	A	5	SER	4.8
4	H	147	GLY	4.0
2	F	140	HIS	3.7
4	H	127	PRO	3.3
1	A	109	ALA	3.1
2	F	142	GLY	3.0
1	C	36	LEU	2.9
3	L	130	ALA	2.8
2	E	49	ASP	2.7
2	F	13	SER	2.7
1	A	108	GLY	2.4
1	A	72	THR	2.4
3	L	161	ASN	2.4
4	H	150	VAL	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
4	H	131	PRO	2.4
3	L	154	GLU	2.3
1	A	106	PRO	2.2
4	H	206	ALA	2.2
4	H	215	ASP	2.2
2	E	44	PRO	2.2
3	L	180	THR	2.1
3	L	179	LEU	2.1
4	H	126	PRO	2.1
4	H	186	SER	2.0
4	H	153	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	B	202	6/6	0.60	0.40	70,89,96,102	0
7	NAG	E	201	14/15	0.65	0.31	92,109,115,115	0
5	GOL	B	201	6/6	0.72	0.41	73,89,96,114	0
5	GOL	L	301	6/6	0.85	0.30	73,83,88,89	0
5	GOL	F	203	6/6	0.88	0.25	56,65,75,86	0
5	GOL	E	203	6/6	0.89	0.24	53,69,73,81	0
5	GOL	F	202	6/6	0.91	0.23	58,69,71,79	0
5	GOL	F	201	6/6	0.93	0.42	67,71,76,77	0
5	GOL	A	201	6/6	0.93	0.17	52,63,67,68	0
5	GOL	E	202	6/6	0.95	0.14	52,58,60,67	0
6	D84	C	201	41/41	0.96	0.20	28,34,42,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.