



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 11:53 PM BST

PDB ID : 4KQM  
Title : Crystal structure of yeast glycogen synthase E169Q mutant in complex with glucose and UDP  
Authors : Chikwana, V.M.; Hurley, T.D.  
Deposited on : 2013-05-15  
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	724	<div> <div>4%</div> <div> <div></div> <div>73%</div> <div>14%</div> <div>•</div> <div>12%</div> </div> </div>
1	B	724	<div> <div>2%</div> <div> <div></div> <div>72%</div> <div>15%</div> <div>•</div> <div>12%</div> </div> </div>
1	C	724	<div> <div>4%</div> <div> <div></div> <div>72%</div> <div>15%</div> <div>•</div> <div>12%</div> </div> </div>
1	D	724	<div> <div>7%</div> <div> <div></div> <div>72%</div> <div>15%</div> <div>•</div> <div>12%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 20825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gsy2p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	638	Total	C	N	O	S	0	0	0
			5157	3292	903	943	19			
1	B	638	Total	C	N	O	S	0	1	0
			5163	3296	904	944	19			
1	C	638	Total	C	N	O	S	0	0	0
			5157	3292	903	943	19			
1	D	635	Total	C	N	O	S	0	0	0
			5132	3276	900	937	19			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	initiating methionine	UNP E7NKU1
A	-17	GLY	-	expression tag	UNP E7NKU1
A	-16	SER	-	expression tag	UNP E7NKU1
A	-15	SER	-	expression tag	UNP E7NKU1
A	-14	HIS	-	expression tag	UNP E7NKU1
A	-13	HIS	-	expression tag	UNP E7NKU1
A	-12	HIS	-	expression tag	UNP E7NKU1
A	-11	HIS	-	expression tag	UNP E7NKU1
A	-10	HIS	-	expression tag	UNP E7NKU1
A	-9	HIS	-	expression tag	UNP E7NKU1
A	-8	SER	-	expression tag	UNP E7NKU1
A	-7	SER	-	expression tag	UNP E7NKU1
A	-6	GLY	-	expression tag	UNP E7NKU1
A	-5	LEU	-	expression tag	UNP E7NKU1
A	-4	VAL	-	expression tag	UNP E7NKU1
A	-3	PRO	-	expression tag	UNP E7NKU1
A	-2	ARG	-	expression tag	UNP E7NKU1
A	-1	GLY	-	expression tag	UNP E7NKU1
A	0	SER	-	expression tag	UNP E7NKU1
A	169	GLN	GLU	engineered mutation	UNP E7NKU1
B	-18	MET	-	initiating methionine	UNP E7NKU1

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	GLY	-	expression tag	UNP E7NKU1
B	-16	SER	-	expression tag	UNP E7NKU1
B	-15	SER	-	expression tag	UNP E7NKU1
B	-14	HIS	-	expression tag	UNP E7NKU1
B	-13	HIS	-	expression tag	UNP E7NKU1
B	-12	HIS	-	expression tag	UNP E7NKU1
B	-11	HIS	-	expression tag	UNP E7NKU1
B	-10	HIS	-	expression tag	UNP E7NKU1
B	-9	HIS	-	expression tag	UNP E7NKU1
B	-8	SER	-	expression tag	UNP E7NKU1
B	-7	SER	-	expression tag	UNP E7NKU1
B	-6	GLY	-	expression tag	UNP E7NKU1
B	-5	LEU	-	expression tag	UNP E7NKU1
B	-4	VAL	-	expression tag	UNP E7NKU1
B	-3	PRO	-	expression tag	UNP E7NKU1
B	-2	ARG	-	expression tag	UNP E7NKU1
B	-1	GLY	-	expression tag	UNP E7NKU1
B	0	SER	-	expression tag	UNP E7NKU1
B	169	GLN	GLU	engineered mutation	UNP E7NKU1
C	-18	MET	-	initiating methionine	UNP E7NKU1
C	-17	GLY	-	expression tag	UNP E7NKU1
C	-16	SER	-	expression tag	UNP E7NKU1
C	-15	SER	-	expression tag	UNP E7NKU1
C	-14	HIS	-	expression tag	UNP E7NKU1
C	-13	HIS	-	expression tag	UNP E7NKU1
C	-12	HIS	-	expression tag	UNP E7NKU1
C	-11	HIS	-	expression tag	UNP E7NKU1
C	-10	HIS	-	expression tag	UNP E7NKU1
C	-9	HIS	-	expression tag	UNP E7NKU1
C	-8	SER	-	expression tag	UNP E7NKU1
C	-7	SER	-	expression tag	UNP E7NKU1
C	-6	GLY	-	expression tag	UNP E7NKU1
C	-5	LEU	-	expression tag	UNP E7NKU1
C	-4	VAL	-	expression tag	UNP E7NKU1
C	-3	PRO	-	expression tag	UNP E7NKU1
C	-2	ARG	-	expression tag	UNP E7NKU1
C	-1	GLY	-	expression tag	UNP E7NKU1
C	0	SER	-	expression tag	UNP E7NKU1
C	169	GLN	GLU	engineered mutation	UNP E7NKU1
D	-18	MET	-	initiating methionine	UNP E7NKU1
D	-17	GLY	-	expression tag	UNP E7NKU1
D	-16	SER	-	expression tag	UNP E7NKU1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-15	SER	-	expression tag	UNP E7NKU1
D	-14	HIS	-	expression tag	UNP E7NKU1
D	-13	HIS	-	expression tag	UNP E7NKU1
D	-12	HIS	-	expression tag	UNP E7NKU1
D	-11	HIS	-	expression tag	UNP E7NKU1
D	-10	HIS	-	expression tag	UNP E7NKU1
D	-9	HIS	-	expression tag	UNP E7NKU1
D	-8	SER	-	expression tag	UNP E7NKU1
D	-7	SER	-	expression tag	UNP E7NKU1
D	-6	GLY	-	expression tag	UNP E7NKU1
D	-5	LEU	-	expression tag	UNP E7NKU1
D	-4	VAL	-	expression tag	UNP E7NKU1
D	-3	PRO	-	expression tag	UNP E7NKU1
D	-2	ARG	-	expression tag	UNP E7NKU1
D	-1	GLY	-	expression tag	UNP E7NKU1
D	0	SER	-	expression tag	UNP E7NKU1
D	169	GLN	GLU	engineered mutation	UNP E7NKU1


- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula:  $C_9H_{14}N_2O_{12}P_2$ ).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 3 is 6-O-phosphono-alpha-D-glucopyranose (three-letter code: G6P) (formula:  $C_6H_{13}O_9P$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 16	C 6	O 9	P 1	0	0
3	B	1	Total 16	C 6	O 9	P 1	0	0
3	C	1	Total 16	C 6	O 9	P 1	0	0
3	D	1	Total 16	C 6	O 9	P 1	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		
4	C	1	Total	C	O	0	0
			7	4	3		
4	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is alpha-D-glucopyranose (three-letter code: GLC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			12	6	6		


- Molecule 6 is beta-D-glucopyranose (three-letter code: BGC) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			12	6	6		









• Molecule 1: Gsy2p



• Molecule 1: Gsy2p





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	192.71Å 204.44Å 206.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.41 – 2.77 48.41 – 2.77	Depositor EDS
% Data completeness (in resolution range)	98.7 (48.41-2.77) 98.7 (48.41-2.77)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.69 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.201 , 0.241 0.201 , 0.242	Depositor DCC
$R_{free}$ test set	5074 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.7	Xtriage
Anisotropy	0.616	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 35.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.005 for -h,-l,-k	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	20825	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, UDP, BGC, PEG, G6P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/5282	0.59	0/7155
1	B	0.51	0/5291	0.67	0/7167
1	C	0.44	0/5282	0.61	0/7155
1	D	0.45	0/5255	0.59	0/7117
All	All	0.46	0/21110	0.62	0/28594

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5157	0	5072	60	0
1	B	5163	0	5080	57	0
1	C	5157	0	5072	64	0
1	D	5132	0	5053	54	1
2	A	25	0	11	0	0
2	B	25	0	11	1	0
2	C	25	0	11	3	0
2	D	25	0	11	0	0
3	A	16	0	11	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	16	0	11	1	0
3	C	16	0	11	0	0
3	D	16	0	11	0	0
4	A	7	0	10	1	0
4	B	7	0	10	1	0
4	C	7	0	10	2	0
4	D	7	0	10	1	0
5	C	12	0	6	0	0
6	C	12	0	6	0	0
All	All	20825	0	20417	226	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 226 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:GLN:HG2	1:C:280:HIS:CE1	2.00	0.96
1:A:280:HIS:CE1	1:D:283:GLN:HG2	2.07	0.88
1:B:280:HIS:CE1	1:C:283:GLN:HG2	2.10	0.86
1:A:379:GLU:HG2	4:B:803:PEG:H41	1.65	0.78
1:B:349:LYS:HE2	1:B:576:VAL:O	1.85	0.77

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:85:SER:OG	1:D:85:SER:OG[2_555]	1.92	0.28

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	636/724 (88%)	610 (96%)	24 (4%)	2 (0%)	41	70
1	B	637/724 (88%)	620 (97%)	16 (2%)	1 (0%)	47	76
1	C	636/724 (88%)	611 (96%)	25 (4%)	0	100	100
1	D	631/724 (87%)	600 (95%)	29 (5%)	2 (0%)	41	70
All	All	2540/2896 (88%)	2441 (96%)	94 (4%)	5 (0%)	47	76

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	205	GLY
1	D	111	VAL
1	D	544	ASN
1	A	111	VAL
1	B	512	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	553/624 (89%)	523 (95%)	30 (5%)	22	50
1	B	554/624 (89%)	519 (94%)	35 (6%)	18	43
1	C	553/624 (89%)	516 (93%)	37 (7%)	16	40
1	D	550/624 (88%)	514 (94%)	36 (6%)	17	41
All	All	2210/2496 (88%)	2072 (94%)	138 (6%)	18	44

5 of 138 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	556	ARG
1	C	283	GLN
1	D	450	ASP
1	B	590	THR
1	C	181	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	6	GLN
1	C	89	HIS
1	D	249	GLN
1	B	89	HIS
1	B	582	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

### 5.6 Ligand geometry ⓘ

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	UDP	B	801	-	20,26,26	0.78	0	25,40,40	1.17	2 (8%)
3	G6P	A	802	-	16,16,16	0.52	0	24,24,24	0.88	2 (8%)
2	UDP	A	801	-	20,26,26	0.91	0	25,40,40	1.20	2 (8%)
4	PEG	A	803	-	6,6,6	0.48	0	5,5,5	0.28	0
4	PEG	D	803	-	6,6,6	0.42	0	5,5,5	0.37	0
3	G6P	D	802	-	16,16,16	0.50	0	24,24,24	0.98	1 (4%)



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEG	B	803	-	6,6,6	0.45	0	5,5,5	0.42	0
2	UDP	D	801	-	20,26,26	0.85	0	25,40,40	1.15	1 (4%)
5	GLC	C	801	-	12,12,12	0.53	0	17,17,17	0.83	0
3	G6P	B	802	-	16,16,16	0.49	0	24,24,24	0.71	0
2	UDP	C	803	-	20,26,26	0.93	1 (5%)	25,40,40	1.21	2 (8%)
3	G6P	C	804	-	16,16,16	0.58	0	24,24,24	0.87	1 (4%)
4	PEG	C	805	-	6,6,6	0.39	0	5,5,5	0.38	0
6	BGC	C	802	-	12,12,12	0.49	0	17,17,17	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	B	801	-	-	5/14/32/32	0/2/2/2
3	G6P	A	802	-	-	1/6/26/26	0/1/1/1
2	UDP	A	801	-	-	8/14/32/32	0/2/2/2
4	PEG	A	803	-	-	3/4/4/4	-
4	PEG	D	803	-	-	3/4/4/4	-
3	G6P	D	802	-	-	0/6/26/26	0/1/1/1
4	PEG	B	803	-	-	3/4/4/4	-
2	UDP	D	801	-	-	5/14/32/32	0/2/2/2
5	GLC	C	801	-	-	0/2/22/22	0/1/1/1
3	G6P	B	802	-	-	2/6/26/26	0/1/1/1
2	UDP	C	803	-	-	8/14/32/32	0/2/2/2
3	G6P	C	804	-	-	0/6/26/26	0/1/1/1
4	PEG	C	805	-	-	3/4/4/4	-
6	BGC	C	802	-	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	803	UDP	O4'-C1'	2.25	1.44	1.41

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	UDP	C3'-C2'-C1'	3.22	105.83	100.98

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	UDP	PA-O3A-PB	-3.04	122.40	132.83
2	D	801	UDP	PA-O3A-PB	-3.01	122.51	132.83
2	B	801	UDP	PA-O3A-PB	-2.61	123.86	132.83
2	A	801	UDP	C3'-C2'-C1'	2.50	104.75	100.98

There are no chirality outliers.

5 of 41 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	B	801	UDP	C2'-C1'-N1-C6
2	B	801	UDP	O4'-C1'-N1-C6
3	B	802	G6P	C4-C5-C6-O6
2	A	801	UDP	O4'-C1'-N1-C6
2	A	801	UDP	C3'-C4'-C5'-O5'



There are no ring outliers.



8 monomers are involved in 13 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	UDP	1	0
3	A	802	G6P	3	0
4	A	803	PEG	1	0
4	D	803	PEG	1	0
4	B	803	PEG	1	0
3	B	802	G6P	1	0
2	C	803	UDP	3	0
4	C	805	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	638/724 (88%)	0.25	28 (4%) 34 28	41, 61, 110, 139	0
1	B	638/724 (88%)	0.04	15 (2%) 59 54	33, 53, 106, 139	0
1	C	638/724 (88%)	0.22	32 (5%) 28 23	40, 59, 111, 142	1 (0%)
1	D	635/724 (87%)	0.38	50 (7%) 12 9	35, 72, 137, 165	0
All	All	2549/2896 (88%)	0.22	125 (4%) 29 24	33, 59, 123, 165	1 (0%)

The worst 5 of 125 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	630	LEU	7.2
1	D	637	ALA	6.5
1	D	626	VAL	6.1
1	D	627	GLY	5.5
1	C	128	ILE	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.






Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PEG	A	803	7/7	0.86	0.18	73,77,85,85	0
4	PEG	B	803	7/7	0.91	0.26	68,69,76,76	0
4	PEG	D	803	7/7	0.93	0.21	55,65,75,78	0
4	PEG	C	805	7/7	0.94	0.13	62,69,77,78	0
2	UDP	B	801	25/25	0.97	0.15	46,57,85,89	0
2	UDP	C	803	25/25	0.97	0.19	39,51,55,59	0
2	UDP	A	801	25/25	0.97	0.14	47,59,72,73	0
5	GLC	C	801	12/12	0.98	0.15	46,49,51,53	12
3	G6P	B	802	16/16	0.98	0.17	36,41,45,47	0
3	G6P	C	804	16/16	0.98	0.14	43,47,49,52	0
2	UDP	D	801	25/25	0.98	0.12	48,58,70,71	0
6	BGC	C	802	12/12	0.98	0.16	53,57,59,60	12
3	G6P	D	802	16/16	0.99	0.16	35,40,44,46	0
3	G6P	A	802	16/16	0.99	0.17	42,48,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around UDP C 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around UDP A 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around G6P B 802:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around G6P C 804:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around UDP D 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around G6P D 802:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around G6P A 802:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.