



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 09:30 pm BST

PDB ID : 3KRB
Title : Structure of Aldose Reductase from Giardia LambliA at 1.75A Resolution
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2009-11-18
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	
1	B	334	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5675 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	307	Total	C	N	O	S	0	4	0
			2455	1580	412	448	15			
1	B	307	Total	C	N	O	S	0	5	0
			2453	1574	417	447	15			


There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP A8BGX6
A	-19	ALA	-	EXPRESSION TAG	UNP A8BGX6
A	-18	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-17	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-16	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-15	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-14	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-13	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-12	MET	-	EXPRESSION TAG	UNP A8BGX6
A	-11	GLY	-	EXPRESSION TAG	UNP A8BGX6
A	-10	THR	-	EXPRESSION TAG	UNP A8BGX6
A	-9	LEU	-	EXPRESSION TAG	UNP A8BGX6
A	-8	GLU	-	EXPRESSION TAG	UNP A8BGX6
A	-7	ALA	-	EXPRESSION TAG	UNP A8BGX6
A	-6	GLN	-	EXPRESSION TAG	UNP A8BGX6
A	-5	THR	-	EXPRESSION TAG	UNP A8BGX6
A	-4	GLN	-	EXPRESSION TAG	UNP A8BGX6
A	-3	GLY	-	EXPRESSION TAG	UNP A8BGX6
A	-2	PRO	-	EXPRESSION TAG	UNP A8BGX6
A	-1	GLY	-	EXPRESSION TAG	UNP A8BGX6
A	0	SER	-	EXPRESSION TAG	UNP A8BGX6
B	-20	MET	-	EXPRESSION TAG	UNP A8BGX6
B	-19	ALA	-	EXPRESSION TAG	UNP A8BGX6
B	-18	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-17	HIS	-	EXPRESSION TAG	UNP A8BGX6

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-15	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-14	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-13	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-12	MET	-	EXPRESSION TAG	UNP A8BGX6
B	-11	GLY	-	EXPRESSION TAG	UNP A8BGX6
B	-10	THR	-	EXPRESSION TAG	UNP A8BGX6
B	-9	LEU	-	EXPRESSION TAG	UNP A8BGX6
B	-8	GLU	-	EXPRESSION TAG	UNP A8BGX6
B	-7	ALA	-	EXPRESSION TAG	UNP A8BGX6
B	-6	GLN	-	EXPRESSION TAG	UNP A8BGX6
B	-5	THR	-	EXPRESSION TAG	UNP A8BGX6
B	-4	GLN	-	EXPRESSION TAG	UNP A8BGX6
B	-3	GLY	-	EXPRESSION TAG	UNP A8BGX6
B	-2	PRO	-	EXPRESSION TAG	UNP A8BGX6
B	-1	GLY	-	EXPRESSION TAG	UNP A8BGX6
B	0	SER	-	EXPRESSION TAG	UNP A8BGX6

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $\text{C}_{21}\text{H}_{28}\text{N}_7\text{O}_{17}\text{P}_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	B	1	Total 48	C 21	N 7	O 17	P 3	0	0

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total X 1 1	0	0
3	A	1	Total X 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	379	Total O 379 379	0	0
5	B	282	Total O 282 282	0	0

- Molecule 1: Aldose reductase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	196.77Å 66.09Å 56.29Å 90.00° 92.26° 90.00°	Depositor
Resolution (Å)	50.00 – 1.75 46.54 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-1.75) 99.9 (46.54-1.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.5.0104	Depositor
R, R_{free}	0.144 , 0.173 0.179 , 0.203	Depositor DCC
R_{free} test set	3672 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5675	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, NAP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/2532	0.74	2/3446 (0.1%)
1	B	0.70	0/2533	0.70	1/3452 (0.0%)
All	All	0.73	0/5065	0.72	3/6898 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	LYS	CD-CE-NZ	5.20	123.65	111.70
1	A	6	ARG	NE-CZ-NH2	-5.09	117.75	120.30
1	B	6	ARG	NE-CZ-NH2	-5.03	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2455	0	2426	5	0
1	B	2453	0	2400	13	0
2	A	48	0	25	2	0
2	B	48	0	25	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	4	0	6	1	0
4	B	4	0	6	0	0
5	A	379	0	0	1	0
5	B	282	0	0	4	0
All	All	5675	0	4888	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177:GLN:NE2	2:B:400:NAP:H2N	2.09	0.67
1:B:13:GLN:HE22	1:B:220:GLN:HE22	1.43	0.64
1:B:44[B]:GLU:OE2	5:B:445:HOH:O	2.15	0.61
1:B:138[B]:ARG:HA	1:B:138[B]:ARG:HE	1.65	0.61
1:B:156:ASN:HD21	2:B:400:NAP:H72N	1.50	0.60
1:B:142:GLN:NE2	5:B:606:HOH:O	2.37	0.57
1:A:99:ASP:OD1	1:A:150:LYS:CE	2.54	0.56
1:A:156:ASN:HD21	2:A:400:NAP:H72N	1.53	0.56
1:A:113:ASP:OD1	5:A:548:HOH:O	2.18	0.55
1:B:13:GLN:NE2	1:B:220:GLN:HE22	2.08	0.51
1:B:237:LYS:HE3	1:B:285:ASP:OD1	2.12	0.49
1:B:177:GLN:HE21	2:B:400:NAP:H2N	1.77	0.49
1:A:177:GLN:NE2	2:A:400:NAP:H2N	2.28	0.48
1:B:138[A]:ARG:HG3	5:B:444:HOH:O	2.17	0.45
1:B:237:LYS:CE	1:B:285:ASP:OD1	2.65	0.44
4:A:401:EDO:H12	1:B:38:TYR:O	2.20	0.42
1:A:44[A]:GLU:OE1	5:B:445:HOH:O	2.22	0.41
1:B:303:PRO:HB3	1:B:307:TRP:CE3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/334 (92%)	302 (98%)	5 (2%)	0	100	100
1	B	308/334 (92%)	303 (98%)	5 (2%)	0	100	100
All	All	615/668 (92%)	605 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/286 (93%)	263 (99%)	2 (1%)	81	72
1	B	262/286 (92%)	259 (99%)	3 (1%)	73	60
All	All	527/572 (92%)	522 (99%)	5 (1%)	76	67

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	SER
1	A	75	TYR
1	B	75	TYR
1	B	129	GLU
1	B	142	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	GLN
1	A	156	ASN
1	A	177	GLN
1	B	13	GLN
1	B	20	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	142	GLN
1	B	156	ASN
1	B	177	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are unknown - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	B	400	-	45,52,52	1.42	5 (11%)	56,80,80	1.33	7 (12%)
4	EDO	B	401	-	3,3,3	0.43	0	2,2,2	0.29	0
2	NAP	A	400	-	45,52,52	1.44	5 (11%)	56,80,80	1.12	4 (7%)
4	EDO	A	401	-	3,3,3	0.40	0	2,2,2	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	B	400	-	-	4/31/67/67	0/5/5/5
4	EDO	B	401	-	-	0/1/1/1	-
2	NAP	A	400	-	-	5/31/67/67	0/5/5/5
4	EDO	A	401	-	-	0/1/1/1	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	NAP	O7N-C7N	6.50	1.36	1.24
2	B	400	NAP	O7N-C7N	6.22	1.36	1.24
2	A	400	NAP	C2A-N3A	3.61	1.37	1.32
2	B	400	NAP	C2A-N3A	2.91	1.36	1.32
2	B	400	NAP	O4B-C1B	2.79	1.45	1.41
2	B	400	NAP	O4D-C1D	2.40	1.44	1.41
2	A	400	NAP	O4D-C1D	2.32	1.44	1.41
2	B	400	NAP	O2D-C2D	2.29	1.48	1.43
2	A	400	NAP	C7N-N7N	-2.09	1.29	1.33
2	A	400	NAP	P2B-O2B	2.01	1.63	1.59

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	NAP	N3A-C2A-N1A	-3.92	122.56	128.68
2	B	400	NAP	C3N-C7N-N7N	-3.70	113.31	117.75
2	B	400	NAP	N3A-C2A-N1A	-3.49	123.22	128.68
2	B	400	NAP	C2N-C3N-C4N	3.12	121.80	118.26
2	B	400	NAP	O7N-C7N-N7N	2.87	126.65	122.58
2	B	400	NAP	C6N-N1N-C2N	-2.79	119.43	121.97
2	B	400	NAP	C5N-C4N-C3N	-2.71	117.14	120.34
2	A	400	NAP	C3N-C7N-N7N	-2.42	114.84	117.75
2	A	400	NAP	C5N-C4N-C3N	-2.29	117.63	120.34
2	B	400	NAP	O2A-PA-O1A	2.11	122.68	112.24
2	A	400	NAP	C2N-C3N-C4N	2.07	120.60	118.26

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	400	NAP	PA-O3-PN-O5D
2	B	400	NAP	PA-O3-PN-O5D
2	A	400	NAP	C4D-C5D-O5D-PN
2	B	400	NAP	C4D-C5D-O5D-PN

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	400	NAP	C5D-O5D-PN-O3
2	B	400	NAP	C5D-O5D-PN-O3
2	A	400	NAP	PN-O3-PA-O1A
2	A	400	NAP	C2B-O2B-P2B-O3X
2	B	400	NAP	C2B-O2B-P2B-O3X


There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	400	NAP	3	0
2	A	400	NAP	2	0
4	A	401	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	307/334 (91%)	0.24	11 (3%) 42 49	9, 12, 24, 33	0
1	B	307/334 (91%)	0.48	26 (8%) 10 14	11, 15, 26, 36	0
All	All	614/668 (91%)	0.36	37 (6%) 21 27	9, 14, 25, 36	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	219	THR	5.1
1	B	211	ALA	4.8
1	A	57	ALA	3.8
1	B	288	ALA	3.7
1	A	58	SER	3.3
1	B	279	VAL	3.3
1	B	295	ASN	3.3
1	B	0	SER	3.2
1	B	57	ALA	3.1
1	B	232	ALA	3.1
1	B	235	ASP	3.0
1	B	281	LEU	3.0
1	B	287	ASP	2.9
1	A	39	VAL	2.9
1	A	219	THR	2.7
1	B	236	ALA	2.6
1	B	268	ALA	2.6
1	B	291	ASN	2.6
1	A	122	ALA	2.6
1	B	230	LEU	2.5
1	B	233	ILE	2.5
1	B	284	ASP	2.4
1	B	223	VAL	2.3
1	B	283	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	105	TRP	2.3
1	B	28	MET	2.3
1	B	239	THR	2.3
1	A	301	CYS	2.2
1	B	227	CYS	2.2
1	B	280	GLN	2.2
1	A	40	TYR	2.1
1	A	103	VAL	2.1
1	B	229	THR	2.1
1	A	102	LEU	2.1
1	B	56	ASP	2.1
1	A	0	SER	2.1
1	B	276	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	401	4/4	0.70	0.26	42,42,43,45	0
4	EDO	B	401	4/4	0.78	0.20	32,32,33,34	0
3	UNX	B	402	1/1	0.89	0.44	24,24,24,24	0
3	UNX	A	402	1/1	0.92	0.60	23,23,23,23	0
2	NAP	B	400	48/48	0.92	0.13	12,18,22,23	0
2	NAP	A	400	48/48	0.97	0.10	7,12,13,13	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP B 400:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP A 400:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.