



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 03:25 pm BST

PDB ID : 5KR6  
Title : Directed Evolution of Transaminases By Ancestral Reconstruction. Using Old Proteins for New Chemistries  
Authors : Wilding, M.; Newman, J.; Peat, T.S.; Scott, C.  
Deposited on : 2016-07-07  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	
1	B	479	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-aminobutyrate transaminase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	457	Total	C	N	O	S	0	2	0
			3534	2236	612	667	19			
1	B	460	Total	C	N	O	S	0	2	0
			3554	2248	615	671	20			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	111	Total 111	O 111	0	0
3	B	107	Total 107	O 107	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-aminobutyrate transaminase



- Molecule 1: 4-aminobutyrate transaminase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.09Å 123.24Å 63.25Å 90.00° 117.61° 90.00°	Depositor
Resolution (Å)	48.80 – 1.99 41.46 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.80-1.99) 98.0 (41.46-1.99)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.17 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.195 , 0.239 0.203 , 0.245	Depositor DCC
$R_{free}$ test set	2651 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.1	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.046 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7336	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/3622	0.81	4/4915 (0.1%)
1	B	0.68	0/3642	0.82	5/4942 (0.1%)
All	All	0.67	0/7264	0.82	9/9857 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	12	ARG	NE-CZ-NH1	7.74	124.17	120.30
1	A	12	ARG	NE-CZ-NH1	6.77	123.69	120.30
1	A	345	ARG	NE-CZ-NH1	6.60	123.60	120.30
1	A	362	ARG	NE-CZ-NH2	-5.99	117.31	120.30
1	A	345	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	B	345	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	B	72	ASP	CB-CG-OD1	5.32	123.09	118.30
1	B	3	ASP	CB-CG-OD1	5.20	122.98	118.30
1	B	398	ASP	CB-CG-OD1	5.06	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3534	0	3425	12	1
1	B	3554	0	3439	15	1
2	A	15	0	6	1	0
2	B	15	0	7	0	0
3	A	111	0	0	0	0
3	B	107	0	0	0	0
All	All	7336	0	6877	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:45:ASP:OD2	1:B:49:ARG:NH1	1.94	1.00
1:B:47:GLU:OE1	1:B:49:ARG:NH2	2.06	0.88
1:B:159:MET:CE	1:B:177:LEU:HD12	2.08	0.83
1:B:159:MET:HE1	1:B:177:LEU:HD12	1.69	0.73
1:B:2:THR:HG23	1:B:7:LEU:HD12	1.76	0.68
1:A:178:GLU:OE1	1:A:179:PRO:HA	1.94	0.67
1:B:2:THR:HG23	1:B:7:LEU:CD1	2.33	0.58
1:B:167:PRO:HG2	1:B:423[A]:HIS:CE1	2.41	0.55
1:A:167:PRO:HG2	1:A:423[B]:HIS:CE1	2.43	0.54
1:B:398:ASP:HB3	1:B:401:LEU:HD12	1.91	0.52
1:A:175:LEU:HB3	1:A:176:PRO:HA	1.94	0.50
1:B:175:LEU:HB3	1:B:176:PRO:HA	1.95	0.49
1:B:60:CYS:O	1:B:292:SER:HA	2.13	0.49
1:A:60:CYS:O	1:A:292:SER:HA	2.12	0.48
1:A:296:PRO:HG2	1:B:296:PRO:HG2	1.94	0.48
1:A:49:ARG:HD3	1:A:51:TYR:OH	2.16	0.46
1:A:362:ARG:HE	1:A:362:ARG:HB3	1.48	0.46
1:A:75:TYR:CD1	1:B:75:TYR:HB2	2.53	0.44
1:A:262:VAL:HG23	2:A:501:PLP:O3	2.18	0.43
1:B:2:THR:CG2	1:B:7:LEU:HD12	2.47	0.43
1:A:398:ASP:HB3	1:A:401:LEU:HD12	2.01	0.43
1:B:261:VAL:HG12	1:B:288:LYS:HD2	2.00	0.43
1:A:421:MET:HB2	1:A:426:ILE:HB	2.01	0.42
1:A:93:GLU:HB3	1:A:94:PRO:HD3	2.02	0.41
1:B:75:TYR:CE2	1:B:79:LYS:HE2	2.55	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:365:GLU:OE2	1:B:354[A]:ASP:OD1[2_855]	1.60	0.60

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/479 (95%)	437 (96%)	17 (4%)	3 (1%)	22	16
1	B	460/479 (96%)	440 (96%)	17 (4%)	3 (1%)	22	16
All	All	917/958 (96%)	877 (96%)	34 (4%)	6 (1%)	22	16

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	288	LYS
1	B	288	LYS
1	B	287	ALA
1	A	287	ALA
1	A	234	ILE
1	B	234	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/381 (96%)	360 (99%)	4 (1%)	73	78
1	B	365/381 (96%)	361 (99%)	4 (1%)	73	78

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	729/762 (96%)	721 (99%)	8 (1%)	73	78

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	PHE
1	A	49	ARG
1	A	59	TYR
1	A	362	ARG
1	B	16	PHE
1	B	59	TYR
1	B	178	GLU
1	B	448	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	275	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	PLP	A	501	1	15,15,16	3.37	3 (20%)	20,22,23	2.12	8 (40%)
2	PLP	B	501	1	15,15,16	3.74	3 (20%)	20,22,23	2.29	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	501	1	-	0/6/6/8	0/1/1/1
2	PLP	B	501	1	-	0/6/6/8	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	PLP	C5-C4	10.74	1.52	1.40
2	A	501	PLP	C3-C2	8.65	1.49	1.40
2	A	501	PLP	C5-C4	8.32	1.49	1.40
2	B	501	PLP	C3-C2	7.81	1.48	1.40
2	B	501	PLP	C3-C4	4.51	1.49	1.40
2	A	501	PLP	C3-C4	4.27	1.49	1.40

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	PLP	C4A-C4-C5	6.42	127.55	120.94
2	A	501	PLP	C4A-C4-C5	5.33	126.43	120.94
2	B	501	PLP	C3-C4-C5	-4.08	114.34	118.74
2	A	501	PLP	C3-C4-C5	-3.66	114.78	118.74
2	A	501	PLP	C6-C5-C4	3.25	120.72	118.16
2	A	501	PLP	O3P-P-O4P	-2.92	98.95	106.73
2	B	501	PLP	C6-C5-C4	2.91	120.45	118.16
2	B	501	PLP	O4P-P-O1P	-2.76	98.74	106.47
2	B	501	PLP	C6-N1-C2	2.51	123.82	119.17
2	A	501	PLP	O3P-P-O1P	2.41	120.13	110.68
2	A	501	PLP	O3P-P-O2P	2.17	115.94	107.64
2	A	501	PLP	O3-C3-C2	2.16	122.20	117.49
2	B	501	PLP	C2A-C2-C3	-2.13	118.26	120.89

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	PLP	O4P-C5A-C5	2.06	113.28	109.35
2	B	501	PLP	C2A-C2-N1	2.02	121.62	117.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	PLP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	457/479 (95%)	0.03	1 (0%) 95 94	13, 20, 36, 51	0
1	B	460/479 (96%)	0.08	7 (1%) 73 72	13, 20, 39, 52	0
All	All	917/958 (95%)	0.06	8 (0%) 84 83	13, 20, 38, 52	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	192	PRO	3.2
1	B	392	ASP	3.1
1	B	400	SER	2.7
1	B	194	GLY	2.3
1	B	389	ALA	2.3
1	B	393	LYS	2.3
1	B	396	ARG	2.2
1	A	392	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PLP	A	501	15/16	0.95	0.14	18,24,31,31	0
2	PLP	B	501	15/16	0.95	0.13	17,19,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.