



Full wwPDB EM Validation Report ⓘ

Nov 16, 2022 – 07:24 AM EST

PDB ID : 7KSR
EMDB ID : EMD-23024
Title : PRC2:EZH1_A from a dimeric PRC2 bound to a nucleosome
Authors : Grau, D.J.; Armache, K.J.
Deposited on : 2020-11-24
Resolution : 4.10 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	747	
2	D	425	
3	C	739	
4	B	441	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase EZH1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	395	Total	C	N	O	S	0	0
			3098	1958	544	560	36		

- Molecule 2 is a protein called Histone-binding protein RBBP4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	367	Total	C	N	O	S	0	0
			2909	1843	501	557	8		

- Molecule 3 is a protein called Polycomb protein SUZ12.


Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	397	Total	C	N	O	S	0	0
			2932	1849	547	514	22		

- Molecule 4 is a protein called Polycomb protein EED.


Mol	Chain	Residues	Atoms					AltConf	Trace
4	B	365	Total	C	N	O	S	0	0
			2917	1849	514	533	21		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
5	A	7	Total	Zn	0
			7	7	



• Molecule 3: Polycomb protein SUZ12



• Molecule 4: Polycomb protein EED



ASN	ALA	PRO	GLY	ARG	LYS	SER	TRP	GLY	LYS	GLY	LYS	TRP	LYS	SER	K76	K77	C78	K79	F101	F113	L135	C150	V165	E195	L196	K197	F198	L206	D310	G334	E337	N381	K385	A399	K400	I426	D431	A432	S433	L440	ARG											
ASN	ALA	PRO	GLY	ARG	LYS	SER	TRP	GLY	LYS	GLY	LYS	TRP	LYS	SER	K76	K77	C78	K79	F101	F113	L135	C150	V165	E195	L196	K197	F198	L206	D310	G334	E337	N381	K385	A399	K400	I426	D431	A432	S433	L440	ARG											
MET	SER	GLU	ARG	GLU	VAL	SER	SER	THR	ALA	PRO	ALA	GLY	THR	ASP	MET	PRO	ALA	ALA	LYS	LYS	GLN	LYS	LEU	SER	SER	ASP	GLU	ASN	SER	ASN	PRO	ASP	ASN	ASP	ALA	VAL	SER	ILE	GLU	GLY	THR	ASN	THR	GLU	ARG	ASP	THR	PRO	THR	ASN	THR	PRO

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	18151	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION; Patch CTF correction	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	56.7	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.043	Depositor
Minimum map value	-0.030	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.007	Depositor
Map size (\AA)	405.0, 405.0, 405.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.35, 1.35, 1.35	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/3170	0.56	0/4283
2	D	0.28	0/2987	0.57	0/4073
3	C	0.27	0/2982	0.58	0/4029
4	B	0.27	0/2992	0.59	0/4058
All	All	0.27	0/12131	0.58	0/16443

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	2945	10	0
2	D	2909	0	2770	18	0
3	C	2932	0	2622	7	0
4	B	2917	0	2802	9	0
5	A	7	0	0	0	0
All	All	11863	0	11139	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:366:LEU:HD21	2:D:369:ILE:HD11	1.61	0.80
3:C:427:ARG:O	3:C:485:ALA:N	2.32	0.63
1:A:459:LEU:O	3:C:682:ARG:NH1	2.33	0.62
2:D:111:VAL:O	2:D:111:VAL:HG13	2.02	0.59
2:D:80:VAL:HG13	2:D:117:ILE:HG23	1.87	0.55
2:D:373:HIS:ND1	2:D:394:SER:OG	2.39	0.55
1:A:307:CYS:SG	1:A:310:HIS:ND1	2.81	0.54
3:C:610:GLU:N	3:C:610:GLU:OE1	2.40	0.54
4:B:195:GLU:OE2	4:B:197:LYS:NZ	2.38	0.54
2:D:135:GLN:NE2	2:D:186:ASN:O	2.40	0.54
3:C:136:ASP:O	3:C:139:SER:OG	2.22	0.53
2:D:254:ILE:O	2:D:266:SER:N	2.41	0.53
4:B:101:PHE:CZ	4:B:426:ILE:HD12	2.44	0.53
1:A:554:CYS:SG	1:A:555:GLN:N	2.83	0.52
1:A:524:CYS:SG	1:A:526:HIS:NE2	2.83	0.52
1:A:274:THR:O	3:C:607:ASN:ND2	2.42	0.51
4:B:381:ASN:ND2	4:B:385:LYS:O	2.44	0.50
2:D:147:SER:OG	2:D:176:HIS:O	2.20	0.50
1:A:597:CYS:SG	1:A:598:LYS:N	2.85	0.49
1:A:544:CYS:SG	1:A:556:ASN:ND2	2.86	0.48
2:D:354:GLN:OE1	2:D:354:GLN:N	2.47	0.48
2:D:390:ILE:HD11	2:D:402:TRP:CZ2	2.50	0.47
3:C:91:GLU:O	3:C:95:GLN:N	2.48	0.45
4:B:101:PHE:HZ	4:B:426:ILE:HD12	1.83	0.43
4:B:431:ASP:OD1	4:B:433:SER:OG	2.24	0.43
2:D:38:HIS:O	2:D:399:MET:N	2.48	0.43
1:A:299:CYS:SG	1:A:310:HIS:ND1	2.92	0.42
2:D:111:VAL:O	2:D:111:VAL:HG22	2.18	0.42
2:D:374:THR:OG1	2:D:396:ASP:OD2	2.38	0.42
4:B:150:CYS:SG	4:B:165:VAL:HG12	2.59	0.42
1:A:68:VAL:HG11	4:B:135:LEU:HD13	2.01	0.42
2:D:311:HIS:CG	2:D:350:ILE:HD11	2.55	0.42
2:D:277:ASN:OD1	2:D:294:ALA:N	2.53	0.42
2:D:173:LEU:HD22	2:D:205:TRP:CG	2.55	0.42
4:B:113:PHE:CZ	4:B:426:ILE:HD13	2.55	0.42
3:C:87:LEU:O	3:C:91:GLU:N	2.53	0.41
4:B:198:PHE:CZ	4:B:206:LEU:HD13	2.55	0.41
2:D:64:HIS:CE1	2:D:86:LEU:HD12	2.56	0.41
1:A:670:LEU:HD11	1:A:707:HIS:HB3	2.03	0.41
2:D:236:HIS:NE2	2:D:286:GLU:O	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:295:ASP:O	2:D:295:ASP:CG	2.59	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/747 (52%)	364 (94%)	21 (6%)	0	100	100
2	D	359/425 (84%)	338 (94%)	21 (6%)	0	100	100
3	C	377/739 (51%)	355 (94%)	22 (6%)	0	100	100
4	B	363/441 (82%)	345 (95%)	18 (5%)	0	100	100
All	All	1484/2352 (63%)	1402 (94%)	82 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/670 (50%)	335 (100%)	0	100	100
2	D	322/375 (86%)	322 (100%)	0	100	100
3	C	267/646 (41%)	267 (100%)	0	100	100
4	B	316/392 (81%)	316 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	1240/2083 (60%)	1240 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	556	ASN
2	D	27	ASN
3	C	104	ASN
4	B	204	ASN
4	B	219	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-23024. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 147



Y Index: 98



Z Index: 119

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.007. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 129 nm³; this corresponds to an approximate mass of 117 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.244 Å⁻¹


8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-23024 and PDB model 7KSR. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.007 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.007).

9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 68% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.007) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.6817	<div></div> 0.2870
A	<div></div> 0.7534	<div></div> 0.3380
B	<div></div> 0.7786	<div></div> 0.3680
C	<div></div> 0.5772	<div></div> 0.2190
D	<div></div> 0.6135	<div></div> 0.2200

