



Full wwPDB X-ray Structure Validation Report ⓘ

May 4, 2022 – 12:08 PM EDT

PDB ID : 7KTM
Title : DNA Polymerase Mu (K438D), 8-oxodGTP:Ct Reaction State Ternary Complex, 50 mM Mn²⁺ (30min)
Authors : Jamsen, J.A.; Wilson, S.H.
Deposited on : 2020-11-24
Resolution : 1.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.28.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.28.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	 85% 7% • 8%
2	T	9	 33% 56% 11%
3	P	5	 40% 20% 40%
4	D	4	 75% 25%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 3570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed DNA/RNA polymerase mu.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	20	0
			2665	1699	479	475	12			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	GLY	-	expression tag	UNP Q9NP87
A	128	SER	-	expression tag	UNP Q9NP87
A	129	ALA	-	expression tag	UNP Q9NP87
A	130	ALA	-	expression tag	UNP Q9NP87
A	131	ALA	-	expression tag	UNP Q9NP87
A	?	-	PRO	deletion	UNP Q9NP87
A	?	-	GLY	deletion	UNP Q9NP87
A	?	-	ALA	deletion	UNP Q9NP87
A	?	-	ALA	deletion	UNP Q9NP87
A	?	-	VAL	deletion	UNP Q9NP87
A	?	-	GLY	deletion	UNP Q9NP87
A	?	-	GLY	deletion	UNP Q9NP87
A	?	-	SER	deletion	UNP Q9NP87
A	?	-	THR	deletion	UNP Q9NP87
A	?	-	ARG	deletion	UNP Q9NP87
A	?	-	PRO	deletion	UNP Q9NP87
A	?	-	CYS	deletion	UNP Q9NP87
A	410	GLY	PRO	conflict	UNP Q9NP87
A	438	ASP	LYS	engineered mutation	UNP Q9NP87

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*GP*GP*CP*CP*TP*AP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	9	Total	C	N	O	P	0	2	0
			218	105	42	62	9			

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*GP*TP*AP*(8OG))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	5	Total	C	N	O	P	0	3	0
			143	69	27	41	6			


- Molecule 4 is a DNA chain called DNA (5'-D(P*GP*CP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	6	Total	Mn	0	0
			6	6		
5	T	1	Total	Mn	0	0
			1	1		
5	P	2	Total	Mn	0	1
			2	2		
5	D	1	Total	Mn	0	0
			1	1		

- Molecule 6 is 8-OXO-2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (three-letter code: 8DG) (formula: C₁₀H₁₆N₅O₁₄P₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	
			32	10	5	14	3	

- Molecule 7 is PYROPHOSPHATE (three-letter code: PPV) (formula: $\text{H}_4\text{O}_7\text{P}_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P		
			9	7	2	0	1

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 3 2 1	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 3 2 1	0	0

- Molecule 9 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total C O S 5 1 3 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	299	Total O 311 311	0	26
10	T	34	Total O 35 35	0	1
10	P	23	Total O 24 24	0	7
10	D	21	Total O 21 21	0	1

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed DNA/RNA polymerase mu




- Molecule 2: DNA (5'-D(*CP*GP*GP*CP*CP*TP*AP*CP*G)-3')



- Molecule 3: DNA (5'-D(*CP*GP*TP*AP*(8OG))-3')



- Molecule 4: DNA (5'-D(P*GP*CP*CP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.24Å 62.22Å 118.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.98 – 1.53 34.98 – 1.53	Depositor EDS
% Data completeness (in resolution range)	98.9 (34.98-1.53) 98.9 (34.98-1.53)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 1.53Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.160 , 0.180 0.168 , 0.187	Depositor DCC
R_{free} test set	3406 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3570	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 8OG, 8DG, EDO, EPE, PPV, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	1/2743 (0.0%)	0.80	8/3713 (0.2%)
2	T	2.01	4/244 (1.6%)	1.66	6/375 (1.6%)
3	P	1.70	0/133	1.47	4/201 (2.0%)
4	D	1.95	1/92 (1.1%)	1.38	1/138 (0.7%)
All	All	0.88	6/3212 (0.2%)	0.96	19/4427 (0.4%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	6	DT	P-O5'	-9.91	1.49	1.59
4	D	1	DG	OP3-P	-9.71	1.49	1.61
2	T	6	DT	C5-C7	-6.62	1.46	1.50
2	T	2[A]	DG	C3'-O3'	-5.71	1.36	1.44
2	T	2[B]	DG	C3'-O3'	-5.71	1.36	1.44
1	A	170	GLU	CG-CD	5.21	1.59	1.51

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	ARG	NE-CZ-NH2	-8.75	115.93	120.30
2	T	6	DT	N3-C4-O4	8.14	124.79	119.90
2	T	3	DG	OP1-P-OP2	7.88	131.41	119.60
3	P	4[A]	DA	O4'-C4'-C3'	-7.38	101.55	104.50
3	P	4[B]	DA	O4'-C4'-C3'	-7.38	101.55	104.50
2	T	9	DG	O4'-C1'-N9	-7.25	102.93	108.00
1	A	145	ARG	NE-CZ-NH1	7.02	123.81	120.30
1	A	175	ARG	NE-CZ-NH1	6.62	123.61	120.30
1	A	393	ARG	NE-CZ-NH1	6.20	123.40	120.30
4	D	1	DG	O4'-C1'-N9	-5.90	103.87	108.00
1	A	438	ASP	CB-CG-OD1	5.89	123.61	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	ASP	CB-CG-OD1	-5.80	113.08	118.30
1	A	438	ASP	CB-CG-OD2	-5.75	113.12	118.30
1	A	445	ARG	NE-CZ-NH2	-5.66	117.47	120.30
2	T	3	DG	O5'-P-OP2	-5.57	100.69	105.70
2	T	7	DA	O4'-C1'-C2'	5.44	110.25	105.90
2	T	6	DT	C5-C4-O4	-5.23	121.24	124.90
3	P	3[A]	DT	N3-C4-O4	5.04	122.92	119.90
3	P	3[B]	DT	N3-C4-O4	5.04	122.92	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2665	0	2581	16	0
2	T	218	0	125	1	0
3	P	143	0	81	4	0
4	D	83	0	45	0	0
5	A	6	0	0	0	0
5	D	1	0	0	0	0
5	P	2	0	0	0	0
5	T	1	0	0	0	0
6	A	32	0	11	3	0
7	A	9	0	0	0	0
8	A	14	0	18	3	0
9	A	5	0	0	0	0
10	A	311	0	0	4	0
10	D	21	0	0	0	0
10	P	24	0	0	1	0
10	T	35	0	0	0	0
All	All	3570	0	2861	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:4[A]:DA:H1'	10:P:212[A]:HOH:O	1.76	0.83
6:A:507[A]:8DG:O1A	10:A:601[A]:HOH:O	2.04	0.74
6:A:507[A]:8DG:H5'1	10:A:715[A]:HOH:O	1.90	0.71
1:A:458[A]:SER:H	8:A:510:EDO:H12	1.59	0.65
1:A:458[B]:SER:H	8:A:510:EDO:H12	1.61	0.64
2:T:1[B]:DC:H2'	2:T:2[B]:DG:C8	2.37	0.59
3:P:5[B]:8OG:O8	3:P:5[B]:8OG:H3'	2.11	0.50
1:A:452:LYS:HE3	1:A:482:LEU:O	2.10	0.50
1:A:226:GLU:OE2	1:A:229[B]:ARG:NH1	2.44	0.49
1:A:362:TYR:O	1:A:390[B]:CYS:HB2	2.13	0.49
1:A:390[B]:CYS:SG	1:A:392[B]:PHE:HE2	2.36	0.48
1:A:170:GLU:HG2	10:A:619[A]:HOH:O	2.17	0.45
1:A:195:THR:HG1	1:A:197:SER:HG	1.64	0.45
1:A:390[B]:CYS:SG	1:A:392[B]:PHE:CE2	3.11	0.44
1:A:351[B]:MET:SD	1:A:419:LEU:HD12	2.58	0.43
10:A:601[A]:HOH:O	3:P:4[A]:DA:H5'	2.19	0.43
1:A:434:TRP:O	6:A:507[A]:8DG:H4'	2.20	0.42
1:A:436:GLY:CA	3:P:5[B]:8OG:H2'	2.50	0.42
1:A:266:ARG:O	1:A:269:PRO:HD3	2.19	0.41
1:A:420:VAL:HG21	1:A:434:TRP:CZ2	2.55	0.41
1:A:457:ASN:HB2	8:A:510:EDO:H12	2.02	0.41
1:A:138:MET:SD	1:A:145:ARG:HD2	2.61	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/356 (96%)	334 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/301 (90%)	272 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	8OG	P	5[B]	5,2,3	18,25,26	1.52	3 (16%)	21,37,40	2.96	9 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	8OG	P	5[B]	5,2,3	-	0/3/21/22	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	5[B]	8OG	C6-N1	4.42	1.40	1.33
3	P	5[B]	8OG	C8-N7	3.19	1.37	1.34
3	P	5[B]	8OG	C2-N1	2.32	1.39	1.35

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	5[B]	8OG	C5-C6-N1	-9.16	110.90	123.43
3	P	5[B]	8OG	C2-N1-C6	6.06	125.56	115.93
3	P	5[B]	8OG	C2-N3-C4	-4.23	110.53	115.36
3	P	5[B]	8OG	C2'-C1'-N9	-2.93	112.82	116.01
3	P	5[B]	8OG	N3-C2-N1	-2.91	123.34	127.22
3	P	5[B]	8OG	O4'-C4'-C3'	-2.57	99.68	105.67
3	P	5[B]	8OG	C4-C5-N7	2.19	111.70	109.47
3	P	5[B]	8OG	C1'-N9-C4	-2.15	123.00	126.99
3	P	5[B]	8OG	C4-C5-C6	-2.02	118.86	120.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	P	5[B]	8OG	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 10 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	EDO	A	509	-	3,3,3	0.46	0	2,2,2	0.57	0
8	EDO	A	511	-	3,3,3	0.48	0	2,2,2	0.28	0
9	EPE	A	513	-	4,4,15	0.91	0	5,6,20	1.79	2 (40%)
6	8DG	A	507[A]	5	26,34,34	1.20	3 (11%)	34,54,54	2.25	7 (20%)
8	EDO	A	510	-	2,2,3	0.78	0	1,1,2	1.05	0
7	PPV	A	508[B]	5	6,8,8	0.83	0	13,13,13	2.19	3 (23%)
8	EDO	A	512	-	2,2,3	0.52	0	1,1,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	EDO	A	509	-	-	0/1/1/1	-
6	8DG	A	507[A]	5	-	5/18/34/34	0/3/3/3
8	EDO	A	511	-	-	1/1/1/1	-
7	PPV	A	508[B]	5	-	0/6/6/6	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	507[A]	8DG	C6-N1	3.72	1.39	1.33
6	A	507[A]	8DG	C8-N7	3.56	1.38	1.34
6	A	507[A]	8DG	C2-N1	2.06	1.39	1.35

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	507[A]	8DG	C5-C6-N1	-8.57	111.72	123.43
6	A	507[A]	8DG	C2-N1-C6	5.40	124.52	115.93
7	A	508[B]	PPV	P2-OPP-P1	-4.93	115.92	132.83
6	A	507[A]	8DG	C2-N3-C4	-4.05	110.73	115.36
7	A	508[B]	PPV	O32-P2-O12	3.46	120.87	107.64
6	A	507[A]	8DG	PA-O3A-PB	-3.40	121.16	132.83
9	A	513	EPE	O2S-S-O1S	-3.18	109.44	118.02
7	A	508[B]	PPV	O21-P1-O11	3.12	119.57	107.64
6	A	507[A]	8DG	O2G-PG-O3B	2.46	112.87	104.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	507[A]	8DG	C2'-C1'-N9	-2.45	113.34	116.01
6	A	507[A]	8DG	C4-C5-C6	-2.22	118.67	120.80
9	A	513	EPE	O2S-S-C10	2.05	111.82	107.54

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	507[A]	8DG	PB-O3B-PG-O2G
8	A	511	EDO	O1-C1-C2-O2
6	A	507[A]	8DG	PA-O3A-PB-O2B
6	A	507[A]	8DG	PB-O3B-PG-O3G
6	A	507[A]	8DG	C5'-O5'-PA-O3A
6	A	507[A]	8DG	PB-O3A-PA-O2A

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	507[A]	8DG	3	0
8	A	510	EDO	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/356 (91%)	-0.13	2 (0%) 89 91	14, 24, 41, 61	0
2	T	9/9 (100%)	-0.49	0 100 100	20, 24, 27, 28	0
3	P	4/5 (80%)	0.24	0 100 100	18, 18, 19, 21	0
4	D	4/4 (100%)	-0.78	0 100 100	21, 21, 24, 27	0
All	All	344/374 (91%)	-0.15	2 (0%) 89 91	14, 24, 40, 61	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	137	TRP	9.9
1	A	384	ALA	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	8OG	P	5[B]	23/24	0.95	0.11	16,29,35,37	23

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MN	A	503	1/1	0.79	0.11	61,61,61,61	1
5	MN	T	101	1/1	0.79	0.22	75,75,75,75	1
8	EDO	A	511	4/4	0.79	0.15	40,43,45,49	4
8	EDO	A	512	3/4	0.82	0.11	49,49,51,51	0
5	MN	D	101	1/1	0.91	0.21	54,54,54,54	1
5	MN	A	505	1/1	0.92	0.08	29,29,29,29	1
6	8DG	A	507[A]	32/32	0.96	0.10	10,17,26,27	32
8	EDO	A	510	3/4	0.96	0.15	16,16,24,31	0
8	EDO	A	509	4/4	0.97	0.18	35,38,38,41	0
7	PPV	A	508[B]	9/9	0.97	0.09	15,18,26,28	9
9	EPE	A	513	5/15	0.98	0.08	29,34,38,39	5
5	MN	A	502	1/1	0.99	0.04	25,25,25,25	1
5	MN	A	504	1/1	0.99	0.10	31,31,31,31	1
5	MN	P	101[B]	1/1	0.99	0.03	19,19,19,19	1
5	MN	P	102	1/1	0.99	0.06	17,17,17,17	0
5	MN	A	506	1/1	1.00	0.07	18,18,18,18	1
5	MN	A	501	1/1	1.00	0.11	15,15,15,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MN A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN T 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN D 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN A 505:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 8DG A 507 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PPV A 508 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN A 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN P 101 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN P 102:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN A 506:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MN A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.