



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 03:11 AM EDT

PDB ID : 1KVL  
Title : X-ray Crystal Structure of AmpC S64G Mutant beta-Lactamase in Complex with Substrate and Product Forms of Cephalothin  
Authors : Beadle, B.M.; Trehan, I.; Focia, P.J.; Shoichet, B.K.  
Deposited on : 2002-01-27  
Resolution : 1.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2556 (1.56-1.52)
Clashscore	141614	2634 (1.56-1.52)
Ramachandran outliers	138981	2580 (1.56-1.52)
Sidechain outliers	138945	2577 (1.56-1.52)
RSRZ outliers	127900	2524 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	
1	B	358	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	THN	B	372	X	-	-	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6146 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	358	Total	C	N	O	S	0	1	0
			2781	1795	473	507	6			
1	B	358	Total	C	N	O	S	0	0	0
			2772	1788	472	506	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	64	GLY	SER	engineered mutation	UNP P00811
B	64	GLY	SER	engineered mutation	UNP P00811

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	1
			5	4	1		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is 2-[CARBOXY-(2-THIOPHEN-2-YL-ACETYLAMINO)-METHYL]-5-METHYL-3,6-DIHYDRO-2H-[1,3]THIAZINE-4-CARBOXYLIC ACID (three-letter code: KCP) (formula:  $C_{14}H_{16}N_2O_5S_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	1
			46	28	4	10	4		

- Molecule 4 is CEPHALOTHIN (three-letter code: CLS) (formula:  $C_{16}H_{16}N_2O_6S_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	1
			26	16	2	6	2		

- Molecule 5 is 2-[CARBOXY-(2-THIOPHEN-2-YL-ACETYLAMINO)-METHYL]-5-METHYLENE-5,6-DIHYDRO-2H-[1,3]THIAZINE-4-CARBOXYLIC ACID (three-letter code: THN) (formula:  $C_{14}H_{14}N_2O_5S_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			23	14	2	5	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	270	Total 270	O 270	0	0
6	B	218	Total 218	O 218	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Beta-lactamase



#### • Molecule 1: Beta-lactamase





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.62Å 77.15Å 98.07Å 90.00° 115.90° 90.00°	Depositor
Resolution (Å)	20.00 – 1.53 19.76 – 1.53	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.53) 82.4 (19.76-1.53)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 1.53Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.195 , 0.220 0.193 , 0.216	Depositor DCC
$R_{free}$ test set	2463 reflections (2.50%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.4	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6146	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCP, THN, CLS, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.71	3/2861 (0.1%)	0.83	4/3912 (0.1%)
1	B	0.60	0/2852	0.77	2/3900 (0.1%)
All	All	0.66	3/5713 (0.1%)	0.80	6/7812 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	164[A]	LYS	C-N	6.27	1.46	1.34
1	A	164[B]	LYS	C-N	6.27	1.46	1.34
1	A	165	PRO	N-CD	6.18	1.56	1.47

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	165	PRO	CA-N-CD	-6.57	102.30	111.50
1	A	164[A]	LYS	C-N-CD	5.83	140.64	128.40
1	A	164[B]	LYS	C-N-CD	5.83	140.64	128.40
1	B	217	ASP	CB-CG-OD1	5.32	123.09	118.30
1	A	312	TRP	N-CA-C	-5.24	96.84	111.00
1	B	121	VAL	N-CA-C	-5.14	97.13	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	164[A]	LYS	Mainchain
1	A	164[B]	LYS	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2781	0	2737	36	0
1	B	2772	0	2729	25	0
2	A	10	0	0	1	0
3	A	46	0	24	2	0
4	A	26	0	15	1	0
5	B	23	0	11	0	0
6	A	270	0	0	1	0
6	B	218	0	0	0	0
All	All	6146	0	5516	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:94:PRO:HG2	1:B:95:GLU:OE1	1.71	0.90
1:A:279:ASN:HD22	1:A:280:PRO:HD2	1.43	0.82
1:B:36:GLY:N	1:B:333:GLU:HG2	2.03	0.73
1:A:7:GLN:O	1:A:11:ILE:HG22	1.89	0.72
1:A:172:GLN:HE21	1:A:176:THR:HG23	1.56	0.70
1:B:262:THR:CG2	1:B:298:VAL:HG12	2.22	0.68
1:A:17:THR:HB	1:A:18:PRO:HD3	1.76	0.67
1:A:172:GLN:HE21	1:A:176:THR:CG2	2.11	0.64
1:A:334:LEU:HG	1:A:357:LEU:HD22	1.79	0.64
4:A:371[A]:CLS:H233	4:A:371[A]:CLS:H22	1.78	0.64
1:A:128:SER:HB3	3:A:380[B]:KCP:O9A	1.99	0.62
1:B:13:HIS:HD2	1:B:17:THR:OG1	1.82	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:PRO:O	1:A:249:GLN:HG3	2.03	0.58
1:A:279:ASN:HD22	1:A:280:PRO:CD	2.14	0.58
1:B:238:LEU:HD12	1:B:330:PRO:HA	1.86	0.58
1:B:262:THR:HG22	1:B:298:VAL:HG12	1.86	0.57
1:A:182:LEU:O	1:A:232:ARG:HD3	2.05	0.56
1:B:334:LEU:HG	1:B:357:LEU:HD22	1.86	0.56
1:B:171:GLU:O	1:B:175:GLN:HG3	2.05	0.55
1:B:94:PRO:CG	1:B:95:GLU:OE1	2.51	0.54
1:A:243:ILE:HD13	1:A:252:ILE:HD12	1.89	0.54
1:A:274:LEU:HD22	1:A:283:ILE:HG22	1.90	0.53
1:B:243:ILE:CD1	1:B:252:ILE:HD12	2.39	0.53
1:A:240:PRO:HA	1:A:243:ILE:HD12	1.90	0.52
1:B:56:GLN:HB3	1:B:228:GLU:HG3	1.91	0.51
1:A:107:LEU:HD22	1:B:303:PRO:HD3	1.93	0.49
1:A:344:TYR:CZ	1:A:349:ARG:HG2	2.48	0.49
1:A:360:LEU:O	1:A:361:GLN:CB	2.60	0.48
1:A:278:VAL:HG21	1:A:283:ILE:CG2	2.43	0.48
1:A:283:ILE:HG13	1:A:284:ILE:N	2.28	0.48
1:B:122:PRO:HG2	1:B:125:VAL:HG23	1.94	0.48
1:A:185:ASN:O	2:A:901:PO4:O1	2.32	0.48
1:B:344:TYR:CE2	1:B:349:ARG:HG2	2.48	0.48
1:B:56:GLN:HB3	1:B:228:GLU:CG	2.44	0.47
1:B:138:TRP:CH2	1:B:140:PRO:HB3	2.48	0.47
1:A:288:ASP:OD1	1:A:290:LYS:HB3	2.16	0.46
1:A:278:VAL:HG21	1:A:283:ILE:HG23	1.97	0.45
1:A:276:TRP:CD2	1:A:277:PRO:HA	2.51	0.45
1:B:17:THR:HB	1:B:18:PRO:HD3	1.97	0.45
1:A:119:LEU:HA	1:A:151:ALA:HA	1.98	0.45
1:A:161:LEU:HD23	1:A:164[B]:LYS:HE3	1.98	0.45
1:B:344:TYR:CZ	1:B:349:ARG:HG2	2.52	0.45
1:A:316:THR:HG22	1:A:325:TYR:CD1	2.52	0.44
1:A:8:ILE:HA	1:A:11:ILE:CG2	2.47	0.44
1:A:262:THR:HG23	1:A:262:THR:O	2.17	0.44
1:A:344:TYR:CE2	1:A:349:ARG:HG2	2.53	0.44
1:A:34:TYR:CE1	1:A:35:GLN:NE2	2.85	0.44
1:A:289:ASN:C	1:A:291:ILE:N	2.71	0.43
1:A:11:ILE:HG23	1:A:12:VAL:N	2.33	0.43
1:B:358:ASN:O	1:B:361:GLN:HG3	2.19	0.43
1:B:37:LYS:HD3	1:B:39:TYR:CZ	2.54	0.42
1:B:231:ALA:O	1:B:235:GLN:HG3	2.19	0.42
1:A:93:TRP:CD2	3:A:380[A]:KCP:H3'3	2.55	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:87:ASP:OD1	1:B:92:TYR:OH	2.30	0.42
1:A:360:LEU:O	1:A:361:GLN:CG	2.68	0.42
1:B:37:LYS:HD3	1:B:39:TYR:OH	2.19	0.42
1:B:164:LYS:HB2	1:B:165:PRO:HD3	2.01	0.41
1:A:360:LEU:O	1:A:361:GLN:HB2	2.19	0.41
1:B:250:GLN:O	1:B:254:LEU:HG	2.20	0.41
1:A:280:PRO:HA	1:A:283:ILE:HG12	2.03	0.40
1:A:309:ARG:NH1	6:A:773:HOH:O	2.53	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/358 (100%)	349 (98%)	8 (2%)	0	100	100
1	B	356/358 (99%)	348 (98%)	8 (2%)	0	100	100
All	All	713/716 (100%)	697 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/291 (98%)	281 (99%)	3 (1%)	73	51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	284/291 (98%)	279 (98%)	5 (2%)	59	29
All	All	568/582 (98%)	560 (99%)	8 (1%)	67	39

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	123	ASP
1	A	279	ASN
1	A	314	HIS
1	B	6	GLN
1	B	124	GLU
1	B	238	LEU
1	B	242	ASP
1	B	314	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	22	GLN
1	A	23	GLN
1	A	35	GLN
1	A	137	ASN
1	A	172	GLN
1	A	198	ASN
1	A	235	GLN
1	A	249	GLN
1	A	279	ASN
1	B	6	GLN
1	B	13	HIS
1	B	22	GLN
1	B	52	GLN
1	B	137	ASN
1	B	198	ASN
1	B	249	GLN
1	B	358	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	KCP	A	380[A]	1	16,24,24	5.56	7 (43%)	13,33,33	2.30	6 (46%)
3	KCP	A	380[B]	1	16,24,24	5.65	7 (43%)	13,33,33	2.28	6 (46%)
4	CLS	A	371[A]	-	25,28,28	2.33	9 (36%)	28,40,40	1.29	3 (10%)
2	PO4	A	901	-	4,4,4	0.73	0	6,6,6	0.52	0
5	THN	B	372	-	13,24,24	2.46	4 (30%)	10,33,33	1.76	4 (40%)
2	PO4	A	902[B]	-	4,4,4	0.88	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	THN	B	372	-	1/1/7/8	0/6/33/33	0/1/2/2
3	KCP	A	380[B]	1	-	2/6/33/33	0/1/2/2
4	CLS	A	371[A]	-	-	3/11/46/46	0/3/3/3
3	KCP	A	380[A]	1	-	0/6/33/33	0/1/2/2

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	380[B]	KCP	C3-C4	19.02	1.58	1.34
3	A	380[A]	KCP	C3-C4	18.63	1.58	1.34
3	A	380[B]	KCP	C3'-C3	-7.50	1.38	1.50
3	A	380[A]	KCP	C3'-C3	-7.25	1.39	1.50
4	A	371[A]	CLS	C13-C14	6.03	1.56	1.51
5	B	372	THN	C2-S1	5.54	1.88	1.81
3	A	380[A]	KCP	C2-C3	5.39	1.58	1.51
3	A	380[B]	KCP	C14-S19	-5.37	1.62	1.73
3	A	380[B]	KCP	C2-C3	5.20	1.58	1.51
5	B	372	THN	C14-S19	-5.16	1.63	1.73
4	A	371[A]	CLS	C7-N10	-4.80	1.35	1.45
3	A	380[A]	KCP	C14-S19	-4.78	1.63	1.73
4	A	371[A]	CLS	C6-N5	4.22	1.55	1.47
3	A	380[A]	KCP	C13-C14	-4.01	1.48	1.51
4	A	371[A]	CLS	C13-C11	-3.51	1.44	1.51
3	A	380[A]	KCP	C2-S1	3.15	1.89	1.82
3	A	380[B]	KCP	C2-S1	3.14	1.89	1.82
4	A	371[A]	CLS	C15-C14	2.98	1.45	1.37
5	B	372	THN	C3'-C3	2.94	1.39	1.32
3	A	380[B]	KCP	C4'-C4	2.79	1.56	1.52
3	A	380[B]	KCP	C4-N5	2.61	1.42	1.33
4	A	371[A]	CLS	C3'-C3	2.46	1.54	1.50
4	A	371[A]	CLS	C14-S19	-2.21	1.69	1.73
4	A	371[A]	CLS	C16-C15	2.06	1.46	1.39
5	B	372	THN	C16-C15	2.03	1.46	1.39
3	A	380[A]	KCP	C4-N5	2.01	1.40	1.33
4	A	371[A]	CLS	C6-C7	2.01	1.60	1.56

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	380[A]	KCP	C3'-C3-C2	3.70	119.23	113.23
3	A	380[B]	KCP	C16-C17-S19	-3.68	110.00	112.98
5	B	372	THN	C16-C17-S19	-3.56	110.09	112.98
3	A	380[A]	KCP	C6-C7-N10	3.43	117.54	110.02
3	A	380[B]	KCP	C14-C13-C11	3.37	120.09	113.39
3	A	380[B]	KCP	C3'-C3-C2	3.09	118.24	113.23
3	A	380[A]	KCP	C16-C17-S19	-3.09	110.48	112.98
3	A	380[B]	KCP	C2-S1-C6	2.99	100.33	94.47
3	A	380[A]	KCP	C3-C2-S1	-2.90	112.86	116.98
3	A	380[A]	KCP	C14-C13-C11	2.86	119.08	113.39
3	A	380[B]	KCP	C3'-C3-C4	-2.78	120.07	123.53
3	A	380[A]	KCP	C3'-C3-C4	-2.73	120.13	123.53

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	371[A]	CLS	C3-C2-S1	-2.66	110.15	115.08
3	A	380[B]	KCP	C6-N5-C4	2.60	124.45	118.32
5	B	372	THN	C2-C3-C3'	-2.43	119.34	123.78
4	A	371[A]	CLS	C7-C6-S1	-2.43	111.99	116.51
4	A	371[A]	CLS	C6-N5-C4	-2.27	118.78	125.35
5	B	372	THN	C8-C7-N10	-2.27	105.61	110.33
5	B	372	THN	C14-C13-C11	2.05	117.48	113.39

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	B	372	THN	C4

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	380[B]	KCP	C6-C7-N10-C11
4	A	371[A]	CLS	C23-C21-O20-C3'
4	A	371[A]	CLS	O22-C21-O20-C3'
3	A	380[B]	KCP	C8-C7-N10-C11
4	A	371[A]	CLS	C4-C3-C3'-O20



There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	380[A]	KCP	1	0
3	A	380[B]	KCP	1	0
4	A	371[A]	CLS	1	0
2	A	901	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	358/358 (100%)	0.44	27 (7%)	14 16	16, 24, 43, 61	0
1	B	358/358 (100%)	0.28	19 (5%)	26 30	18, 26, 41, 47	0
All	All	716/716 (100%)	0.36	46 (6%)	19 22	16, 25, 42, 61	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	284	ILE	7.7
1	A	285	ASN	7.0
1	A	4	ALA	5.3
1	A	288	ASP	4.8
1	A	34	TYR	4.7
1	A	289	ASN	4.6
1	B	4	ALA	4.5
1	B	244	ASN	4.5
1	A	279	ASN	4.4
1	A	281	ASP	4.3
1	B	242	ASP	4.2
1	A	287	SER	4.2
1	A	283	ILE	4.2
1	A	361	GLN	4.0
1	A	290	LYS	3.8
1	B	49	ALA	3.7
1	B	128	SER	3.5
1	A	35	GLN	3.5
1	B	127	SER	3.5
1	A	244	ASN	3.3
1	A	205	GLU	3.2
1	A	280	PRO	3.2
1	B	5	PRO	3.0
1	B	102	ASN	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	241	LEU	2.7
1	A	6	GLN	2.7
1	A	7	GLN	2.7
1	B	206	GLY	2.6
1	A	286	GLY	2.5
1	A	123	ASP	2.5
1	A	278	VAL	2.4
1	A	291	ILE	2.4
1	B	52	GLN	2.3
1	A	22	GLN	2.3
1	B	246	LYS	2.3
1	B	204	ARG	2.3
1	A	102	ASN	2.2
1	A	336	ILE	2.2
1	B	245	GLU	2.2
1	B	207	LYS	2.2
1	A	322	PHE	2.2
1	B	94	PRO	2.2
1	B	124	GLU	2.2
1	A	126	LYS	2.1
1	B	6	GLN	2.1
1	B	129	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CLS	A	371[A]	26/26	0.64	0.32	34,44,64,64	26

*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	A	901	5/5	0.69	0.20	42,48,50,53	0
3	KCP	A	380[B]	23/23	0.73	0.26	36,41,43,44	23
3	KCP	A	380[A]	23/23	0.73	0.26	19,29,35,35	23
2	PO4	A	902[B]	5/5	0.78	0.20	35,35,37,37	5
5	THN	B	372	23/23	0.84	0.16	29,42,49,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around CLS A 371 (A):**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around KCP A 380 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around KCP A 380 (A):**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around THN B 372:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.