



# Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 06:52 am BST

PDB ID : 6KVJ  
Title : Crystal structure of UDPX-SrUGT76G1  
Authors : Li, J.X.; Liu, Z.F.; Wang, Y.; Zhang, P.  
Deposited on : 2019-09-04  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	<div> <div>5%</div> <div> <div></div> <div>68%</div> <div>25%</div> <div>...</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called UDP-glycosyltransferase 76G1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	448	3578	2295	601	671	11	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q6VAB4
A	460	THR	-	expression tag	UNP Q6VAB4
A	461	SER	-	expression tag	UNP Q6VAB4

- Molecule 2 is URIDINE-5'-DIPHOSPHATE-XYLOPYRANOSE (three-letter code: UDX) (formula: C<sub>14</sub>H<sub>22</sub>N<sub>2</sub>O<sub>16</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	34	14	2	16	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	12	Total	O	0	0
			12	12		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-glycosyltransferase 76G1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.24Å 98.24Å 90.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.12 – 2.50 49.12 – 2.50	Depositor EDS
% Data completeness (in resolution range)	71.1 (49.12-2.50) 46.7 (49.12-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.64 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, $R_{free}$	0.181 , 0.230 0.182 , 0.214	Depositor DCC
$R_{free}$ test set	432 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.8	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.038 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3624	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: UDX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	1/3667 (0.0%)	0.83	13/4976 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	21	VAL	C-N	5.15	1.44	1.34

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	GLU	CB-CA-C	-16.27	77.87	110.40
1	A	327	ASP	CB-CA-C	-16.18	78.03	110.40
1	A	327	ASP	N-CA-C	12.16	143.82	111.00
1	A	328	GLY	N-CA-C	-10.81	86.09	113.10
1	A	318	GLY	N-CA-C	-7.33	94.77	113.10
1	A	322	VAL	N-CA-C	6.75	129.21	111.00
1	A	332	GLU	N-CA-C	6.33	128.09	111.00
1	A	321	TRP	N-CA-C	-6.16	94.36	111.00
1	A	68	LEU	CB-CA-C	-5.47	99.80	110.20
1	A	322	VAL	N-CA-CB	-5.42	99.57	111.50
1	A	179	GLN	N-CA-C	5.38	125.52	111.00
1	A	203	GLU	OE1-CD-OE2	-5.37	116.86	123.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	86	LEU	CB-CG-CD2	5.05	119.59	111.00

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	178	GLU	Peptide
1	A	180	ALA	Peptide
1	A	326	PRO	Peptide
1	A	327	ASP	Peptide
1	A	330	LEU	Peptide
1	A	70	ASN	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3578	0	3528	113	0
2	A	34	0	20	3	0
3	A	12	0	0	1	0
All	All	3624	0	3548	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:GLY:O	1:A:332:GLU:HB2	1.36	1.11
1:A:330:LEU:HD13	1:A:330:LEU:H	1.33	0.94
1:A:289:ASP:OD2	1:A:292:ASP:OD2	1.92	0.87
1:A:178:GLU:OE2	1:A:178:GLU:N	2.08	0.87
1:A:418:ASP:O	1:A:425:ARG:NH2	2.11	0.84
1:A:177:GLU:C	1:A:178:GLU:OE2	2.16	0.83
1:A:333:ARG:HH11	1:A:333:ARG:HG3	1.42	0.82

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:422:GLU:OE1	1:A:425:ARG:NH1	2.13	0.81
1:A:71:ASP:H	1:A:72:PRO:HD2	1.46	0.81
1:A:331:GLY:O	1:A:332:GLU:CB	2.23	0.80
1:A:71:ASP:N	1:A:72:PRO:HD2	1.96	0.78
1:A:203:GLU:O	1:A:207:LYS:HG3	1.84	0.77
1:A:208:MET:O	1:A:212:THR:HG23	1.86	0.76
1:A:333:ARG:NH1	1:A:333:ARG:HG3	1.97	0.75
1:A:174:THR:HG23	1:A:175:ARG:HD3	1.68	0.74
1:A:161:GLN:O	1:A:165:LEU:HD12	1.89	0.72
1:A:262:ASP:OD1	1:A:264:THR:OG1	2.07	0.72
1:A:295:GLU:OE1	1:A:298:ARG:NH1	2.19	0.72
1:A:298:ARG:HG2	1:A:329:PHE:HZ	1.54	0.72
1:A:321:TRP:O	1:A:323:GLU:OE1	2.09	0.71
1:A:199:GLN:HA	1:A:202:LYS:HE3	1.73	0.70
1:A:282:PHE:O	1:A:286:SER:OG	2.08	0.70
1:A:168:LEU:HD12	1:A:195:TYR:HB2	1.74	0.70
1:A:330:LEU:CD1	1:A:330:LEU:H	2.06	0.66
1:A:323:GLU:N	1:A:323:GLU:OE1	2.25	0.66
1:A:51:PHE:N	1:A:69:ASP:OD1	2.27	0.65
1:A:156:HIS:HB3	1:A:209:ILE:HD11	1.81	0.62
1:A:198:TRP:CZ2	1:A:202:LYS:HD3	2.35	0.62
1:A:414:ARG:HA	1:A:418:ASP:HB2	1.82	0.62
1:A:20:PRO:HB3	1:A:30:ILE:HG21	1.83	0.61
1:A:81:PRO:HD2	1:A:86:LEU:HD13	1.83	0.60
1:A:330:LEU:N	1:A:330:LEU:HD13	2.10	0.60
1:A:224:LYS:HG3	1:A:231:LEU:HD11	1.84	0.59
1:A:71:ASP:N	1:A:72:PRO:CD	2.66	0.58
1:A:323:GLU:N	1:A:324:PRO:HD2	2.19	0.58
1:A:197:ASN:O	1:A:201:LEU:HG	2.04	0.57
2:A:801:UDX:H1'	2:A:801:UDX:O2A	2.05	0.57
1:A:329:PHE:N	1:A:330:LEU:HD13	2.19	0.57
1:A:161:GLN:HA	1:A:164:GLU:HG2	1.87	0.57
1:A:204:ILE:HD12	1:A:204:ILE:H	1.71	0.56
1:A:50:ASN:H	1:A:69:ASP:HB3	1.71	0.55
1:A:169:ASP:O	1:A:192:LYS:NZ	2.36	0.54
1:A:20:PRO:HB2	1:A:27:ILE:HG22	1.88	0.54
1:A:420:GLU:OE1	1:A:420:GLU:N	2.34	0.53
1:A:132:GLN:HB2	1:A:142:ARG:NH1	2.24	0.53
1:A:199:GLN:HA	1:A:202:LYS:CE	2.40	0.52
1:A:166:GLY:HA3	1:A:173:LYS:HD2	1.92	0.51
1:A:333:ARG:N	1:A:333:ARG:CD	2.74	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:TYR:O	1:A:192:LYS:NZ	2.37	0.51
1:A:213:LYS:HD3	1:A:237:GLU:O	2.11	0.50
1:A:419:GLU:H	1:A:419:GLU:CD	2.14	0.50
1:A:289:ASP:O	1:A:292:ASP:N	2.45	0.50
1:A:289:ASP:OD2	1:A:292:ASP:CG	2.50	0.50
1:A:199:GLN:HG3	1:A:199:GLN:O	2.11	0.50
1:A:330:LEU:N	1:A:330:LEU:CD1	2.72	0.49
1:A:329:PHE:C	1:A:329:PHE:CD1	2.85	0.49
1:A:40:LYS:HG3	1:A:453:VAL:HG21	1.94	0.49
1:A:23:PHE:HE1	1:A:91:ILE:HD11	1.75	0.48
1:A:89:MET:O	1:A:92:PRO:HD2	2.14	0.48
1:A:17:ILE:HD11	1:A:109:LEU:HD22	1.96	0.48
1:A:324:PRO:O	1:A:326:PRO:HD3	2.13	0.48
1:A:159:LEU:HD21	1:A:195:TYR:CD2	2.49	0.47
1:A:197:ASN:CG	1:A:200:ILE:HD12	2.35	0.47
1:A:259:LEU:HD12	1:A:339:TRP:CE3	2.49	0.47
1:A:197:ASN:ND2	1:A:200:ILE:HD12	2.30	0.47
1:A:196:SER:N	3:A:902:HOH:O	2.47	0.47
1:A:408:ILE:O	1:A:412:ILE:HG13	2.15	0.47
1:A:303:SER:O	1:A:305:GLN:NE2	2.44	0.47
1:A:81:PRO:HD2	1:A:86:LEU:CD1	2.45	0.47
1:A:333:ARG:HH11	1:A:333:ARG:CG	2.17	0.46
1:A:89:MET:O	1:A:93:ILE:HG13	2.15	0.46
1:A:198:TRP:CG	1:A:202:LYS:NZ	2.84	0.45
1:A:429:ARG:O	1:A:433:GLN:HG3	2.16	0.45
1:A:233:THR:O	1:A:237:GLU:HG2	2.15	0.45
1:A:325:LEU:HA	1:A:325:LEU:HD23	1.74	0.45
1:A:332:GLU:HB3	1:A:333:ARG:HD2	1.98	0.45
1:A:329:PHE:CD1	1:A:329:PHE:O	2.70	0.45
1:A:197:ASN:OD1	1:A:200:ILE:HD12	2.17	0.45
1:A:45:THR:HG21	1:A:109:LEU:CD1	2.47	0.45
1:A:289:ASP:OD2	1:A:292:ASP:HB2	2.16	0.44
1:A:48:HIS:O	1:A:66:PHE:HA	2.17	0.44
1:A:168:LEU:HD11	1:A:195:TYR:HD2	1.82	0.44
1:A:202:LYS:HB2	1:A:202:LYS:HE3	1.40	0.44
1:A:86:LEU:HD12	1:A:87:ALA:N	2.33	0.44
1:A:177:GLU:HA	1:A:188:VAL:HB	1.99	0.44
1:A:399:LEU:HD21	1:A:411:ALA:HB2	2.00	0.44
1:A:259:LEU:HD12	1:A:339:TRP:CD2	2.53	0.43
1:A:342:GLN:NE2	2:A:801:UDX:H2A1	2.33	0.43
1:A:159:LEU:HD22	1:A:198:TRP:CH2	2.52	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:TRP:CH2	1:A:202:LYS:HD3	2.53	0.43
1:A:22:PRO:HB3	1:A:48:HIS:CE1	2.53	0.43
1:A:178:GLU:HG3	1:A:187:LYS:HD2	2.01	0.43
1:A:301:VAL:HG21	1:A:329:PHE:CE1	2.53	0.43
1:A:73:GLN:HG2	1:A:77:ILE:HD12	2.00	0.43
1:A:91:ILE:HB	1:A:92:PRO:HD3	1.99	0.43
1:A:18:LEU:HA	1:A:123:ILE:O	2.18	0.43
2:A:801:UDX:H1'	2:A:801:UDX:PA	2.59	0.42
1:A:328:GLY:C	1:A:330:LEU:HD13	2.38	0.42
1:A:231:LEU:HD23	1:A:231:LEU:HA	1.79	0.42
1:A:104:ARG:O	1:A:108:LEU:HG	2.19	0.42
1:A:77:ILE:HG22	1:A:80:LEU:HD12	2.01	0.42
1:A:174:THR:HG23	1:A:175:ARG:HH11	1.83	0.42
1:A:288:VAL:CG1	1:A:293:PHE:HB2	2.50	0.41
1:A:389:MET:SD	1:A:393:LEU:HD23	2.59	0.41
1:A:174:THR:OG1	1:A:175:ARG:N	2.53	0.41
1:A:200:ILE:O	1:A:204:ILE:HD12	2.19	0.41
1:A:199:GLN:NE2	1:A:203:GLU:OE1	2.53	0.41
1:A:278:LEU:HD22	1:A:412:ILE:HD13	2.02	0.41
1:A:88:GLY:O	1:A:91:ILE:HG12	2.21	0.41
1:A:176:LEU:HA	1:A:176:LEU:HD22	1.87	0.41
1:A:20:PRO:HD2	1:A:47:PHE:O	2.21	0.41
1:A:140:LEU:HD23	1:A:141:ARG:O	2.21	0.41
1:A:132:GLN:OE1	1:A:142:ARG:HD3	2.21	0.40
1:A:159:LEU:HA	1:A:159:LEU:HD23	1.89	0.40
1:A:327:ASP:HB2	1:A:328:GLY:H	1.41	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/461 (97%)	418 (94%)	24 (5%)	4 (1%)	17	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	290	GLU
1	A	322	VAL
1	A	68	LEU
1	A	326	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	397/412 (96%)	377 (95%)	20 (5%)	24	46

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	GLU
1	A	113	SER
1	A	152	ASN
1	A	176	LEU
1	A	183	PHE
1	A	212	THR
1	A	262	ASP
1	A	285	THR
1	A	288	VAL
1	A	306	SER
1	A	320	THR
1	A	325	LEU
1	A	327	ASP
1	A	330	LEU
1	A	332	GLU
1	A	333	ARG
1	A	335	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	393	LEU
1	A	448	SER
1	A	454	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	ASN
1	A	261	HIS
1	A	271	GLN
1	A	343	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	UDX	A	801	-	29,36,36	1.71	8 (27%)	38,55,55	1.46	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDX	A	801	-	-	4/19/54/54	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	UDX	C2-N3	-3.69	1.30	1.38
2	A	801	UDX	C6-N1	-3.63	1.31	1.35
2	A	801	UDX	C2D-C1D	-2.90	1.49	1.53
2	A	801	UDX	PB-O2B	-2.72	1.42	1.55
2	A	801	UDX	PA-O2A	-2.58	1.43	1.55
2	A	801	UDX	PA-O1A	-2.28	1.42	1.50
2	A	801	UDX	C4'-C3'	-2.09	1.49	1.52
2	A	801	UDX	PB-O3B	-2.07	1.55	1.60

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	UDX	O5'-C1'-C2'	-3.79	104.22	110.04
2	A	801	UDX	O4'-C4'-C3'	-3.10	103.93	110.14
2	A	801	UDX	O3B-C1'-C2'	2.83	113.57	108.38
2	A	801	UDX	O2'-C2'-C1'	-2.53	103.89	110.05

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	UDX	C1'-O3B-PB-O3A
2	A	801	UDX	C2'-C1'-O3B-PB
2	A	801	UDX	PB-O3A-PA-O5D
2	A	801	UDX	O4D-C4D-C5D-O5D

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	UDX	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	448/461 (97%)	0.08	25 (5%) <span style="border: 1px solid red; padding: 0 2px;">24</span> <span style="border: 1px solid red; padding: 0 2px;">25</span>	27, 43, 93, 132	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	172	ASP	8.0
1	A	460	THR	7.5
1	A	198	TRP	6.3
1	A	171	ASP	6.0
1	A	71	ASP	5.3
1	A	173	LYS	4.8
1	A	168	LEU	4.7
1	A	169	ASP	4.4
1	A	175	ARG	4.3
1	A	176	LEU	4.3
1	A	174	THR	4.0
1	A	166	GLY	3.9
1	A	327	ASP	3.8
1	A	116	ASP	3.5
1	A	167	TYR	3.5
1	A	177	GLU	3.4
1	A	329	PHE	2.8
1	A	163	ASP	2.7
1	A	115	GLU	2.6
1	A	195	TYR	2.4
1	A	164	GLU	2.3
1	A	72	PRO	2.2
1	A	332	GLU	2.1
1	A	199	GLN	2.1
1	A	197	ASN	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	UDX	A	801	34/34	0.97	0.15	29,38,68,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.