



wwPDB EM Validation Summary Report ⓘ

Nov 15, 2022 – 11:45 AM JST

PDB ID : 6KWX
EMDB ID : EMD-0780
Title : cryo-EM structure of human PA200
Authors : Ouyang, S.; Hongxin, G.
Deposited on : 2019-09-09
Resolution : 3.75 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1878	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IHP	A	1902	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14227 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome activator complex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1751	Total	C	N	O	S	0	0
			14147	9105	2414	2549	79		

There are 37 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-34	MET	-	initiating methionine	UNP Q14997
A	-33	GLY	-	expression tag	UNP Q14997
A	-32	THR	-	expression tag	UNP Q14997
A	-31	THR	-	expression tag	UNP Q14997
A	-30	ARG	-	expression tag	UNP Q14997
A	-29	SER	-	expression tag	UNP Q14997
A	-28	THR	-	expression tag	UNP Q14997
A	-27	MET	-	expression tag	UNP Q14997
A	-26	SER	-	expression tag	UNP Q14997
A	-25	TYR	-	expression tag	UNP Q14997
A	-24	TYR	-	expression tag	UNP Q14997
A	-23	HIS	-	expression tag	UNP Q14997
A	-22	HIS	-	expression tag	UNP Q14997
A	-21	HIS	-	expression tag	UNP Q14997
A	-20	HIS	-	expression tag	UNP Q14997
A	-19	HIS	-	expression tag	UNP Q14997
A	-18	HIS	-	expression tag	UNP Q14997
A	-17	ASP	-	expression tag	UNP Q14997
A	-16	TYR	-	expression tag	UNP Q14997
A	-15	ASP	-	expression tag	UNP Q14997
A	-14	ILE	-	expression tag	UNP Q14997
A	-13	PRO	-	expression tag	UNP Q14997
A	-12	THR	-	expression tag	UNP Q14997
A	-11	THR	-	expression tag	UNP Q14997
A	-10	GLU	-	expression tag	UNP Q14997
A	-9	ASN	-	expression tag	UNP Q14997
A	-8	LEU	-	expression tag	UNP Q14997
A	-7	TYR	-	expression tag	UNP Q14997

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	PHE	-	expression tag	UNP Q14997
A	-5	GLN	-	expression tag	UNP Q14997
A	-4	GLY	-	expression tag	UNP Q14997
A	-3	ALA	-	expression tag	UNP Q14997
A	-2	MET	-	expression tag	UNP Q14997
A	-1	ASP	-	expression tag	UNP Q14997
A	0	PRO	-	expression tag	UNP Q14997
A	821	ILE	LEU	conflict	UNP Q14997
A	822	LEU	ILE	conflict	UNP Q14997

- Molecule 2 is [(1 {S},2 {R},3 {R},4 {S},5 {S},6 {R})-2-[oxidanyl(phosphonooxy)phosphoryl]oxy-3,4,5,6-tetraphosphonooxy-cyclohexyl] phosphono hydrogen phosphate (three-letter code: KOW) (formula: $C_6H_{20}O_{30}P_8$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	O	P	0
			44	6	30	8	

- Molecule 3 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula: $C_6H_{18}O_{24}P_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
3	A	1	Total	C	O	P	0
			36	6	24	6	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proteasome activator complex subunit 4



PRO	T1779	W1780	M1781	P1782	Q1783	L1784	L1785	M1786	M1787	L1788	S1789	A1790	H1791	L1792	M1793	D1794	P1795	Q1796	P1797	I1798	E1799	M1800	T1801	V1802	K1803	K1804	T1805	L1806	S1807	M1808	F1809	R1810	R1811	T1812	H1813	ASP	ASN	TRP	GLN	GLU	GLY	LYS	GLN	GLN	PHE	THR	ASP	GLN	LEU	LEU	VAL	LEU	THR	ASP	LEU	VAL	SER					
	N1716	T1719	M1720	P1782	S1682	P1721	S1683		R1658	T1660	V1661	L1662	L1665	Q1666	M1667	V1669	F1670	Y1671	M1672	L1673	F1674		L1677	M1678	N1679	E1680	D1681	A1682	D1685	I1686	R1687	W1688		T1691	S1692	L1693	L1694	E1695	D1696	E1697	Q1698		V1701	R1702	E1703	M1704	A1705	A1706	T1707	T1708	L1709		L1712	L1713	Q1714	C1715		D1776	V1777	P1778		
	V1555	MET	GLU	ASN	GLY	ILE	GLY	GLU	D1565	E1566	R1567		L1571	K1572	L1573	K1574	K1575		L1578	L1581		R1587	A1592	E1595		L1600	F1603	K1605	I1606	A1607	P1608	V1609	E1610	M1611	D1612	M1613	S1614	Y1615	D1616	E1617	L1618	K1619	R1620	D1621	A1622	K1623		L1628	Q1631		Q1638											
	D1461	A1462		L1465		L1468	Q1469		Q1475	E1476	W1477	R1478	V1479	P1480	A1481	L1482		R1485	L1486		L1490	K1493	L1494		V1497	Y1498	K1499	N1500	V1501	R1502		I1505	G1506	S1507	W1508	L1509	T1510	Y1511	I1512	F1513	M1514	I1515	D1516		I1537		K1540	L1541		M1545	D1546	V1547	D1548	E1549	E1550	I1551	Q1552	N1553	H1554			
	H1363		L1367		D1370	S1371	H1372		A1380	E1381	I1382	I1383	A1384	G1385		W1393	V1398	W1402	E1403	L1404	C1406		L1409		A1412		N1415		E1419		C1427		C1432	E1433	S1434	R1435		K1439	L1440		E1443	F1444	E1445	L1446	L1447	L1448	E1449	S1450	P1451	L1452	S1453	G1454	E1455	G1456	G1457							
	T1236	Q1237		N1246		L1249		E1262		V1269	K1271	T1272	H1273	W1274		N1282		V1285	Y1286		E1290		R1299	E1300	D1301		E1306	Q1307	T1308		K1317		Q1321	L1322	I1323		L1326	S1327	L1328	E1329	D1330	R1331	K1332	G1333	K1334	D1335		R1341		G1347		R1350	N1351	F1352	D1353		L1357		L1360			
	E1123	K1124		R1131		K1135		D1138		R1141	N1142		L1146	V1147	D1148		D1152	G1153	Q1154	E1155	Q1156	R1157		K1162	F1163	E1164		I1168	Q1169	L1170		R1176	D1177	D1178		L1181	P1182	L1183		I1186	R1187		E1191	N1192	L1193	N1194	H1195	D1196		K1213		K1216		I1224		C1227		K1234	P1235			
	L1012		D1015	R1016		V1019	T1020		G1033	N1034	H1035	S1036		A1041	N1042	L1043	H1044		Q1050	Q1051	T1052	W1053	P1054		A1055	I1056	V1057		L1061	S1062	Q1063	A1064	M1065	S1066	L1067		V1073	R1074		D1077		D1078	A1079	E1081	K1082		R1085	Q1086		I1086		Q1109	S1110	K1111	N1112	P1113		N1116	Q1117		L1120	
F918			Y921		M925	E926	N927	R928	L929		K932	K933	Q934	H935	I936		L939	L940	D942	R943		E949	L950	R951	T952	L953		G957	C958	E959	Y960	K961	K962	I963		D966	H967		I968		R969		D970	L971	L972	L974		S980		R983	H984	K985		F990		D1003	I1004		L1009	E1010	F1011	

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	103000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	18.901	Depositor
Minimum map value	-11.801	Depositor
Average map value	-0.010	Depositor
Map value standard deviation	1.005	Depositor
Recommended contour level	4.0	Depositor
Map size (\AA)	208.0, 208.0, 208.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.04, 1.04, 1.04	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KOW, IHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/14476	0.79	39/19631 (0.2%)

There are no bond length outliers.

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1353	ASP	CB-CG-OD1	10.33	127.60	118.30
1	A	516	LEU	CA-CB-CG	8.77	135.48	115.30
1	A	1177	ASP	CB-CG-OD1	8.71	126.14	118.30
1	A	1468	LEU	CA-CB-CG	8.20	134.15	115.30
1	A	302	LEU	CA-CB-CG	8.04	133.78	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	14147	0	14364	393	0
2	A	44	0	0	14	0
3	A	36	0	6	11	0
All	All	14227	0	14370	393	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 393 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:972:LEU:HD11	1:A:1004:ILE:CD1	1.30	1.60
1:A:972:LEU:CD1	1:A:1004:ILE:HD11	1.34	1.53
1:A:300:ARG:NH1	1:A:1631:GLN:HG2	1.26	1.49
1:A:298:VAL:CG2	1:A:299:PRO:HD3	1.57	1.33
1:A:324:LYS:O	1:A:326:VAL:N	1.63	1.31

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1743/1878 (93%)	1580 (91%)	157 (9%)	6 (0%)	41 74

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	325	LEU
1	A	477	GLY
1	A	962	LYS
1	A	1113	PRO
1	A	1480	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1593/1705 (93%)	1555 (98%)	38 (2%)	49 71

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1468	LEU
1	A	1712	LEU
1	A	1494	LEU
1	A	1545	MET
1	A	1787	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1474	GLN
1	A	1787	ASN
1	A	1613	ASN
1	A	1714	GLN
1	A	789	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	IHP	A	1902	-	36,36,36	0.72	0	54,60,60	0.94	0
2	K0W	A	1901	-	40,44,44	1.27	4 (10%)	66,74,74	1.25	8 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IHP	A	1902	-	-	2/30/54/54	0/1/1/1
2	K0W	A	1901	-	-	8/42/66/66	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1901	K0W	PA3-O13	3.41	1.65	1.59
2	A	1901	K0W	PA6-O16	2.92	1.64	1.59
2	A	1901	K0W	PA4-O14	2.77	1.64	1.59
2	A	1901	K0W	PA2-O12	2.75	1.64	1.59

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1901	K0W	O77-PA5-O15	3.39	109.32	102.48
2	A	1901	K0W	C5-C6-C1	3.37	117.78	110.41
2	A	1901	K0W	C5-C4-C3	2.79	116.52	110.41
2	A	1901	K0W	PA1-O76-PB1	-2.69	123.58	132.83
2	A	1901	K0W	C4-C3-C2	2.61	116.12	110.41

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1901	K0W	C4-C3-O13-PA3
2	A	1901	K0W	C2-O12-PA2-O22

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	1902	IHP	C3-O13-P3-O43
2	A	1901	K0W	C6-O11-PA1-O76
2	A	1901	K0W	C3-O13-PA3-O23


There are no ring outliers.

2 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1902	IHP	11	0
2	A	1901	K0W	14	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-0780. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 100



Y Index: 100




Z Index: 100


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 118



Y Index: 97



Z Index: 102


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 4.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 71 nm³; this corresponds to an approximate mass of 64 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.267 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.267 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.75	-	-
Author-provided FSC curve	3.75	4.25	3.83
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-0780 and PDB model 6KWX. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 4.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (4.0).

9.4 Atom inclusion ⓘ



At the recommended contour level, 76% of all backbone atoms, 58% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (4.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.5840	<div></div> 0.3570
A	<div></div> 0.5840	<div></div> 0.3570

