



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2020 – 03:19 AM BST

PDB ID : 6KXL  
Title : Crystal structure of the catalytic domain of Chitinophilus shinanonensis chitinase ChiL (CsChiL) complexed with N,N'-diacetylchitobiose  
Authors : Ueda, M.; Shimosaka, M.; Arai, R.  
Deposited on : 2019-09-12  
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.35 Å.





Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1551 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	372	 96% ..
1	B	372	 94% ..
2	C	2	 100%
2	D	2	 100%

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 7354 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Family 18 chitinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	368	Total	C	N	O	S	0	9	0
			2970	1892	503	569	6			
1	B	367	Total	C	N	O	S	0	10	0
			2963	1894	498	565	6			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	GLY	-	expression tag	UNP F8WSX2
A	36	GLY	-	expression tag	UNP F8WSX2
A	37	SER	-	expression tag	UNP F8WSX2
A	38	GLY	-	expression tag	UNP F8WSX2
A	39	GLY	-	expression tag	UNP F8WSX2
A	40	SER	-	expression tag	UNP F8WSX2
B	35	GLY	-	expression tag	UNP F8WSX2
B	36	GLY	-	expression tag	UNP F8WSX2
B	37	SER	-	expression tag	UNP F8WSX2
B	38	GLY	-	expression tag	UNP F8WSX2
B	39	GLY	-	expression tag	UNP F8WSX2
B	40	SER	-	expression tag	UNP F8WSX2

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			29	16	2	11			

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	0	0	0
			29	16	2	11			

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 2-METHOXYETHANOL (three-letter code: MXE) (formula:  $C_3H_8O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			5	3	2		

- Molecule 5 is MALONATE ION (three-letter code: MLI) (formula:  $C_3H_2O_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	3	4		
5	A	1	Total	C	O	0	0
			7	3	4		
5	A	1	Total	C	O	0	0
			7	3	4		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	3	4		
5	B	1	Total	C	O	0	0
			7	3	4		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			15	8	1	6		
8	B	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 9 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula:  $C_8H_{18}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			13	8	5		
9	B	1	Total	C	O	0	0
			13	8	5		

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	641	Total	O	0	1
			642	642		
10	B	569	Total	O	0	1
			570	570		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: Family 18 chitinase

Chain A:  96%




- Molecule 1: Family 18 chitinase

Chain B:  94%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%



## 4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.98Å 82.50Å 131.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.21 – 1.35	Depositor
% Data completeness (in resolution range)	95.2 (31.21-1.35)	Depositor
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.70 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.8.0253	Depositor
R, $R_{free}$	0.118 , 0.148	Depositor
Wilson B-factor (Å <sup>2</sup> )	12.0	Xtriage
Anisotropy	0.553	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7354	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, NAG, MLI, EDO, PG4, ACT, MXE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/3062	0.69	0/4143
1	B	0.60	0/3067	0.69	0/4149
All	All	0.60	0/6129	0.69	0/8292

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2970	0	2843	9	0
1	B	2963	0	2861	13	0
2	C	29	0	27	2	0
2	D	29	0	27	2	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
4	A	5	0	8	0	0
5	A	21	0	6	1	0
5	B	14	0	4	0	0
6	A	7	0	9	0	0
6	B	7	0	9	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	12	0	18	1	0
7	B	8	0	12	2	0
8	A	15	0	15	5	0
8	B	15	0	15	4	0
9	B	39	0	54	1	0
10	A	642	0	0	4	0
10	B	570	0	0	4	0
All	All	7354	0	5914	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:B:609:HOH:O	2:D:2:NAG:O7	2.04	0.73
1:B:386:TRP:CZ2	8:B:510:NAG:H3	2.23	0.72
1:B:406:THR:HG22	10:B:941:HOH:O	1.91	0.70
8:A:510:NAG:H4	2:C:2:NAG:H4	1.74	0.69
10:A:612:HOH:O	2:C:2:NAG:O7	2.12	0.67
5:A:505:MLI:H12	10:A:997:HOH:O	1.95	0.66
1:B:386:TRP:CE2	8:B:510:NAG:H3	2.33	0.63
1:A:386:TRP:CE2	8:A:510:NAG:H1	2.35	0.62
1:A:386:TRP:CZ2	8:A:510:NAG:H1	2.35	0.61
1:B:159:GLU:OE2	8:B:510:NAG:O7	2.22	0.58
1:A:234:TYR:HE1	8:A:510:NAG:H62	1.69	0.57
9:B:505:PG4:H41	10:B:1007:HOH:O	2.11	0.50
7:A:509:EDO:H21	10:A:1013:HOH:O	2.12	0.49
1:B:159:GLU:HA	1:B:160:TYR:CD1	2.51	0.45
1:A:159:GLU:HA	1:A:160:TYR:CD1	2.51	0.45
1:B:104:LYS:HG3	1:B:112:ASN:ND2	2.31	0.45
1:B:127[A]:ASP:OD1	1:B:172:ARG:NH2	2.50	0.44
1:A:256:TYR:CE2	1:A:261:ILE:HG21	2.53	0.44
1:B:347:LYS:HZ1	7:B:508:EDO:C1	2.30	0.44
1:B:378:LYS:HB3	1:B:380:LEU:HG	2.00	0.44
1:A:316:GLY:HA3	1:A:319:GLU:O	2.19	0.43
8:B:510:NAG:H2	2:D:2:NAG:O4	2.19	0.42
1:A:322[A]:ASN:OD1	10:A:601:HOH:O	2.22	0.41
1:A:180:THR:HA	1:A:222:VAL:HG12	2.03	0.41
1:A:234:TYR:CE1	8:A:510:NAG:H62	2.53	0.41
1:B:105:LYS:NZ	10:B:621:HOH:O	2.54	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:115[B]:SER:OG	1:B:157:ASP:OD2	2.38	0.41
1:B:376:LYS:CE	1:B:403:GLU:O	2.69	0.40
1:B:347:LYS:CE	7:B:508:EDO:H11	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	2	15,15,15	0.63	0	21,21,21	1.17	2 (9%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	2	2	14,14,15	0.52	0	17,19,21	0.79	0
2	NAG	D	1	2	15,15,15	0.46	0	21,21,21	1.14	2 (9%)
2	NAG	D	2	2	14,14,15	0.51	0	17,19,21	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2	-	0/6/26/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	2	-	0/6/26/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	NAG	C1-O5-C5	-2.32	109.29	113.66
2	D	1	NAG	C3-C4-C5	-2.25	106.23	110.24
2	D	1	NAG	C1-O5-C5	-2.16	109.58	113.66
2	C	1	NAG	C3-C2-N2	-2.13	106.60	110.62

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	NAG	2	0
2	D	2	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	MXE	A	502	-	4,4,4	0.33	0	3,3,3	0.29	0
3	ACT	A	501	-	1,3,3	3.04	1 (100%)	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MLI	A	504	-	0,6,6	0.00	-	0,7,7	0.00	-
3	ACT	B	501	-	1,3,3	2.61	1 (100%)	0,3,3	0.00	-
5	MLI	B	503	-	0,6,6	0.00	-	0,7,7	0.00	-
5	MLI	A	505	-	0,6,6	0.00	-	0,7,7	0.00	-
7	EDO	A	507	-	3,3,3	0.12	0	2,2,2	0.28	0
7	EDO	A	508	-	3,3,3	0.18	0	2,2,2	0.30	0
6	PGE	B	507	-	6,6,9	0.16	0	5,5,8	0.18	0
9	PG4	B	504	-	12,12,12	0.22	0	11,11,11	0.24	0
7	EDO	B	508	-	3,3,3	0.14	0	2,2,2	0.41	0
8	NAG	B	510	-	15,15,15	0.74	0	21,21,21	1.29	1 (4%)
8	NAG	A	510	-	15,15,15	0.78	0	21,21,21	2.21	9 (42%)
9	PG4	B	505	-	12,12,12	0.21	0	11,11,11	0.19	0
5	MLI	B	502	-	0,6,6	0.00	-	0,7,7	0.00	-
5	MLI	A	503	-	0,6,6	0.00	-	0,7,7	0.00	-
9	PG4	B	506	-	12,12,12	0.19	0	11,11,11	0.08	0
7	EDO	B	509	-	3,3,3	0.14	0	2,2,2	0.47	0
6	PGE	A	506	-	6,6,9	0.11	0	5,5,8	0.19	0
7	EDO	A	509	-	3,3,3	0.13	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MXE	A	502	-	-	1/2/2/2	-
5	MLI	A	505	-	-	0/0/4/4	-
5	MLI	A	504	-	-	0/0/4/4	-
5	MLI	B	503	-	-	0/0/4/4	-
7	EDO	A	507	-	-	0/1/1/1	-
6	PGE	B	507	-	-	1/4/4/7	-
7	EDO	A	508	-	-	1/1/1/1	-
7	EDO	B	509	-	-	1/1/1/1	-
9	PG4	B	504	-	-	4/10/10/10	-
7	EDO	B	508	-	-	0/1/1/1	-
8	NAG	B	510	-	-	0/6/26/26	0/1/1/1
8	NAG	A	510	-	-	4/6/26/26	0/1/1/1
5	MLI	B	502	-	-	0/0/4/4	-
5	MLI	A	503	-	-	0/0/4/4	-
9	PG4	B	506	-	-	7/10/10/10	-
9	PG4	B	505	-	-	5/10/10/10	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	A	506	-	-	0/4/4/7	-
7	EDO	A	509	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	ACT	CH3-C	3.04	1.52	1.48
3	B	501	ACT	CH3-C	2.61	1.52	1.48

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	510	NAG	C1-C2-N2	-4.58	105.42	110.73
8	A	510	NAG	O3-C3-C4	-4.40	100.17	110.35
8	A	510	NAG	C1-C2-C3	3.93	115.90	110.54
8	A	510	NAG	O3-C3-C2	3.83	117.39	109.66
8	A	510	NAG	C3-C4-C5	3.50	116.48	110.24
8	A	510	NAG	O4-C4-C3	-3.28	102.78	110.35
8	A	510	NAG	C4-C3-C2	2.67	114.26	110.34
8	A	510	NAG	C1-O5-C5	-2.28	109.36	113.66
8	A	510	NAG	O5-C1-C2	2.26	111.79	109.52
8	A	510	NAG	O4-C4-C5	2.01	114.29	109.30

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	B	504	PG4	O3-C5-C6-O4
8	A	510	NAG	C4-C5-C6-O6
6	B	507	PGE	O2-C3-C4-O3
9	B	505	PG4	O1-C1-C2-O2
9	B	505	PG4	C6-C5-O3-C4
8	A	510	NAG	O5-C5-C6-O6
8	A	510	NAG	C8-C7-N2-C2
9	B	506	PG4	O2-C3-C4-O3
7	A	508	EDO	O1-C1-C2-O2
9	B	506	PG4	O4-C7-C8-O5
9	B	504	PG4	O4-C7-C8-O5
4	A	502	MXE	C1-C2-O2-C3
9	B	506	PG4	C1-C2-O2-C3
9	B	505	PG4	C8-C7-O4-C6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
8	A	510	NAG	O7-C7-N2-C2
9	B	504	PG4	C1-C2-O2-C3
9	B	504	PG4	C6-C5-O3-C4
7	B	509	EDO	O1-C1-C2-O2
9	B	506	PG4	O1-C1-C2-O2
9	B	506	PG4	O3-C5-C6-O4
7	A	509	EDO	O1-C1-C2-O2
9	B	506	PG4	C3-C4-O3-C5
9	B	505	PG4	O4-C7-C8-O5
9	B	506	PG4	C5-C6-O4-C7
9	B	505	PG4	O2-C3-C4-O3

There are no ring outliers.

6 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	505	MLI	1	0
7	B	508	EDO	2	0
8	B	510	NAG	4	0
8	A	510	NAG	5	0
9	B	505	PG4	1	0
7	A	509	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.


### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


EDS failed to run properly - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands ⓘ

EDS failed to run properly - this section is therefore empty.

## 6.5 Other polymers ⓘ

EDS failed to run properly - this section is therefore empty.