



wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 10:33 am BST

PDB ID : 3KZ4
Title : Crystal Structure of the Rotavirus Double Layered Particle
Authors : McClain, B.; Settembre, E.C.; Bellamy, A.R.; Harrison, S.C.
Deposited on : 2009-12-07
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1288 (4.00-3.60)
Ramachandran outliers	138981	1243 (4.00-3.60)
Sidechain outliers	138945	1237 (4.00-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	880	
1	B	880	
2	C	397	
2	D	397	
2	E	397	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 54109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inner capsid protein VP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	781	Total	C	N	O	S	0	0	0
			6374	4049	1099	1190	36			
1	B	810	Total	C	N	O	S	0	0	0
			6624	4211	1138	1239	36			

- Molecule 2 is a protein called Intermediate capsid protein VP6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	D	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	E	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	F	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	G	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	H	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	I	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	J	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	K	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	L	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	M	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			
2	N	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	397	Total	C	N	O	S	0	0	0
			3162	2001	550	596	15			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	O	1	Total	Zn	0	0
			1	1		
3	I	1	Total	Zn	0	0
			1	1		
3	L	1	Total	Zn	0	0
			1	1		
3	C	1	Total	Zn	0	0
			1	1		
3	F	1	Total	Zn	0	0
			1	1		

SEQUENCE-PLOTS INFOmissingINFO

3 Data and refinement statistics

EDS was not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	740.75Å 1198.07Å 1345.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.80	Depositor
% Data completeness (in resolution range)	15.9 (30.00-3.80)	Depositor
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.49 (at 3.78Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.328 , 0.339	Depositor
Wilson B-factor (Å ²)	167.1	Xtriage
Anisotropy	0.043	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	54109	wwPDB-VP
Average B, all atoms (Å ²)	154.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality

4.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/6491	0.84	9/8806 (0.1%)
1	B	0.50	0/6745	0.82	10/9149 (0.1%)
2	C	0.51	0/3232	0.78	5/4397 (0.1%)
2	D	0.51	0/3232	0.78	5/4397 (0.1%)
2	E	0.51	0/3232	0.77	5/4397 (0.1%)
2	F	0.50	0/3232	0.76	5/4397 (0.1%)
2	G	0.50	0/3232	0.77	5/4397 (0.1%)
2	H	0.50	0/3232	0.76	5/4397 (0.1%)
2	I	0.50	0/3232	0.77	5/4397 (0.1%)
2	J	0.51	0/3232	0.78	5/4397 (0.1%)
2	K	0.51	0/3232	0.77	5/4397 (0.1%)
2	L	0.51	0/3232	0.77	5/4397 (0.1%)
2	M	0.51	0/3232	0.77	5/4397 (0.1%)
2	N	0.51	0/3232	0.77	5/4397 (0.1%)
2	O	0.52	0/3232	0.79	7/4397 (0.2%)
All	All	0.50	0/55252	0.79	86/75116 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 86 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	273	TYR	CB-CG-CD2	10.70	127.42	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	273	TYR	CB-CG-CD1	-9.86	115.08	121.00
1	A	273	TYR	CA-CB-CG	9.85	132.12	113.40
1	B	273	TYR	CB-CG-CD1	9.85	126.91	121.00
1	B	273	TYR	CB-CG-CD2	-9.21	115.48	121.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	273	TYR	Sidechain

4.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6374	0	6394	992	0
1	B	6624	0	6652	1122	0
2	C	3162	0	3101	170	0
2	D	3162	0	3101	179	0
2	E	3162	0	3101	164	0
2	F	3162	0	3101	160	0
2	G	3162	0	3101	149	0
2	H	3162	0	3101	175	0
2	I	3162	0	3101	215	0
2	J	3162	0	3101	187	0
2	K	3162	0	3101	166	0
2	L	3162	0	3101	178	0
2	M	3162	0	3101	173	0
2	N	3162	0	3101	191	0
2	O	3162	0	3101	163	0
3	C	1	0	0	0	0
3	F	1	0	0	0	0
3	I	1	0	0	0	0
3	L	1	0	0	0	0
3	O	1	0	0	0	0
All	All	54109	0	53359	4148	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 4148 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:771:VAL:HB	1:B:809:PHE:HB3	1.23	1.18
1:A:333:VAL:HG11	1:A:380:LYS:HA	1.27	1.15
1:A:428:GLN:OE1	1:A:456:PHE:HB2	1.46	1.13
1:A:563:MET:HA	1:A:563:MET:HE3	1.22	1.12
2:G:145:ARG:HB3	2:G:145:ARG:HH11	1.10	1.12

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	779/880 (88%)	431 (55%)	199 (26%)	149 (19%)	0	2
1	B	808/880 (92%)	458 (57%)	204 (25%)	146 (18%)	0	2
2	C	395/397 (100%)	318 (80%)	54 (14%)	23 (6%)	1	21
2	D	395/397 (100%)	315 (80%)	50 (13%)	30 (8%)	1	15
2	E	395/397 (100%)	323 (82%)	49 (12%)	23 (6%)	1	21
2	F	395/397 (100%)	324 (82%)	47 (12%)	24 (6%)	1	20
2	G	395/397 (100%)	323 (82%)	49 (12%)	23 (6%)	1	21
2	H	395/397 (100%)	323 (82%)	50 (13%)	22 (6%)	2	21
2	I	395/397 (100%)	320 (81%)	51 (13%)	24 (6%)	1	20
2	J	395/397 (100%)	320 (81%)	53 (13%)	22 (6%)	2	21
2	K	395/397 (100%)	318 (80%)	52 (13%)	25 (6%)	1	20
2	L	395/397 (100%)	322 (82%)	47 (12%)	26 (7%)	1	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	M	395/397 (100%)	322 (82%)	52 (13%)	21 (5%)	2	23
2	N	395/397 (100%)	317 (80%)	50 (13%)	28 (7%)	1	17
2	O	395/397 (100%)	317 (80%)	47 (12%)	31 (8%)	1	15
All	All	6722/6921 (97%)	5051 (75%)	1054 (16%)	617 (9%)	1	12

5 of 617 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	LYS
1	A	130	GLN
1	A	193	SER
1	A	198	LYS
1	A	220	ALA

4.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	715/809 (88%)	603 (84%)	112 (16%)	2	17
1	B	744/809 (92%)	635 (85%)	109 (15%)	3	19
2	C	350/350 (100%)	325 (93%)	25 (7%)	14	45
2	D	350/350 (100%)	322 (92%)	28 (8%)	12	42
2	E	350/350 (100%)	325 (93%)	25 (7%)	14	45
2	F	350/350 (100%)	327 (93%)	23 (7%)	16	48
2	G	350/350 (100%)	327 (93%)	23 (7%)	16	48
2	H	350/350 (100%)	329 (94%)	21 (6%)	19	50
2	I	350/350 (100%)	327 (93%)	23 (7%)	16	48
2	J	350/350 (100%)	327 (93%)	23 (7%)	16	48
2	K	350/350 (100%)	327 (93%)	23 (7%)	16	48
2	L	350/350 (100%)	321 (92%)	29 (8%)	11	40
2	M	350/350 (100%)	325 (93%)	25 (7%)	14	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	N	350/350 (100%)	324 (93%)	26 (7%)	13	44
2	O	350/350 (100%)	329 (94%)	21 (6%)	19	50
All	All	6009/6168 (97%)	5473 (91%)	536 (9%)	9	38

5 of 536 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	374	TYR
2	F	13	ASP
2	N	73	LEU
2	D	19	VAL
2	D	382	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 220 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	345	ASN
2	H	26	ASN
2	N	167	ASN
2	F	72	ASN
2	F	345	ASN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

4.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data ⓘ

5.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

5.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

5.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

5.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.