



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 08:41 pm BST

PDB ID : 5L0F
Title : Human metavinculin quadruple mutant (residues 959-1134)
Authors : Chinthalapudi, K.; Izard, T.
Deposited on : 2016-07-27
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>17%</div> </div> </div>
1	B	176	<div> <div>6%</div> <div> <div></div> <div>74%</div> <div>26%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3038 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vinculin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	176	Total	C	N	O	S	0	1	0
			1373	845	255	264	9			
1	B	176	Total	C	N	O	S	0	1	0
			1366	843	251	263	9			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	975	GLN	ARG	engineered mutation	UNP P18206
A	979	GLN	LYS	engineered mutation	UNP P18206
A	1107	GLN	ARG	engineered mutation	UNP P18206
A	1128	GLN	ARG	engineered mutation	UNP P18206
B	975	GLN	ARG	engineered mutation	UNP P18206
B	979	GLN	LYS	engineered mutation	UNP P18206
B	1107	GLN	ARG	engineered mutation	UNP P18206
B	1128	GLN	ARG	engineered mutation	UNP P18206

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	137	Total	O	0	0
			137	137		
3	B	150	Total	O	0	0
			150	150		

- Molecule 1: Vinculin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.15Å 70.91Å 118.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.79 – 2.76 60.79 – 2.76	Depositor EDS
% Data completeness (in resolution range)	100.0 (60.79-2.76) 100.0 (60.79-2.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.78 (at 2.77Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.201 , 0.247 0.208 , 0.259	Depositor DCC
R_{free} test set	526 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 77.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3038	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 65.57 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.9328e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1386	0.60	0/1867
1	B	0.47	0/1382	0.61	0/1863
All	All	0.47	0/2768	0.60	0/3730

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1373	0	1420	29	1
1	B	1366	0	1417	63	1
2	A	6	0	8	0	0
2	B	6	0	8	1	0
3	A	137	0	0	1	0
3	B	150	0	0	2	0
All	All	3038	0	2853	87	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1114:ILE:HG23	3:B:1343:HOH:O	1.57	1.03
1:B:1039:GLN:HE21	1:B:1120:ALA:HB2	1.21	1.00
1:B:987:ILE:HD12	1:B:1124:LEU:HD13	1.44	0.98
1:B:1039:GLN:NE2	1:B:1120:ALA:CB	2.33	0.91
1:B:1039:GLN:NE2	1:B:1120:ALA:HB2	1.84	0.91
1:B:987:ILE:HD12	1:B:1124:LEU:CD1	2.01	0.90
1:A:987:ILE:HD12	1:A:1124:LEU:CD2	2.02	0.89
1:A:987:ILE:HD12	1:A:1124:LEU:HD21	1.56	0.88
1:B:1039:GLN:HE21	1:B:1120:ALA:CB	1.87	0.88
1:B:1056:ILE:HD11	1:B:1101:SER:HB3	1.62	0.81
1:B:1056:ILE:CD1	1:B:1101:SER:HB3	2.10	0.80
1:B:1036:VAL:CG2	1:B:1124:LEU:CD1	2.62	0.78
1:B:1117:ARG:NH1	1:B:1119:ASP:HB3	2.00	0.77
1:B:1036:VAL:HG21	1:B:1124:LEU:HD12	1.66	0.77
1:B:1036:VAL:HG21	1:B:1124:LEU:CD1	2.13	0.76
1:A:979:GLN:HG2	1:A:980:TRP:CE3	2.20	0.75
1:B:1039:GLN:NE2	1:B:1120:ALA:HB1	2.05	0.72
1:B:1036:VAL:CG2	1:B:1124:LEU:HD12	2.20	0.71
1:A:987:ILE:CD1	1:A:1124:LEU:HD21	2.21	0.69
1:B:1113:SER:HA	1:B:1116:ILE:HD12	1.73	0.69
1:B:1035:GLU:OE2	1:B:1038:LYS:NZ	2.26	0.69
1:B:1035:GLU:HB3	1:B:1122:PHE:CE2	2.28	0.68
1:B:1113:SER:HA	1:B:1116:ILE:CD1	2.24	0.67
1:B:1036:VAL:CG2	1:B:1124:LEU:HD11	2.23	0.66
1:A:1073:MET:HG2	1:A:1079:ILE:HD13	1.79	0.65
1:B:1073:MET:HG2	1:B:1079:ILE:HD13	1.78	0.65
1:A:1042:ASP:HB2	1:A:1115:LYS:HG2	1.79	0.65
1:B:1035:GLU:OE2	1:B:1038:LYS:CE	2.46	0.64
1:B:987:ILE:CD1	1:B:1124:LEU:CD1	2.77	0.62
1:B:1054:GLU:O	1:B:1057:PRO:HD2	2.01	0.61
1:B:1042:ASP:HB2	1:B:1115:LYS:HD2	1.84	0.60
1:B:1053:CYS:O	1:B:1057:PRO:HD3	2.03	0.58
1:A:974:HIS:CE1	1:A:978:THR:HG21	2.38	0.58
1:B:1052:VAL:O	1:B:1056:ILE:HG12	2.05	0.56
1:B:1034:LYS:O	1:B:1038:LYS:HG3	2.05	0.56
1:B:974:HIS:CE1	1:B:978:THR:HG21	2.39	0.56
1:B:987:ILE:CD1	1:B:1124:LEU:HD11	2.35	0.56
1:A:979:GLN:CG	1:A:980:TRP:CE3	2.88	0.55
1:B:963:ASN:HD21	1:B:1092:VAL:CG2	2.19	0.55
1:B:1117:ARG:HH12	1:B:1119:ASP:HB3	1.69	0.55
1:A:1020:LYS:HE3	1:B:1065:ILE:CD1	2.38	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:979:GLN:CD	1:A:980:TRP:CZ3	2.82	0.54
1:B:1129:LYS:HD2	2:B:1201:GOL:H2	1.90	0.53
1:B:1049:LEU:HD11	1:B:1108:GLU:HG3	1.89	0.53
1:B:1035:GLU:OE2	1:B:1038:LYS:HE3	2.09	0.53
1:B:1049:LEU:CD1	1:B:1108:GLU:HG3	2.39	0.53
1:B:1036:VAL:HG21	1:B:1124:LEU:HD11	1.89	0.52
1:A:1001:MET:O	1:A:1005:VAL:HG13	2.10	0.52
1:A:979:GLN:OE1	1:A:980:TRP:CZ3	2.63	0.52
1:B:1033:ALA:HB3	1:B:1053:CYS:SG	2.48	0.52
1:A:963:ASN:HD22	1:A:966:ILE:H	1.56	0.52
1:A:1063:LEU:HB2	1:A:1094:ASN:HD21	1.74	0.51
1:A:1020:LYS:CE	1:B:1065:ILE:CD1	2.89	0.50
1:B:1053:CYS:O	1:B:1057:PRO:CD	2.60	0.50
1:B:1065:ILE:O	1:B:1069:VAL:HG23	2.12	0.49
1:B:1030:THR:HG23	1:B:1053:CYS:HB3	1.95	0.47
1:B:1049:LEU:HD12	1:B:1108:GLU:HG2	1.96	0.47
1:A:1119:ASP:HB3	1:A:1122:PHE:HB3	1.96	0.47
1:B:1045:ILE:O	1:B:1049:LEU:HD13	2.13	0.47
1:B:1096:GLN:NE2	3:B:1301:HOH:O	2.27	0.47
1:A:1119:ASP:OD2	1:A:1122:PHE:CD2	2.68	0.47
1:A:979:GLN:HE21	1:A:1128:GLN:HE22	1.63	0.47
1:B:1049:LEU:CD1	1:B:1108:GLU:CG	2.93	0.47
1:B:1056:ILE:CD1	1:B:1101:SER:CB	2.89	0.46
1:B:963:ASN:ND2	1:B:1092:VAL:HG22	2.30	0.45
1:A:987:ILE:HD12	1:A:1124:LEU:HD23	1.95	0.45
1:B:1049:LEU:HD12	1:B:1108:GLU:CG	2.47	0.45
1:B:979:GLN:HE21	1:B:1128:GLN:HE22	1.64	0.45
1:B:1112:ALA:O	1:B:1116:ILE:HG13	2.17	0.44
1:A:961:PRO:HG2	3:A:1330:HOH:O	2.17	0.44
1:B:1086:GLN:O	1:B:1090:MET:HG2	2.19	0.43
1:A:1119:ASP:OD2	1:A:1122:PHE:HB3	2.19	0.43
1:A:1086:GLN:O	1:A:1090:MET:HG2	2.20	0.42
1:A:1033:ALA:HB1	1:A:1049:LEU:HD22	2.01	0.42
1:A:1020:LYS:HE2	1:B:1065:ILE:HD12	2.01	0.42
1:B:1056:ILE:HB	1:B:1057:PRO:HD3	2.02	0.41
1:B:1068:THR:O	1:B:1072:THR:HG23	2.20	0.41
1:B:963:ASN:ND2	1:B:1092:VAL:CG2	2.83	0.41
1:A:1065:ILE:O	1:A:1069:VAL:HG23	2.20	0.41
1:B:987:ILE:HD11	1:B:1124:LEU:HD11	2.02	0.41
1:A:963:ASN:ND2	1:A:966:ILE:HG13	2.35	0.41
1:B:1117:ARG:HG3	1:B:1117:ARG:HH11	1.86	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1065:ILE:CD1	1:B:1020:LYS:HE2	2.50	0.40
1:B:1041:THR:OG1	1:B:1115:LYS:HB3	2.21	0.40
1:B:1036:VAL:HG22	1:B:1124:LEU:CD1	2.51	0.40
1:A:1020:LYS:HE2	1:B:1065:ILE:CD1	2.51	0.40
1:A:1068:THR:O	1:A:1072:THR:HG23	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1125:ARG:NH1	1:B:1119:ASP:OD1[3_756]	1.84	0.36

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	175/176 (99%)	174 (99%)	1 (1%)	0	100	100
1	B	175/176 (99%)	174 (99%)	1 (1%)	0	100	100
All	All	350/352 (99%)	348 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	149/148 (101%)	149 (100%)	0	100	100
1	B	149/148 (101%)	149 (100%)	0	100	100
All	All	298/296 (101%)	298 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	960	GLN
1	A	963	ASN
1	A	979	GLN
1	A	1094	ASN
1	B	963	ASN
1	B	974	HIS
1	B	979	GLN
1	B	1039	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	1201	-	5,5,5	0.06	0	5,5,5	0.18	0
2	GOL	A	1201	-	5,5,5	0.05	0	5,5,5	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	1201	-	-	0/4/4/4	-
2	GOL	A	1201	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1201	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	176/176 (100%)	-0.07	4 (2%) 60 69	9, 21, 51, 66	0
1	B	176/176 (100%)	0.06	10 (5%) 23 28	10, 22, 53, 74	0
All	All	352/352 (100%)	-0.00	14 (3%) 38 45	9, 21, 52, 74	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1122	PHE	8.1
1	B	1119	ASP	3.9
1	B	1115	LYS	3.7
1	B	1114	ILE	3.3
1	B	1121	GLY	2.9
1	B	1134	GLN	2.6
1	A	1077	THR	2.5
1	B	1054	GLU	2.5
1	B	1122	PHE	2.5
1	B	1077	THR	2.4
1	A	1121	GLY	2.4
1	B	1116	ILE	2.3
1	A	1078	ASN	2.1
1	B	1078	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	B	1201	6/6	0.72	0.20	49,53,54,54	0
2	GOL	A	1201	6/6	0.80	0.16	63,65,67,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.