



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2021 – 06:07 PM EDT

PDB ID : 7L0F  
Title : Monobody 12VC3 Bound to HRAS(WT)  
Authors : Teng, K.W.; Hattori, T.; Tsai, S.; Koide, S.  
Deposited on : 2020-12-11  
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.17.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.17.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>%</span> <span>93%</span> <span>6% .</span> </div> </div>
1	E	167	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>%</span> <span>95%</span> <span>. .</span> </div> </div>
1	G	167	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>4%</span> <span>91%</span> <span>8% ..</span> </div> </div>
1	L	167	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span></span> <span>93%</span> <span>6% .</span> </div> </div>
2	B	97	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>3%</span> <span>90%</span> <span>6% .</span> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	97	<div><div><div></div><div></div><div></div></div><div><div>2%</div><div>85%</div><div>11%</div><div></div></div><div></div></div>
2	H	97	<div><div><div></div><div></div><div></div></div><div><div>2%</div><div>91%</div><div>6%</div><div></div></div><div></div></div>
2	M	97	<div><div><div></div><div></div><div></div></div><div><div>%</div><div>87%</div><div>7%</div><div></div></div><div></div></div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8875 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTPase HRas.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	166	Total	C	N	O	S	0	0	0
			1322	824	228	263	7			
1	A	166	Total	C	N	O	S	0	0	0
			1322	824	228	263	7			
1	G	166	Total	C	N	O	S	0	0	0
			1322	824	228	263	7			
1	L	166	Total	C	N	O	S	0	0	0
			1322	824	228	263	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	0	SER	-	expression tag	UNP P01112
A	0	SER	-	expression tag	UNP P01112
G	0	SER	-	expression tag	UNP P01112
L	0	SER	-	expression tag	UNP P01112


- Molecule 2 is a protein called Monobody 12VC3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	93	Total	C	N	O	0	0	0
			702	459	113	130			
2	B	93	Total	C	N	O	0	0	0
			702	459	113	130			
2	H	94	Total	C	N	O	0	0	0
			708	462	114	132			
2	M	93	Total	C	N	O	0	0	0
			702	459	113	130			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0
3	G	1	Total Mg 1 1	0	0
3	L	1	Total Mg 1 1	0	0

- Molecule 4 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>S).



*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	F	86	Total 86	O 86	0	0
5	A	97	Total 97	O 97	0	0
5	B	50	Total 50	O 50	0	0
5	G	41	Total 41	O 41	0	0
5	H	57	Total 57	O 57	0	0
5	L	118	Total 118	O 118	0	0
5	M	74	Total 74	O 74	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GTPase HRas



- Molecule 1: GTPase HRas




- Molecule 1: GTPase HRas




- Molecule 1: GTPase HRas




- Molecule 2: Monobody 12VC3




- Molecule 2: Monobody 12VC3



• Molecule 2: Monobody 12VC3



• Molecule 2: Monobody 12VC3



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.01Å 64.82Å 127.12Å 90.00° 102.37° 90.00°	Depositor
Resolution (Å)	44.88 – 1.98 44.84 – 1.98	Depositor EDS
% Data completeness (in resolution range)	95.5 (44.88-1.98) 95.6 (44.84-1.98)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.65 (at 1.98Å)	Xtriage
Refinement program	REFMAC v5.8.0238	Depositor
R, $R_{free}$	0.159 , 0.191 0.170 , 0.200	Depositor DCC
$R_{free}$ test set	3851 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.8	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8875	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GSP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/1341	0.83	0/1810
1	E	0.74	0/1341	0.86	0/1810
1	G	0.66	0/1341	0.80	0/1810
1	L	0.69	0/1341	0.81	0/1810
2	B	0.63	0/723	0.77	0/990
2	F	0.65	0/723	0.80	0/990
2	H	0.64	0/729	0.79	0/998
2	M	0.68	0/723	0.83	0/990
All	All	0.68	0/8262	0.82	0/11208

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1322	0	1294	7	0
1	E	1322	0	1294	2	0
1	G	1322	0	1294	11	0
1	L	1322	0	1294	7	0
2	B	702	0	706	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	702	0	706	6	0
2	H	708	0	711	4	0
2	M	702	0	706	9	0
3	A	1	0	0	0	0
3	E	1	0	0	0	0
3	G	1	0	0	0	0
3	L	1	0	0	0	0
4	A	32	0	12	1	0
4	E	32	0	12	0	0
4	G	32	0	12	2	0
4	L	32	0	12	0	0
5	A	97	0	0	1	0
5	B	50	0	0	0	0
5	E	118	0	0	0	0
5	F	86	0	0	0	0
5	G	41	0	0	0	0
5	H	57	0	0	0	0
5	L	118	0	0	3	0
5	M	74	0	0	1	0
All	All	8875	0	8053	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:5:PRO:O	2:M:6:THR:HG23	1.72	0.89
1:L:27:HIS:ND1	5:L:302:HOH:O	2.23	0.70
1:L:68:ARG:HG2	5:L:305:HOH:O	1.97	0.63
1:G:52:LEU:C	1:G:52:LEU:HD13	2.19	0.63
1:L:102:ARG:NH2	5:L:303:HOH:O	2.32	0.61
2:B:66:GLY:HA2	2:B:95:THR:HG22	1.84	0.59
1:G:23:LEU:C	1:G:23:LEU:HD13	2.24	0.58
1:A:67:MET:SD	2:B:30:PHE:HZ	2.27	0.57
1:A:68:ARG:HG2	5:A:303:HOH:O	2.04	0.57
1:A:23:LEU:HD22	1:A:156:PHE:CG	2.40	0.56
1:E:23:LEU:HD22	1:E:156:PHE:CG	2.43	0.54
2:H:15:PRO:HA	2:H:95:THR:HG22	1.89	0.53
2:B:34:ILE:HD12	2:B:34:ILE:N	2.24	0.53
2:M:34:ILE:HD12	2:M:34:ILE:N	2.25	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:112:VAL:CG1	1:L:142:ILE:HD13	2.39	0.52
1:G:141:TYR:O	1:G:142:ILE:HD13	2.09	0.51
2:B:94:ARG:HG2	2:B:95:THR:HG23	1.92	0.51
1:L:112:VAL:HG11	1:L:142:ILE:HD13	1.92	0.51
2:F:15:PRO:HA	2:F:95:THR:HG22	1.93	0.50
1:L:23:LEU:HD22	1:L:156:PHE:CG	2.47	0.50
2:H:66:GLY:O	2:H:94:ARG:NH1	2.44	0.50
1:G:84:ILE:HD12	1:G:123:ARG:HB2	1.94	0.49
2:H:34:ILE:HD12	2:H:34:ILE:N	2.26	0.49
2:F:66:GLY:O	2:F:94:ARG:NH1	2.46	0.49
1:A:84:ILE:HD12	1:A:123:ARG:HB2	1.93	0.49
1:G:23:LEU:HD13	1:G:23:LEU:O	2.14	0.48
2:M:14:THR:HB	2:M:15:PRO:HD2	1.96	0.47
1:L:141:TYR:OH	1:L:143:GLU:HG3	2.14	0.47
2:H:34:ILE:O	2:H:48:ALA:HA	2.16	0.45
1:A:161:ARG:HD2	1:G:150:GLN:OE1	2.17	0.45
1:E:94:HIS:HD2	1:E:137:TYR:OH	2.00	0.45
2:M:5:PRO:HG3	2:M:32:TYR:HE1	1.82	0.44
1:G:3:GLU:HG3	1:G:52:LEU:HD12	1.99	0.44
2:F:5:PRO:HD2	2:F:89:ILE:HG22	1.99	0.44
2:F:34:ILE:O	2:F:48:ALA:HA	2.18	0.43
2:M:15:PRO:HA	2:M:95:THR:HG22	2.00	0.43
1:G:84:ILE:HD13	1:G:118:CYS:HA	2.02	0.42
2:B:34:ILE:O	2:B:48:ALA:HA	2.20	0.42
1:G:16:LYS:NZ	4:G:202:GSP:O2G	2.46	0.41
2:M:77:GLY:HA3	5:M:105:HOH:O	2.20	0.41
2:M:5:PRO:HG3	2:M:32:TYR:CE1	2.54	0.41
2:M:5:PRO:O	2:M:6:THR:CG2	2.57	0.41
1:A:67:MET:SD	2:B:30:PHE:CZ	3.11	0.41
1:A:32:TYR:CD1	4:A:202:GSP:H5'2	2.56	0.40
2:F:4:VAL:HG11	2:F:84:TYR:OH	2.21	0.40
1:G:101:LYS:HD2	1:G:107:ASP:HA	2.03	0.40
1:G:117:LYS:HG2	4:G:202:GSP:C6	2.56	0.40
2:M:34:ILE:O	2:M:48:ALA:HA	2.20	0.40
2:F:4:VAL:HG13	2:F:5:PRO:HA	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	164/167 (98%)	161 (98%)	3 (2%)	0	100	100
1	E	164/167 (98%)	160 (98%)	4 (2%)	0	100	100
1	G	164/167 (98%)	160 (98%)	4 (2%)	0	100	100
1	L	164/167 (98%)	161 (98%)	3 (2%)	0	100	100
2	B	91/97 (94%)	89 (98%)	2 (2%)	0	100	100
2	F	91/97 (94%)	90 (99%)	1 (1%)	0	100	100
2	H	92/97 (95%)	90 (98%)	2 (2%)	0	100	100
2	M	91/97 (94%)	88 (97%)	1 (1%)	2 (2%)	6	1
All	All	1021/1056 (97%)	999 (98%)	20 (2%)	2 (0%)	47	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	M	5	PRO
2	M	6	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/145 (99%)	142 (99%)	2 (1%)	67	62
1	E	144/145 (99%)	141 (98%)	3 (2%)	53	47
1	G	144/145 (99%)	142 (99%)	2 (1%)	67	62

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	144/145 (99%)	143 (99%)	1 (1%)	84	83
2	B	75/78 (96%)	75 (100%)	0	100	100
2	F	75/78 (96%)	74 (99%)	1 (1%)	69	64
2	H	76/78 (97%)	76 (100%)	0	100	100
2	M	75/78 (96%)	75 (100%)	0	100	100
All	All	877/892 (98%)	868 (99%)	9 (1%)	76	73

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	67	MET
1	E	79	LEU
1	E	105	ASP
2	F	3	SER
1	A	6	LEU
1	A	79	LEU
1	G	31	GLU
1	G	52	LEU
1	L	135	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	27	HIS
1	E	94	HIS
2	F	47	GLN
2	F	81	GLN
1	A	27	HIS
1	G	27	HIS
1	G	95	GLN
2	H	92	ASN
1	L	131	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GSP	L	202	3	26,34,34	1.31	4 (15%)	28,54,54	1.89	6 (21%)
4	GSP	A	202	3	26,34,34	1.23	5 (19%)	28,54,54	1.89	7 (25%)
4	GSP	G	202	3	26,34,34	1.39	5 (19%)	28,54,54	2.05	8 (28%)
4	GSP	E	202	3	26,34,34	1.18	2 (7%)	28,54,54	1.99	6 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GSP	L	202	3	-	2/17/38/38	0/3/3/3
4	GSP	A	202	3	-	2/17/38/38	0/3/3/3
4	GSP	G	202	3	-	1/17/38/38	0/3/3/3
4	GSP	E	202	3	-	2/17/38/38	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	202	GSP	C6-C5	4.23	1.48	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	L	202	GSP	C6-C5	3.91	1.48	1.41
4	E	202	GSP	C6-C5	3.55	1.47	1.41
4	A	202	GSP	C6-C5	3.42	1.47	1.41
4	G	202	GSP	PG-S1G	2.75	1.96	1.90
4	L	202	GSP	PG-S1G	2.63	1.96	1.90
4	E	202	GSP	PG-S1G	2.40	1.95	1.90
4	A	202	GSP	PG-S1G	2.37	1.95	1.90
4	G	202	GSP	C5-C4	2.34	1.47	1.40
4	L	202	GSP	C5-C4	2.28	1.47	1.40
4	A	202	GSP	C5-C4	2.21	1.46	1.40
4	L	202	GSP	O4'-C1'	2.12	1.44	1.41
4	G	202	GSP	C2'-C1'	-2.05	1.50	1.53
4	A	202	GSP	O4'-C1'	2.04	1.43	1.41
4	A	202	GSP	C2'-C1'	-2.03	1.50	1.53
4	G	202	GSP	O4'-C1'	2.01	1.43	1.41

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	202	GSP	C6-N1-C2	5.04	123.94	115.93
4	E	202	GSP	C5-C6-N1	-4.87	116.78	123.43
4	G	202	GSP	C2-N3-C4	4.57	120.58	115.36
4	L	202	GSP	C6-N1-C2	4.57	123.19	115.93
4	E	202	GSP	C6-C5-C4	-4.43	116.56	120.80
4	L	202	GSP	C6-C5-C4	-4.42	116.58	120.80
4	A	202	GSP	C6-C5-C4	-4.38	116.62	120.80
4	A	202	GSP	C6-N1-C2	4.36	122.86	115.93
4	G	202	GSP	C6-C5-C4	-4.33	116.66	120.80
4	A	202	GSP	C5-C6-N1	-4.15	117.76	123.43
4	G	202	GSP	C6-N1-C2	4.11	122.46	115.93
4	G	202	GSP	C5-C6-N1	-3.98	117.98	123.43
4	L	202	GSP	C5-C6-N1	-3.98	117.99	123.43
4	G	202	GSP	N3-C2-N1	-3.45	122.63	127.22
4	A	202	GSP	C2-N3-C4	3.40	119.24	115.36
4	E	202	GSP	N3-C2-N1	-3.33	122.78	127.22
4	L	202	GSP	N3-C2-N1	-3.30	122.82	127.22
4	L	202	GSP	C2-N3-C4	3.26	119.08	115.36
4	A	202	GSP	N3-C2-N1	-3.08	123.11	127.22
4	E	202	GSP	C2-N3-C4	2.84	118.60	115.36
4	G	202	GSP	C4-C5-N7	-2.55	106.74	109.40
4	A	202	GSP	C1'-N9-C4	-2.35	122.52	126.64
4	E	202	GSP	C1'-N9-C4	-2.35	122.52	126.64

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	202	GSP	C4-C5-N7	-2.27	107.03	109.40
4	L	202	GSP	C1'-N9-C4	-2.21	122.75	126.64
4	G	202	GSP	C1'-N9-C4	-2.16	122.85	126.64
4	G	202	GSP	C3'-C2'-C1'	2.12	104.17	100.98

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	202	GSP	PA-O3A-PB-O2B
4	A	202	GSP	PA-O3A-PB-O2B
4	G	202	GSP	PA-O3A-PB-O2B
4	L	202	GSP	PA-O3A-PB-O1B
4	L	202	GSP	PA-O3A-PB-O2B
4	E	202	GSP	PA-O3A-PB-O1B
4	A	202	GSP	PA-O3A-PB-O1B


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	202	GSP	1	0
4	G	202	GSP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

## Ligand GSP L 202



## Ligand GSP A 202





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	166/167 (99%)	-0.06	1 (0%) 89 90	19, 31, 56, 69	0
1	E	166/167 (99%)	-0.27	2 (1%) 79 80	17, 27, 50, 66	0
1	G	166/167 (99%)	0.06	6 (3%) 42 45	25, 45, 70, 89	0
1	L	166/167 (99%)	-0.30	0 100 100	17, 29, 50, 66	0
2	B	93/97 (95%)	-0.16	3 (3%) 47 50	21, 37, 61, 69	0
2	F	93/97 (95%)	-0.36	2 (2%) 62 63	18, 28, 51, 61	0
2	H	94/97 (96%)	-0.17	2 (2%) 63 65	22, 34, 55, 82	0
2	M	93/97 (95%)	-0.30	1 (1%) 80 82	18, 33, 57, 62	0
All	All	1037/1056 (98%)	-0.18	17 (1%) 72 73	17, 32, 60, 89	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	2	SER	5.2
1	G	166	HIS	4.1
1	A	166	HIS	3.5
2	B	5	PRO	3.5
1	G	121	ALA	3.4
2	H	3	SER	3.3
2	F	3	SER	3.2
2	B	3	SER	2.9
1	E	166	HIS	2.7
2	B	4	VAL	2.3
2	M	5	PRO	2.2
1	G	122	ALA	2.2
1	G	52	LEU	2.2
1	G	1	MET	2.1
2	F	4	VAL	2.1
1	G	107	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	E	121	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	G	201	1/1	0.98	0.05	24,24,24,24	0
4	GSP	G	202	32/32	0.98	0.07	19,30,44,47	0
3	MG	E	201	1/1	0.99	0.07	16,16,16,16	0
3	MG	L	201	1/1	0.99	0.09	16,16,16,16	0
4	GSP	E	202	32/32	0.99	0.09	13,18,21,22	0
4	GSP	A	202	32/32	0.99	0.10	17,22,26,27	0
3	MG	A	201	1/1	0.99	0.10	20,20,20,20	0
4	GSP	L	202	32/32	0.99	0.09	15,21,25,26	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around GSP G 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around GSP E 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around GSP A 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GSP L 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.